

ÜBERSICHT

1 Zur Effektivität von Aphasietherapie

H. Grötzbach

ORIGINALARBEIT

6 D-Amphetamin zur Förderung der Rehabilitation nach Schlaganfall

– Wunsch und Wirklichkeit

S. Hesse, T. Treig, M. Sachse, C. Werner

9 Neurogene Blasenfunktionsstörungen bei Multipler Sklerose: Ursachen, Abklärung und neue therapeutische Aspekte durch die Behandlung mit Botulinumtoxin-A

H. Schulte-Baukloh, C. Weiß, J. Schobert, T. Stolze, H. H. Knispel

13 Strukturelle Validität des Short Form 36 Health Survey (SF-36) bei neurologischen Patienten

C. Maurischat, A. Bestmann, U. Hasenbein, M. Bullinger

JOURNAL SCREEN

21 Cortical remapping in amputees and dysmelic patients: A functional

MRI study | NeuroRehabilitation 2003; 18 (4): 299-305

Cognitive impairment in patients with chronic whiplash-associated disorder | NeuroRehabilitation 2003; 18 (4): 307-315

Benefit of inpatient multidisciplinary rehabilitation up to 1 year after stroke | Arch Phys Med Rehabil 2003; 84 (11): 1687-1691

Does neurorehabilitation have a role in relapsing-remitting multiple sclerosis? | J Neurol 2003; 250 (10): 1214-1218

The cognitive ability of an incident cohort of Parkinson's patients in the UK. The CamPaIGN study | Brain 2004; 127 (3): 550-560

Does the representation of time depend on the cerebellum? Effect of cerebellar stroke | Brain 2004; 127 (3): 561-574

Subthalamic nucleus stimulation and dysarthria in Parkinson's disease: a PET study | Brain 2004; 127 (3): 602-615

Triptans in migraine – The risks of stroke, cardiovascular disease, and death in practice | Neurology 2004; 62: 563-568

Structural brain abnormalities in multiple sclerosis patients with major depression | Neurology 2004; 62: 586-590

Long-term effects of temporal lobectomy on intelligence | Neurology 2004; 62: 607-611

