

A. Akyol Erikci, M. Ozyurt, H. Terekeci, A. Ozturk, O. Karabudak, K. Oncu. Oesophageal aspergillosis in a case of acute lymphoblastic leukaemia successfully treated with caspofungin alone due to liposomal amphotericin B induced severe hepatotoxicity	84	S. Verghese, T. Chellamma, K. M. Cherian. Osteomyelitis of the rib caused by <i>Aspergillus flavus</i> following cardiac surgery	91
T.-A. Vyzantiadis, A. Kioumi, E. Papadakis, M. Braimi, E. Dermitzakis, I. Tsitouridis, A. Antoniadis. Rhino-cerebral zygomycosis resistant to antimycotic treatment: a case report	87	Book Review	94
		Corrigenda	95
		Congress Calendar	97

CAPTION OF THE COVER ILLUSTRATION. Histopathological analysis of skin from untreated control (a) and guinea pigs treated with terbinafine HCl nail solutions (TNS) (terbinafine HCl 1%, dodecyl-2-N,N-dimethylaminopropionate hydrochloride 5%) (b). Sections were stained with Grocott Methenamine Silver stain. As can be seen in (a), dark-coloured *Trichophyton mentagrophytes* hyphae were present (arrows) in the stratum corneum of the epidermis and (b) shows the structure of intact skin from TNS-treated animals. See M. A. Ghannoum *et al.*, Determination of the efficacy of terbinafine hydrochloride nail solution in the topical treatment of dermatophytosis in a guinea pig model, pp. 35-43 in this issue.
