

Contents

■ CNS-active Drugs · Hypnotics · Psychotropics · Sedatives

Cánovas M, Torres F, Domenech G, Cebrecos J, Pelagio P, Martínez G, Polonio F, Cabré F E 489

Bioequivalence evaluation of two oral formulations of quetiapine fumarate in healthy volunteers

■ Antiemetics · Gastrointestinal Drugs · Urologic Drugs

Huang M, Tian Y, Zhang Z, Zhang H, Lin H E 494

Determination of propiverine hydrochloride in human plasma by high performance liquid chromatography-tandem mass spectrometry: application to the pharmacokinetic study of a sustained release formulation

Cai W, Wang ZT, Li J, Hu JJ, Zhong J E 502

Bioequivalence study of two esomeprazole enteric coated formulations in healthy Chinese volunteers

■ Analgesics · Anti-inflammatories · Antiphlogistics · Antirheumatic Drugs

Venkatesh S, Durga KD, Padmavathi Y, Reddy BM, Mullangi R E 506

Influence of piperine on ibuprofen induced antinociception and its pharmacokinetics

Ruikar AD, Misar AV, Jadhav RB, Rojatkari SR, Mujumdar AM, Puranik VG, Deshpande NR E 510

Sesquiterpene lactone, a potent drug molecule from *Artemisia pallens* Wall with anti-inflammatory activity

■ Sex Hormones and Inhibitors

Kobayashi H, Gotanda K, Shibata Y, Watanabe JJ, Nakano Y, Shinbo A, Suzuki K E 515

Suppressive effects of the antiandrogen agent, chlormadinone acetate and the 5 α -reductase inhibitor, dutasteride on prostate weight and intraprostatic androgen levels in rats

■ Antibiotics · Antimycotics · Antiparasitics · Antiviral Drugs · Chemotherapeutics · Cytostatics

Bashandy MS, Al-Said MS, Al-Qasoumi SI, Ghorab MM E 521

Design and synthesis of some novel hydrazide, 1,2-dihydropyridine, chromene derivatives carrying biologically active sulfone moieties with potential anticancer activity

Al-Said MS, Bashandy MS, Ghorab MM E 527

Novel quinolines bearing a biologically active trimethoxyphenyl moiety as a new class of anti-tumor agents

Wahajuddin, Singh SP, Patel K, Pradhan T, Siddiqui HH, Singh SK E 532

Prediction of human absorption of a trioxane antimalarial drug (CDRI 99/411) using an in-house validated *in situ* single-pass intestinal perfusion model

Sanki UK, Mandal BK, Chandrakala V E 538

Comparative pharmacokinetics study of two different clindamycin capsule formulations: a randomized, two-period, two-sequence, two-way crossover clinical trial in healthy volunteers