

**CONTENTS**

Volume 159 Issue 10 2011

**SPECIAL ISSUE SECTION: Nitrogen Deposition, Critical Loads and Biodiversity**

**Introduction**

- 2211 Special issue on nitrogen deposition, critical loads, and biodiversity**  
C.L. Goodale, N.B. Dise, M.A. Sutton

**Special Section Papers**

- 2214 Atmospheric nitrogen deposition: Revisiting the question of the importance of the organic component**  
S.E. Cornell

*Routinely including the organic component of atmospheric deposition (known to be around 25–35% worldwide) would make the understanding and prediction of nitrogen biogeochemistry more robust. This paper makes a preliminary global synthesis based on literature reports.*

- 2223 Nitrogen critical loads using biodiversity-related critical limits**  
M. Posch, J. Aherne, J.-P. Hettelingh

*Biodiversity-related critical limits lead to optimal nitrogen and sulphur deposition envelopes for plant species or species compositions.*

- 2228 Effects of experimental nitrogen additions on plant diversity in tropical forests of contrasting disturbance regimes in southern china**  
X. Lu, J. Mo, F.S. Gilliam, G. Yu, W. Zhang, Y. Fang, J. Huang

*Research in disturbed forests of southeastern China demonstrates that land-use history can substantially alter effects of excess nitrogen deposition on plant diversity of tropical forest ecosystems.*

- 2236 Diversity of shrub tree layer, leaf litter decomposition and N release in a Brazilian Cerrado under N, P and N plus P additions**  
T.K.B. Jacobson, M.M.da.C. Bustamante, A.R. Kozovits

*Shrub-tree diversity and functioning of Brazilian savanna are affected by increasing nutrient availability.*

- 2243 The impact of nitrogen deposition on acid grasslands in the Atlantic region of Europe**  
C.J. Stevens, C. Duprè, E. Dorland, C. Gaudnik, D.J.G. Gowing, A. Bleeker, M. Diekmann, D. Alard, R. Bobbink, D. Fowler, E. Corcket, J.O. Mountford, V. Vandvik, P.A. Aarrestad, S. Muller, N.B. Dise

*Atmospheric nitrogen deposition affects soils, plant-tissue chemistry and plant species composition in acid grasslands in the Atlantic biogeographic region of Europe.*

- 2251 Nitrogen deposition and its ecological impact in China: An overview**  
X. Liu, L. Duan, J. Mo, E. Du, J. Shen, X. Lu, Y. Zhang, X. Zhou, C. He, F. Zhang

*This paper reviews current knowledge on nitrogen deposition and its effects across China*

- 2265 Nitrogen deposition effects on Mediterranean-type ecosystems: An ecological assessment**  
R. Ochoa-Hueso, E.B. Allen, C. Branquinho, C. Cruz, T. Dias, M.E. Fenn, E. Manrique, M.E. Pérez-Corona, L.J. Sheppard, W.D. Stock

*Nitrogen deposition threatens the Mediterranean regions of the world.*

- 2280 N deposition as a threat to the World's protected areas under the Convention on Biological Diversity**  
A. Bleeker, W.K. Hicks, F. Dentener, J. Galloway, J.W. Erisman

*Significant areas of the UNEP Protected Areas Programme under the CBD receive high N deposition rates that are likely to increase in the future, especially in Asia, and may pose a significant threat to biodiversity.*

# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from outside back cover

- 2289 Modelling the impact of nitrogen deposition, climate change and nutrient limitations on tree carbon sequestration in Europe for the period 1900–2050**  
W. de Vries, M. Posch

*A modelling exercise indicates that nitrogen deposition was the main factor that enhanced tree carbon sequestration in Europe in the past, while climate change will do so in the future*

## Recognition of Frequent Reviewers

- 2300 Recognition of frequent reviewers 2007–2010**

### Reviews

- 2301 Regional dynamics of persistent organic pollutants (POPs) in the Pearl River Delta, China: Implications and perspectives**  
K. Zhang, B.-Z. Zhang, S.-M. Li, E.Y. Zeng

*Regional dynamics of p,p'-DDT and BDE-209 in the Pearl River Delta (South China) are examined and used to predict temporal variances.*

- 2310 Sources of ambient volatile organic compounds and their contributions to photochemical ozone formation at a site in the Pearl River Delta, southern China**  
Z.H. Ling, H. Guo, H.R. Cheng, Y.F. Yu

*Solvent usage, diesel vehicular emissions and biomass/biofuel burning were the major contributors to the photochemical O<sub>3</sub> formation in the PRD region.*

- 2320 *In vitro* and *in vivo* approaches for the measurement of oral bioavailability of lead (Pb) in contaminated soils: A review**  
M.H. Zia, E.E. Codling, K.G. Scheckel, R.L. Chaney

*Improved risk evaluation and recommendations for Pb contaminated soils should be based on bioavailability-correlated bioaccessible soil Pb rather than total soil Pb.*

- 2328 Fate and effects of anthropogenic chemicals in mangrove ecosystems: A review**  
M. Lewis, R. Pryor, L. Wilking

*Chemical risk assessments and resource management are restricted by the limited chemical fate and effects database for mangroves.*

### Commentary

- 2347 Trophically available metal – A variable feast**  
P.S. Rainbow, S.N. Luoma, W.-X. Wang

*Trophically Available Metal (TAM), the component of accumulated metal in food that is taken up by a feeding animal, varies with food type and consumer.*

### Rapid Communication

- 2350 Exposure to moderate concentrations of tropospheric ozone impairs tree stomatal response to carbon dioxide**  
G. Onandia, A.-K. Olsson, S. Barth, J.S. King, J. Uddling

*Exposure to moderate concentrations of tropospheric ozone impairs stomatal CO<sub>2</sub> responsiveness of birch in the Aspen FACE experiment.*

### New Initiatives

- 2355 Sorption isotherms of brominated diphenyl ethers on natural soils with different organic carbon fractions**  
W. Liu, W. Li, B. Xing, J. Chen, S. Tao

*Both linear distribution and nonlinear Freundlich model could well describe the sorption isotherms of PBDEs on natural soils.*

### Regular Papers

- 2359 Assessing the diurnal variability of pharmaceutical and personal care products in a full-scale activated sludge plant**  
R. Salgado, R. Marques, J.P. Noronha, J.T. Mexia, G. Carvalho, A. Oehmen, M.A.M. Reis

*The diurnal variations of pharmaceuticals and musks were studied in an activated sludge plant, where the loadings of the musks were more repeatable than the pharmaceuticals.*

- 2368 Species- and age-related variation in metal exposure and accumulation of two passerine bird species**  
Å.M.M. Berglund, M.J. Koivula, T. Eeva

*Accumulation of metals in liver of two insectivorous passerines reflects inter-specific differences in diet, absorption rate and physiological requirements.*

- 2375 Sorption-desorption behavior of polybrominated diphenyl ethers in soils**  
Y. Olshansky, T. Polubesova, W. Vetter, B. Chefetz

*Pronounced desorption hysteresis was observed for BDE-15 in natural soils.*

- 2380 Wastewater treatment plants and landfills emit volatile methyl siloxanes (VMSs) to the atmosphere: Investigations using a new passive air sampler**

Y. Cheng, M. Shoeb, L. Ahrens, T. Harner, J. Ma

*Wastewater treatment plants and landfills are shown to emit large quantities of chemicals used in personal care products (volatile methyl siloxanes) to the atmosphere.*

- 2387 Global DNA methylation in earthworms: A candidate biomarker of epigenetic risks related to the presence of metals/metalloids in terrestrial environments**

M.M. Santoyo, C.R. Flores, A.L. Torres, K. Wrobel, K. Wrobel

*Biomethylation of As, Hg in earthworms versus DNA methylation – a candidate biomarker of epigenetic risks related to the presence of metals/metalloids in soil.*

# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from inside back cover

- 2393 Mobile passive samplers: Concept for a novel mode of exposure**  
I.J. Allan, H.C. Nilson, I. Tjensvoll, C. Bradshaw, K. Næs  
*The first application of a novel mode of exposure for passive sampling devices in water is presented.*
- 2398 Arsenite toxicity and uptake rate of rice (*Oryza sativa* L.) in vivo**  
H. Hoffmann, M.K. Schenk  
*Uptake of arsenite, water, and nutrients by rice seedlings was impaired by arsenite concentrations higher than the toxicity threshold of 2.4 µM.*
- 2405 The effect of air pollution and other environmental stressors on leaf fluctuating asymmetry and specific leaf area of *Salix alba* L**  
T. Wuytack, K. Wuyts, S. Van Dongen, L. Baeten, F. Kardel, K. Verheyen, R. Samson  
*Specific leaf area of *S. alba* increased with increasing shade and, in less extent, with increasing NO<sub>x</sub> and decreasing O<sub>3</sub> concentration, while leaf asymmetry did not respond to air pollution*
- 2412 Concentration of trace elements in feathers of three Antarctic penguins: Geographical and interspecific differences**  
S. Jerez, M. Motas, M.J. Palacios, F. Valera, J.J. Cuervo, A. Barbosa  
*Trace element levels in feathers of three penguin species from the Antarctic Peninsula indicate the presence of pollution in certain locations.*
- 2420 Within season and carry-over effects following exposure of grassland species mixtures to increasing background ozone**  
F. Hayes, G. Mills, H. Harmens, K. Wyness  
*Increased background ozone enhanced leaf senescence and caused carry-over effects including reduced winter die-back and an acceleration in flowering the next year.*
- 2427 Distribution, speciation and availability of antimony (Sb) in soils and terrestrial plants from an active Sb mining area**  
G. Okkenhaug, Y.-G. Zhu, L. Luo, M. Lei, X. Li, J. Mulder  
*Antimony (Sb) in soils from an active Sb mining area is highly bioavailable and controlled by the solubility of calcium antimonate.*
- 2435 Inorganic arsenic and trace elements in Ghanaian grain staples**  
E.E. Adomako, P.N. Williams, C. Deacon, A.A. Meharg  
*Ghanaian rice is generally low in potentially toxic trace elements and is of higher nutritional quality than rice imported into the country.*
- 2443 Characterization of PBDEs in soils and vegetations near an e-waste recycling site in South China**  
Y. Wang, C. Luo, J. Li, H. Yin, X. Li, G. Zhang  
*E-waste recycling sites are heavily contaminated with PBDEs, which may pose a potential health risk to the local ecosystem and inhabitants.*
- 2449 Reduction in uptake by rice and soybean of aromatic arsenicals from diphenylarsinic acid contaminated soil amended with activated charcoal**  
T. Arao, Y. Maejima, K. Baba  
*Activated charcoal amendment to soil contaminated with diphenylarsinic acid reduced aromatic arsenicals in rice and soybean.*
- 2454 Seabird feathers as monitors of the levels and persistence of heavy metal pollution after the *Prestige* oil spill**  
R. Moreno, L. Jover, C. Diez, C. Sanpera  
*Monitoring heavy metal in seabird feathers indicated chronic oil pollution in the marine food web for at least three years after the *Prestige* oil spill.*
- 2461 Groundwater quality assessment of one former industrial site in Belgium using a TRIAD-like approach**  
S. Crévecoeur, V. Debacker, C. Joaquim-Justo, S. Gobert, M.-L. Scippo, W. Dejonghe, P. Martin, J.-P. Thomé  
*This study presents the first application of the TRIAD approach on groundwater system. None of the TRIAD components (chemistry, physico-chemistry and ecotoxicity) could reliably predict the other one.*
- 2467 Long-term effects of clear-cutting and selective cutting on soil methane fluxes in a temperate spruce forest in southern Germany**  
X. Wu, N. Brüggemann, R. Gasche, H. Papen, G. Willibald, K. Butterbach-Bahl  
*Forest management affects the soil CH<sub>4</sub> sink strength, with clear-cutting reducing uptake rates for at least eight years.*
- 2476 Binational school-based monitoring of traffic-related air pollutants in El Paso, Texas (USA) and Ciudad Juárez, Chihuahua (México)**  
A.U. Raysoni, J.A. Sarnat, S.E. Sarnat, J.H. Garcia, F. Holguin, S.F. Luévano, W.-W. Li  
*Spatial variability in traffic-mediated pollutant concentrations can exist at the intra-urban level and ambient monitoring sites may not accurately represent these concentration gradients.*
- 2487 Reporting and evaluation criteria as means towards a transparent use of ecotoxicity data for environmental risk assessment of pharmaceuticals**  
M. Ågerstrand, A. Küster, J. Bachmann, M. Breitholtz, I. Ebert, B. Rechenberg, C. Rudén  
*The need for reporting and evaluation criteria towards a transparent and reliable use of ecotoxicity data.*

**CONTENTS** – Continued from page VII

- 2493 Enhanced nitrogen deposition exacerbates the negative effect of increasing background ozone in *Dactylis glomerata*, but not *Ranunculus acris***  
K. Wyness, G. Mills, L. Jones, J.D. Barnes, D.L. Jones  
*Synergistic effects of elevated O<sub>3</sub> and N were observed in below ground C-partitioning in the grass Dactylis glomerata, but not in the forb Ranunculus acris.*
- 2500 Mercury accumulation in marine bivalves: Influences of biodynamics and feeding niche**  
K. Pan, W.-X. Wang  
*The species-specific body concentrations of MeHg and THg in bivalves reflect the complicated interaction between the biodynamics of Hg(II) and MeHg and the different food sources.*
- 2507 Simultaneous adsorption and degradation of  $\gamma$ -HCH by nZVI/Cu bimetallic nanoparticles with activated carbon support**  
C. Chang, F. Lian, L. Zhu  
*Impregnation of Cu amended iron on AC enhances the removal efficiency of  $\gamma$ -HCH and reduces the concentrations of its intermediates in aqueous solution.*
- 2515 Environmental impact of sunscreen nanomaterials: Ecotoxicity and genotoxicity of altered TiO<sub>2</sub> nanocomposites on *Vicia faba***  
A.-S. Foltête, J.-F. Masfaraud, E. Bigorgne, J. Nahmani, P. Chaurand, C. Botta, J. Labille, J. Rose, J.-F. Férard, S. Cotelle  
*Artificially aged TiO<sub>2</sub>-containing sunscreen nanocomposites caused few damages to *Vicia faba* in 48 h exposure, but particles clogged on roots and Ti entered the root tissues.*
- 2523 Blood dynamics of mercury and selenium in northern elephant seals during the lactation period**  
S. Habran, C. Debier, D.E. Crocker, D.S. Houser, K. Das  
*Blood levels of total Hg and Se are modified during the 4-week lactating period in northern elephant seals.*
- 2530 Effects of organic matter and ageing on the bioaccessibility of arsenic**  
L. Meunier, I. Koch, K.J. Reimer  
*Adding organic carbon may increase arsenic bioaccessibility, especially in samples originally containing >30% bioaccessible arsenic in organic carbon-rich soils (>25%).*
- 2537 Can arbuscular mycorrhizal fungi improve grain yield, As uptake and tolerance of rice grown under aerobic conditions?**  
H. Li, Z.H. Ye, W.F. Chan, X.W. Chen, F.Y. Wu, S.C. Wu, M.H. Wong  
*Different rice/AMF combinations had very different effects on arsenic and phosphorus uptake.*
- 2546 Oxidation of bisphenol F (BPF) by manganese dioxide**  
Z. Lu, K. Lin, J. Gan  
*The commonly occurring MnO<sub>2</sub> shows a high reactivity toward bisphenol F, which may account for its natural attenuation and suggest a beneficial use of MnO<sub>2</sub> for managed removal of bisphenol F.*
- 2552 Organic persistent toxic substances in soils, waters and sediments along an altitudinal gradient at Mt. Sagarmatha, Himalayas, Nepal**  
L. Guzzella, G. Poma, A. De Paolis, C. Roscioli, G. Viviano  
*Organic PTSs in environmental matrices in remote regions of the Himalayan ridge.*
- 2565 Distributions of polyhalogenated compounds in Hudson River (New York, USA) fish in relation to human uses along the river**  
L.C. Skinner  
*Residues of polyhalogenated compounds in resident and migratory fish from the Hudson River are compared with human uses of the compounds in the river basin.*
- 2575 Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze River basin**  
Y. Yi, Z. Yang, S. Zhang  
*Possible ecological risk of sediment and slight non-cancer health risk of eating fish were found in the middle and lower reaches of the Yangtze River.*
- 2586 Long-term patterns in soil acidification due to pollution in forests of the Eastern Sudetes Mountains**  
R. Hédli, P. Petřík, K. Boublík  
*Strong acidification decreasing with altitude was observed in forest soils resurveyed after more than half a century.*
- 2594 Sorption of simazine to corn straw biochars prepared at different pyrolytic temperatures**  
G. Zhang, Q. Zhang, K. Sun, X. Liu, W. Zheng, Y. Zhao  
*The corn straw biochar prepared at higher temperature with stronger hydrophobicity, more aromatic C and larger surface area had higher sorption capacity for simazine.*
- 2602 Impacts of atmospheric pollution on the plant communities of British acid grasslands**  
R.J. Payne, C.J. Stevens, N.B. Dise, D.J. Gowing, M.G. Pilkington, G.K. Phoenix, B.A. Emmett, M.R. Ashmore  
*Ozone exposure and nitrogen deposition have distinct but additive impacts on the plant communities of British acid grasslands.*
- 2609 Monitoring plastic ingestion by the northern fulmar *Fulmarus glacialis* in the North Sea**  
J.A. van Franeker, C. Blaize, J. Danielsen, K. Fairclough, J. Gollan, N. Guse, P.-L. Hansen, M. Heubeck, J.-K. Jensen, G. Le Guillou, B. Olsen, K.-O. Olsen, J. Pedersen, E.W.M. Stienen, D.M. Turner  
*Trends and patterns in the abundance of plastics in northern fulmar stomachs show that seabirds can be a valuable tool for monitoring marine litter.*

- 2616 Adsorption of sulfamethoxazole on functionalized carbon nanotubes as affected by cations and anions**  
D. Zhang, B. Pan, M. Wu, B. Wang, H. Zhang, H. Peng, D. Wu, P. Ning  
*The balance between decreasing and increasing roles determines the apparent sulfamethoxazole adsorption on carbon nanotubes depending on environmental conditions.*
- 2622 The impact of heavy metal pollution gradients in sediments on benthic macrofauna at population and community levels**  
J. Ryu, J.S. Khim, S.-G. Kang, D. Kang, C.-h. Lee, C.-h. Koh  
*Community and population level response to the polluted environment of the harbor reflected an integration effect, together with biological interactions.*
- 2630 In situ evaluation of cadmium biomarkers in green algae**  
D.F. Simon, T.A. Davis, M.-L. Tercier-Waeber, R. England, K.J. Wilkinson  
*Algal transcription levels of several biomarkers were studied in two natural waters in situ.*
- 2637 Arsenic distribution in soils and plants of an arsenic impacted former mining area**  
V. Otones, E. Álvarez-Ayuso, A. García-Sánchez, I. Santa Regina, A. Murciego  
*The plants *Salix atrocinerea*, *Genista scorpius* and *Scirpus holoschoenus* are suitable for revegetation or phytostabilisation approaches of As-polluted soils.*
- 2648 Developmental toxicity of two common corn pesticides to the endangered southern bell frog (*Litoria raniformis*)**  
C.B. Choung, R.V. Hyne, R.M. Mann, M.M. Stevens, G.C. Hose  
*A breakdown product of the insecticide terbufos retards development and delays metamorphosis of southern bell frog tadpoles.*
- 2656 Seasonal and spatial variability of polychlorinated biphenyls (PCBs) in vegetation and cow milk from a high altitude pasture in the Italian Alps**  
L. Tato, P. Tremolada, C. Ballabio, N. Guazzoni, M. Parolini, M. Caccianiga, A. Binelli  
*In a mountain pasture, PCBs concentrations in vegetation were found to be related to emission flux from soil and to the mean temperature of the site, and a direct transfer to milk was also observed.*
- 2665 Sediment-associated aliphatic and aromatic hydrocarbons in coastal British Columbia, Canada: Concentrations, composition, and associated risks to protected sea otters**  
K.A. Harris, M.B. Yunker, N. Dangerfield, P.S. Ross  
*Anthropogenically-derived hydrocarbons in coastal sediments in British Columbia may pose a risk to protected sea otters.*
- 2675 Hydrocarbon degradation, plant colonization and gene expression of alkane degradation genes by endophytic *Enterobacter ludwigii* strains**  
S. Yousaf, M. Afzal, T.G. Reichenauer, C.L. Brady, A. Sessitsch  
*Enterobacter ludwigii strains belonging to the *E. cloacae* complex are able to efficiently degrade alkanes when associated with plants and to promote plant growth.*
- 2684 Field controlled experiments of mercury accumulation in crops from air and soil**  
Z. Niu, X. Zhang, Z. Wang, Z. Ci  
*Capsule Mercury accumulated in the aboveground organs of crop was mainly from the air.*
- 2690 Variations in concentrations and compositions of polycyclic aromatic hydrocarbons (PAHs) in coals related to the coal rank and origin**  
S. Laumann, V. Micić, M.A. Krüge, C. Achten, R.F. Sachsenhofer, J. Schwarzbauer, T. Hofmann  
*The origin of coal has a strong influence on the concentration and composition of indigenous PAHs.*
- 2698 Ecotoxicological assessment of TiO<sub>2</sub> byproducts on the earthworm *Eisenia fetida***  
E. Bigorgne, L. Foucaud, E. Lapiéd, J. Labille, C. Botta, C. Sirguey, J. Falla, J. Rose, E.J. Joner, F. Rodius, J. Nahmani  
*A short time exposure to TiO<sub>2</sub> byproducts can induce sublethal effects on the earthworm, *Eisenia fetida*.*
- 2706 Stable carbon isotope analysis ( $\delta^{13}\text{C}$  values) of polybrominated diphenyl ethers and their UV-transformation products**  
N. Rosenfelder, P. Bendig, W. Vetter  
*Diametrically progressing  $\delta^{13}\text{C}$  values in technical mixtures and UV-transformation products of DE-79 may be useful for source appointment of PBDEs in environmental samples*
- 2713 Chronic toxicity of ZnO nanoparticles, non-nano ZnO and ZnCl<sub>2</sub> to *Folsomia candida* (Collembola) in relation to bioavailability in soil**  
P.L. Kool, M.D. Ortiz, C.A.M. van Gestel  
*ZnO nanoparticle toxicity to springtails in soil can be explained from Zn dissolution but not from particle size.*
- 2720 Reproductive toxicity assessment of surface water of the Tai section of the Yangtze River, China by *in vitro* bioassays coupled with chemical analysis**  
X. Wang, J. Wu, Y. Hao, B. Zhu, W. Shi, G. Hu, X. Han, J.P. Giesy, H. Yu  
*In vitro bioassay responses observed in Yangtze River source water extracts showed great reproductive toxicity, and PAHs were responsible.*
- 2726 Will PM control undermine China's efforts to reduce soil acidification?**  
Y. Zhao, L. Duan, Y. Lei, J. Xing, C.P. Nielsen, J. Hao  
*The control of anthropogenic PM emission in China will increase the risk of soil acidification even with reduced SO<sub>2</sub> emission. Such implication is stronger than that in developed countries.*

- 2733 The effect of mercury on trees and their mycorrhizal fungi**  
S.R. Jean-Philippe, J.A. Franklin, D.S. Buckley, K. Hughes  
*Though evidence suggests that mercury-contaminated soils may reduce tree and fungal populations, there are tolerant species that may remain and survive following contamination.*
- 2740 Terrestrial exposure of oilfield flowline additives diminish soil structural stability and remediative microbial function**  
S.J. George, J. Sherbone, C. Hinz, M. Tibbett  
*Scale inhibitor and biocide oilfield flowline additives interactively affect soil physical and microbial properties*
- 2750 Changes in the chemistry of acidified Adirondack streams from the early 1980s to 2008**  
G.B. Lawrence, H.A. Simonin, B.P. Baldigo, K.M. Roy, S.B. Capone  
*Despite a 50% reduction in atmospheric sulfur deposition since the mid 1980s, chemical recovery of Adirondack streams has been slow, with minimal recovery from 2001 to 2008.*
- 2759 Atmospheric phenanthrene pollution modulates carbon allocation in red clover (*Trifolium pratense* L.)**  
D. Desalme, P. Binet, D. Epron, N. Bernard, D. Gilbert, M.-L. Toussaint, C. Plain, G. Chiapusio  
*Exposure of clover to atmospheric PHE affected not only its growth, but also biomass partitioning and C allocation among its organs.*
- 2766 Reproductive toxicity of bisphenol A and cadmium in *Potamopyrgus antipodarum* and modulation of bisphenol A effects by different test temperature**  
A. Sieratowicz, D. Stange, U. Schulte-Oehlmann, J. Oehlmann  
*A reproduction test with *Potamopyrgus antipodarum* with 2 substances for subsequent pre-validation is presented and bisphenol A effects show a temperature dependency.*
- 2775 Heavy metal pollution exerts reduction/adaptation in the diversity and enzyme expression profile of heterotrophic bacteria in Cochin estuary, India**  
J. Jose, R. Giridhar, A. Anas, P.A. Loka Bharathi, S. Nair  
*Heavy metal pollution exerts pressure on the diversity and enzyme expression profile of estuarine bacteria.*
- 2781 Dispersal networks for enhancing bacterial degradation in heterogeneous environments**  
T. Banitz, L.Y. Wick, I. Fetzer, K. Frank, H. Harms, K. Johst  
*Simulation modelling demonstrates that fungus-mediated bacterial dispersal can considerably improve the bioavailability of organic pollutants under spatially heterogeneous abiotic conditions typical for water-unsaturated soils.*
- 2789 Arsenic and Mn levels in *Isaza* (*Gymnogobius isaza*) during the mass mortality event in Lake Biwa, Japan**  
S.H. Hirata, D. Hayase, A. Eguchi, T. Itai, K. Nomiya, T. Isobe, T. Agusa, T. Ishikawa, M. Kumagai, S. Tanabe  
*As and Mn levels in dead *Isaza* caused by mass mortality.*
- 2797 Substituting missing data in compositional analysis**  
C. Real, J. Ángel Fernández, J.R. Aboal, A. Carballeira  
*A method is presented for the assessment of the possible distortions in multivariate analysis caused by the substitution of missing values.*
- 2801 Inherent mineralization of 2,6-dichlorobenzamide (BAM) in unsaturated zone and aquifers – Effect of initial concentrations and adaptation**  
G.S. Janniche, L. Clausen, H.-J. Albrechtsen  
*BAM mineralization in subsurface and groundwater was demonstrated.*
- 2808 Temporal genetic variability and host sources of *Escherichia coli* associated with fecal pollution from domesticated animals in the shellfish culture environment of Xiangshan Bay, East China Sea**  
L.-L. Fu, J.-B. Shuai, Y. Wang, H.-J. Ma, J.-R. Li  
*The temporal genetic variability and dominant host sources of *E.coli* in fecal contaminated shellfish growing waters of Xiangshan Bay was characterized.*
- 2815 Distribution and bioaccumulation of steroidal and phenolic endocrine disrupting chemicals in wild fish species from Dianchi Lake, China**  
J. Liu, R. Wang, B. Huang, C. Lin, Y. Wang, X. Pan  
*Contaminants of endocrine disrupting chemicals in wild fish.*
- 2823 Ozone affects pollen viability and NAD(P)H oxidase release from *Ambrosia artemisiifolia* pollen**  
S. Pasqualini, E. Tedeschini, G. Frenguelli, N. Wopfner, F. Ferreira, G. D'Amato, L. Ederli  
*This study focuses on the effects of the atmospheric pollutant ozone on ROS content and NAD(P)H oxidase activity of ragweed pollen grains.*
- 2831 Using advanced surface complexation models for modelling soil chemistry under forests: Solling forest, Germany**  
L.T.C. Bonten, J.E. Groenenberg, H. Meesenburg, W. de Vries  
*Surface complexation models with generic parameters make calibration of sorption superfluous in dynamic modelling of deposition impacts on soil chemistry under nature areas.*
- 2840 Floodplain methylmercury biomagnification factor higher than that of the contiguous river (South River, Virginia USA)**  
M.C. Newman, X. Xu, A. Condon, L. Liang  
*Higher methylmercury biomagnification on South River floodplain than the associated river likely explain high mercury in floodplain birds.*

- 2845 <sup>1</sup>H NMR-based metabolomics of time-dependent responses of *Eisenia fetida* to sub-lethal phenanthrene exposure**  
B.P. Lankadurai, D.M. Wolfe, A.J. Simpson, M.J. Simpson  
*<sup>1</sup>H NMR metabolomics is used to determine the relationship between phenanthrene exposure and the metabolic response of the earthworm *E. fetida* over time and also to elucidate the phenanthrene mode of toxicity.*
- 2852 Nitrogen concentrations in mosses indicate the spatial distribution of atmospheric nitrogen deposition in Europe**  
H. Harmens, D.A. Norris, D.M. Cooper, G. Mills, E. Steinnes, E. Kubin, L. Thöni, J.R. Aboal, R. Alber, A. Carballeira, M. Coşkun, L. De Temmerman, M. Frolova, L. González-Miqueo, Z. Jeran, S. Leblond, S. Liiv, B. Maňková, R. Pesch, J. Poikolainen, Å. Rühling, J.M. Santamaría, P. Simonè, W. Schröder, I. Suchara, L. Yurukova, H.G. Zechmeister  
*Mosses as biomonitors of atmospheric nitrogen deposition in Europe.*
- 2861 Accumulation of mercury and methylmercury by mushrooms and earthworms from forest soils**  
S.R. Rieder, I. Brunner, M. Horvat, A. Jacobs, B. Frey  
*Accumulation of Hg and methyl-Hg by mushrooms and earthworms is species- and ecophysiological group dependent.*
- 2870 Indoor metallic pollution related to mining activity in the Bolivian Altiplano**  
F.E. Fontúrbel, E. Barbieri, C. Herbas, F.L. Barbieri, J. Gardon  
*In a typical Andean mining city, the urban indoor pollution with toxic metallic elements is directly related to the closeness of the mining activities.*
- 2876 Comparison of competitive and synergetic adsorption of three phenolic compounds on river sediment**  
P. Gao, Y. Feng, Z. Zhang, J. Liu, N. Ren  
*The coagulation theory can explain the adsorption among phenolic compounds on river sediment.*
- 2882 The fate of the aqueous phase polycyclic aromatic hydrocarbon fraction in a detention pond system**  
K. Neary, T.B. Boving  
*Aqueous-phase PAH concentrations in stormwater are relatively unaffected by treatment in a conventional detention pond system.*
- 2891 Biomarker responses associated with halogenated organic contaminants in northern fulmars (*Fulmarus glacialis*) breeding in the Canadian Arctic**  
B.M. Braune, S. Trudeau, D.A. Jeffrey, M.L. Mallory  
*EROD induction occurs in northern fulmars in the Canadian Arctic at environmentally-relevant concentrations.*
- 2899 Interspecies and spatial trends in polycyclic aromatic hydrocarbons (PAHs) in Atlantic and Mediterranean pelagic seabirds**  
J.L. Roscales, J. González-Solís, P. Calabuig, B. Jiménez  
*PAH levels in Atlantic and Mediterranean pelagic seabirds show specific inter-species patterns related to trophic ecology but a minor influence of their geographic origin.*
- 2906 How seasonality affects the flow of estrogens and their conjugates in one of Japan's most populous catchments**  
V. Kumar, N. Nakada, N. Yamashita, A.C. Johnson, H. Tanaka  
*This manuscript provides a comprehensive survey of estrogen conjugates in a densely populated river catchment.*
- 2913 Occurrence and distribution of antibiotics in coastal water of the Bohai Bay, China: Impacts of river discharge and aquaculture activities**  
S. Zou, W. Xu, R. Zhang, J. Tang, Y. Chen, G. Zhang  
*River discharge and aquaculture were suggested to be important sources for antibiotics occurred in the coastal water of the Bohai Bay, North China.*
- 2921 Toxicity of dispersant application: Biomarkers responses in gills of juvenile golden grey mullet (*Liza aurata*)**  
T. Milinkovitch, J. Godefroy, M. Théron, H. Thomas-Guyon  
*When the meteorological conditions induce the dispersion of the oil slick (e.g. wave), the application of dispersant does not increase the toxicity of petroleum.*
- 2929 Screening of pharmaceuticals and hormones at the regional scale, in surface and groundwaters intended to human consumption**  
E. Vulliet, C. Cren-Olivé  
*52 pharmaceuticals and hormones investigated in 71 surface waters and 70 groundwaters intended to human consumption.*
- 2935 Vertical emission profiles for Europe based on plume rise calculations**  
J. Bieser, A. Aulinger, V. Matthias, M. Quante, H.A.C. Denier van der Gon  
*Vertical emission profiles that take into account the spatial and temporal variability of effective emission heights improve the usability of European point-source emissions in CTMs.*
- 2947 Spectral estimation of global levels of atmospheric pollutants**  
J. Fernández-Macho  
*A structural spatial model is proposed for the estimation of underlying smooth levels of atmospheric pollutants by spectral methods.*
- 2954 Modeling vehicle emissions in different types of Chinese cities: Importance of vehicle fleet and local features**  
H. Huo, Q. Zhang, K. He, Z. Yao, X. Wang, B. Zheng, D.G. Streets, Q. Wang, Y. Ding  
*Vehicle emission characteristics of Chinese cities are remarkably different, and local features need to be taken into account in vehicle emission studies and control strategy.*
- 2961 Modeling redistribution of  $\alpha$ -HCH in Chinese soil induced by environment factors**  
C. Tian, L. Liu, J. Ma, J. Tang, Y.-F. Li  
*This study provides the first comprehensive overview to redistribution of a toxic chemical incurred by long-term variation of environmental factors across China.*

- 2968 The application of zero-water discharge system in treating diffuse village wastewater and its benefits in community afforestation**  
Y. Wu, L. Xia, Z. Hu, S. Liu, H. Liu, B. Nath, N. Zhang, L. Yang  
*A zero-water discharge system has proven highly-effective for removing COD, SS, TP, and TN from diffuse village wastewater and benefiting the community afforestation.*
- 2974 Comparison of solid and liquid-phase bioassays using ecoscores to assess contaminated soils**  
C. Lors, J.-F. Ponge, M. Martínez Aldaya, D. Damidot  
*Aqueous and solid phases of contaminated soils give similar results in terms of toxicity but are complementary for the evaluation of environmental hazards by ecoscores.*
- 2982 Particle-bound Dechlorane Plus and polybrominated diphenyl ethers in ambient air around Shanghai, China**  
Z. Yu, R. Liao, H. Li, L. Mo, X. Zeng, G. Sheng, J. Fu  
*The moderate PBDEs levels with deca-BDE being the most important contributor and low DP levels in ambient air around Shanghai reflected their different usage in this area.*
- 2989 Spatio-temporal variations of nitrogen in an agricultural watershed in eastern China: Catchment export, stream attenuation and discharge**  
D. Chen, J. Lu, Y. Shen, D. Gong, O. Deng  
*High flow rate brings large catchment nitrogen export but low in-stream attenuation efficiency, and then enhances stream nitrogen discharge.*
- 2996 Modelling arsenic toxicity in wheat: Simultaneous application of diffusive gradients in thin films to arsenic and phosphorus in soil**  
O. Mojsilovic, R.G. McLaren, L.M. Condon  
*Diffusive gradient in thin films (DGT) measure of soil arsenic-to-phosphorus ratio provides a promising indicator of arsenic toxicity in wheat.*
- 3003 Synergistic toxic effect of nano- $Al_2O_3$  and As(V) on *Ceriodaphnia dubia***  
D. Wang, J. Hu, B.E. Forthaus, J. Wang  
*Nano- $Al_2O_3$  could accumulate background As(V) and enhance As(V) toxicity on *C. dubia* through the uptake of As(V)-loaded nano- $Al_2O_3$  particles.*
- 3009 Functional redundancy and food web functioning in linuron-exposed ecosystems**  
F. De Laender, P.J. Van den Brink, C.R. Janssen  
*Food web modelling reveals functional redundancy of the planktonic community in microcosms treated with linuron.*
- 3018 Phytostabilization of semiarid soils residually contaminated with trace elements using by-products: Sustainability and risks**  
A. Pérez-de-Mora, P. Madejón, P. Burgos, F. Cabrera, N.W. Lepp, E. Madejón  
*Phytostabilization using by-products as amendments is a suitable approach for long-term immobilization of various trace elements in semiarid contaminated soils.*
- 3028 Trace element uptake by *Eleocharis equisetina* (spike rush) in an abandoned acid mine tailings pond, northeastern Australia: Implications for land and water reclamation in tropical regions**  
B.G. Lottermoser, P.M. Ashley  
*The metal-excluding aquatic macrophyte *Eleocharis equisetina* is of use in land and water remediation in tropical regions.*
- 3036 Patterns of insect communities along a stress gradient following decommissioning of a Cu–Ni smelter**  
J. Babin-Fenske, M. Anand  
*Our study finds both unexpected and expected responses of insect communities to a landscape gradient of past pollution suggesting the emergence of novel ecosystems.*
- 3044 Polybrominated diphenyl ethers (PBDEs) in feathers of colonial water-bird species from Pakistan**  
R.N. Malik, C. Moeckel, K.C. Jones, D. Hughes  
*Feathers are used as a biomonitoring tool for PBDEs contamination.*
- 3051 Pyrethroid insecticides in urban salmon streams of the Pacific Northwest**  
D.P. Weston, A.M. Asbell, S.A. Hecht, N.L. Scholz, M.J. Lydy  
*Pyrethroid insecticides are present in sediments of urban creeks of Oregon and Washington, though less commonly than in studies elsewhere in the U.S.*
- 3057 Assessment of heavy metal pollution from a Fe-smelting plant in urban river sediments using environmental magnetic and geochemical methods**  
C. Zhang, Q. Qiao, J.D.A. Piper, B. Huang  
*Heavy metal (HM) contamination of river sediments from industrial input by surface water transport and deposition can be detected by using magnetic methods providing a convenient assessment of HM pollution in industrialized cities.*
- 3071 The use of environmental metabolomics to determine glyphosate level of exposure in rapeseed (*Brassica napus* L.) seedlings**  
I.L. Petersen, G. Tomasi, H. Sørensen, E.S. Boll, H.C.B. Hansen, J.H. Christensen  
*A novel untargeted environmental metabolomic approach is used to detect low-level glyphosate exposure of rapeseed seedlings.*
- 3078 Field sampling of soil pore water to evaluate trace element mobility and associated environmental risk**  
E. Moreno-Jiménez, L. Beesley, N.W. Lepp, N.M. Dickinson, W. Hartley, R. Clemente  
*In situ pore water sampling from a wide variety of soils proves to be a beneficial application to monitor the stability of pollutants in soils and subsequent risk through mobility.*



**CONTENTS – Continued from page II**

**3086 Global gene expression profile induced by the UV-filter 2-ethyl-hexyl-4-trimethoxycinnamate (EHMC) in zebrafish (*Danio rerio*)**

S. Zucchi, D.M. Oggier, K. Fent

*Gene expression changes by 2-ethyl-hexyl-4-trimethoxycinnamate in zebrafish.*

**3097 Accumulation, subcellular distribution and toxicity of inorganic mercury and methylmercury in marine phytoplankton**

Y. Wu, W.-X. Wang

*The inter-species difference in methylmercury and inorganic mercury toxicity in phytoplankton can be explained by cellular accumulation and subcellular distribution.*

**3106 Does small mammal prey guild affect the exposure of predators to anticoagulant rodenticides?**

D.G. Tosh, R.A. McDonald, S. Bearhop, N.R. Lilewellyn, S. Fee, E.A. Sharp, E.A. Barnett, R.F. Shore

*Restriction of the small mammal prey guild is associated with enhanced exposure of predators to anticoagulant rodenticides.*

**3113 Seasonal and spatial variation of diffuse (non-point) source zinc pollution in a historically metal mined river catchment, UK**

E. Gozzard, W.M. Mayes, H.A.B. Potter, A.P. Jarvis

*Diffuse zinc sources are an important source of instream contamination to mine-impacted rivers under varying flow conditions.*

**Short Communications**

**3123 Identifying genes and gene networks involved in chromium metabolism and detoxification in *Crambe abyssinica***

A. Zulfiqar, B. Paulose, S. Chhikara, O.P. Dhankher

*This study describes the identification and isolation of differentially expressed genes involved in chromium metabolism and detoxification in a non-food industrial oil crop *Crambe abyssinica*.*

**3129 Contrasting controls on arsenic and lead budgets for a degraded peatland catchment in Northern England**

J.J. Rothwell, K.G. Taylor, M.G. Evans, T.E.H. Allott

*Degraded peatlands can be significant sources of previously deposited arsenic and lead*

**3134 Degradability of aged aquatic suspensions of C<sub>60</sub> nanoparticles**

N.B. Hartmann, I.M. Buendia, J. Bak, A. Baun

*Aged suspensions of nC<sub>60</sub> were found not to be readily biodegradable according to the OECD test procedure. The biomass was not inhibited by the presence of nC<sub>60</sub>*

**3138 Trends in atmospheric reactive nitrogen for the Eastern United States**

R.W. Pinder, K.W. Appel, R.L. Dennis

*Ambient observations and simulation results indicate decreasing oxidized nitrogen levels over the Eastern US, but no trend in reduced nitrogen.*

**Corrigendum**

**3142 Corrigendum to “Time trends of perfluorinated compounds from the sediment core of Tokyo Bay, Japan (1950s–2004)” [Environ. Pollut. 158 (2010) 756–763]**

Y. Zushi, M. Tamada, Y. Kanai, S. Masunaga