

AQUACULTURE INTERNATIONAL

Volume 19 No. 1 February 2011

- Evaluation of corn steep liquor as a supplementary feed for *Labeo rohita* (Ham.) fingerlings / **S.G. Chovatiya, S.S. Bhatt and A.R. Shah** 1–12
- Effects of dietary protein and lipid levels and protein to energy ratios on growth performance and feed utilization of hatchery-reared juvenile spotted babylon (*Babylonia areolata*) / **N. Chaitanawisuti, S. Kritsanapuntu and W. Santaweesuk** 13–21
- Differences in fatty acid composition of egg capsules from broodstock spotted babylon, *Babylonia areolata*, fed a local trash fish and formulated diet under hatchery conditions / **N. Chaitanawisuti, S. Sangsawangchote and S. Piyatiratitivorakul** 23–31
- Socio-economic factors critical for intensification of fish farming technology. A case of selected villages in Morogoro and Dar es Salaam regions, Tanzania / **K. Wetengere** 33–49
- The effects of freshwater rearing on the whole body and muscle tissue fatty acid profile of the European sea bass (*Dicentrarchus labrax*) / **A.Ö. Hunt, F. Özkan, K. Engin and N. Tekelioğlu** 51–61
- Effects of early weaning strategies on growth, survival and digestive enzyme activities in cobia (*Rachycentron canadum* L.) larvae / **H.Q. Nguyen, H. Reinertsen, P.-A. Wold, T.M. Tran and E. Kjørsvik** 63–78
- Effects of dietary lipid levels on growth performance, feed utilization and fatty acid composition of juvenile Japanese seabass (*Lateolabrax japonicus*) reared in seawater / **J.-H. Xu, J. Qin, B.-L. Yan, M. Zhu and G. Luo** 79–89
- Antibacterial effect of medium-chain fatty acid: caprylic acid on gnotobiotic *Artemia franciscana* nauplii against shrimp pathogens *Vibrio harveyi* and *V. parahaemolyticus* / **G. Immanuel, M. Sivagnanavelmurugan and A. Palavesam** 91–101
- Effects of atrazine on the viability and the formation of byssus of the mussel *Mytilus galloprovincialis* / **D. Karagiannis, I.N. Vatsos and P. Angelidis** 103–110
- A preliminary study on the use of mannan oligosaccharides (MOS) in freshwater crayfish, *Astacus leptodactylus* Eschscholtz, 1823 juvenile diets / **Y. Mazlum, E. Yilmaz, M.A. Genç and O. Guner** 111–119

- Behavioural response of carp (*Cyprinus carpio*, L.) pond stock upon occurrence of hunting great cormorant (*Phalacrocorax carbo sinensis*) flocks / **J. Kortan and Z. Adámek** 121–129
- Effect of the interaction between body weight and temperature on growth and maximum daily food intake in sharpnout sea bream (*Diplodus puntazzo*) / **B. García García, J. Cerezo Valverde, F. Aguado-Giménez, J. García García and M.D. Hernández** 131–141
- Effect of partial fish meal replacement by soybean meal on the growth performance and biochemical indices of juvenile Japanese flounder *Paralichthys olivaceus* / **J. Ye, X. Liu, Z. Wang and K. Wang** 143–153
- Growth and survival of grouper *Epinephelus coioides* (Hamilton) larvae fed free-living nematode *Panagrellus redivivus* at first feeding / **O.S. Reyes, M.N. Duray, C.B. Santiago and M. Ricci** 155–164
- Effects of two dietary protein levels on energy balance and digestive capacity of *Octopus maya* / **C. Rosas, A. Sánchez, C. Pascual, J. Aguila, T. Maldonado and P. Domingues** 165–180
- Comparison of spawning induction techniques on *Paracentrotus lividus* (Echinodermata: Echinoidea) broodstock / **J. Gago and O.J. Luís** 181–191
- Tolerance response to ammonia and nitrite in hatchlings paralarvae of *Octopus vulgaris* and its toxic effects on prey consumption rate and chromatophores activity / **P. Feyjoo, R. Riera, B.C. Felipe, A. Skalli and E. Almansa** 193–204
- Antibacterial activities of a new combination of essential oils against marine bacteria / **S.M. Mousavi, G. Wilson, D. Raftos, S.S. Mirzargar and R. Omidbaigi** 205–214