

## ORIGINAL ARTICLES

<b>J. ROMEIS, M. MEISSEL</b>	Non-target risk assessment of <i>Bt</i> crops – Cry protein uptake by aphids	<b>1–6</b>
<b>C. GASPERS, B. D. SIEGFRIED, T. SPENCER, A. P. ALVES, N. P. STORER, I. SCHUPHAN, S. EBER</b>	Susceptibility of European and North American populations of the European corn borer to the Cry1F insecticidal protein	<b>7–16</b>
<b>E. J. G. PEREIRA, N. P. STORER, B. D. SIEGFRIED</b>	Fitness costs of Cry1F resistance in laboratory-selected European corn borer (Lepidoptera: Crambidae)	<b>17–24</b>
<b>D. L. DORHOUT, T. W. SAPPINGTON, L. C. LEWIS, M. E. RICE</b>	Flight behaviour of European corn borer infected with <i>Nosema pyrausta</i>	<b>25–37</b>
<b>V. DE LUIGI, L. FURLAN, S. PALMIERI, M. VETTORAZZO, G. ZANINI, C. R. EDWARDS, G. BURGIO</b>	Results of WCR monitoring plans and evaluation of an eradication programme using GIS and Indicator Kriging	<b>38–46</b>
<b>E. J. BERNKLAU, L. B. BJOSTAD, B. E. HIBBARD</b>	Synthetic feeding stimulants enhance insecticide activity against western corn rootworm larvae, <i>Diabrotica virgifera virgifera</i> (Coleoptera: Chrysomelidae)	<b>47–54</b>
<b>M. LESKINEN, I. MARKKULA, J. KOISTINEN, P. PYLKKÖ, S. OOPERI, P. SILJAMO, H. OJANEN, S. RAISKIO, K. TIILIKKALA</b>	Pest insect immigration warning by an atmospheric dispersion model, weather radars and traps	<b>55–67</b>
<b>M. BORGES, M. C. B. MORAES, M. F. PEIXOTO, C. S. S. PIRES, E. R. SUJII, R. A. LAUMANN</b>	Monitoring the Neotropical brown stink bug <i>Euschistus heros</i> (F.) (Hemiptera: Pentatomidae) with pheromone-baited traps in soybean fields	<b>68–80</b>
<b>R. O. OLATINWO, T. PRABHA, J. O. PAZ, D. G. RILEY, G. HOOGENBOOM</b>	The Weather Research and Forecasting (WRF) model: application in prediction of TSWV-vectors populations	<b>81–90</b>
<b>J. M. MARSHALL, A. J. STORER, I. FRASER, V. C. MASTRO</b>	A predictive model for detection of <i>Agrilus planipennis</i> (Col., Buprestidae) larvae in girdled ash ( <i>Fraxinus spp.</i> )	<b>91–97</b>
<b>A. MANRAKHAN, C. KOTZE</b>	Attraction of <i>Ceratitis capitata</i> , <i>C. rosa</i> and <i>C. cosyra</i> (Diptera: Tephritidae) to proteinaceous baits	<b>98–105</b>
<b>A. L. KNIGHT, C. PICKEL, L. HAWKINS, C. ABBOTT, R. HANSEN, L. HULL</b>	Monitoring oriental fruit moth (Lepidoptera: Tortricidae) and peach twig borer (Lepidoptera: Gelechiidae) with clear delta-shaped traps	<b>106–114</b>
<b>R. R. COSTA, J. C. MORAES, R. R. DACOSTA</b>	Feeding behaviour of the greenbug <i>Schizaphis graminum</i> on wheat plants treated with imidacloprid and/or silicon	<b>115–120</b>
<b>M. HORI, H. NAKAMURA, Y. FUJII, Y. SUZUKI, K. MATSUDA</b>	Chemicals affecting the feeding preference of the Solanaceae-feeding lady beetle <i>Henosepilachna vigintioctomaculata</i> (Coleoptera: Coccinellidae)	<b>121–131</b>
<b>A. KOMONEN, L. M. SCHROEDER, J. WESLIEN</b>	<i>Ips typographus</i> population development after a severe storm in a nature reserve in southern Sweden	<b>132–141</b>
<b>T. MALAUSA, A. FENIS, S. WAROT, J.-F. GERMAIN, N. RIS, E. PRADO, M. BOTTON, F. VANLERBERGHE-MASUTTI, R. SFORZA, C. CRUAUD, A. COULOUX, P. KREITER</b>	DNA markers to disentangle complexes of cryptic taxa in mealybugs (Hemiptera: Pseudococcidae)	<b>142–155</b>
<b>X. R. WANG, X. W. ZHU, X. C. KONG, M. M. MOTA</b>	A rapid detection of the pinewood nematode, <i>Bursaphelenchus xylophilus</i> in stored <i>Monochamus alternatus</i> by rDNA amplification	<b>156–159</b>