

CONTENTS

Original Articles

- Effects of taxonomic group, spatial scale and descriptor on the relationship between human activity and stream biota
A.G. Yates and R.C. Bailey (Ontario, Canada) 759
- Macrophytes in the anthropic constructions of the Venice littorals and their ecological assessment by an integration of the “CARLIT” index
A. Sfriso and C. Facca (Venice, Italy) 772
- Sensitivity of a cirolanid isopod to human pressure
V.G. Veloso (Rio de Janeiro, Brazil), G. Neves (Niterói, Brazil) and L. de Almeida Capper (Rio de Janeiro, Brazil) 782
- The landscape infrastructure footprint of oil development: Venezuela’s heavy oil belt
C.W. Baynard (Jacksonville, FL, USA) 789
- Dynamic sustainability assessment of countries at the macro level: A principal component analysis
H. Mirshojaeian Hosseini and S. Kaneko (Hiroshima, Japan) 811
- Trace elements accumulation and temporal trends in leaves of urban deciduous trees (*Aesculus hippocastanum* and *Tilia* spp.)
M. Aničić, T. Spasić, M. Tomašević, S. Rajšić and M. Tasić (Belgrade, Serbia) 824
- The influence of landscape structure on ants and dung beetles diversity in a Mediterranean savanna—Forest ecosystem
J.R. Verdú, C. Numa and O. Hernández-Cuba (Alicante, Spain) 831
- Macroinvertebrate-based multimetric predictive models for evaluating the human impact on biotic condition of Bolivian streams
N. Moya (Cochabamba, Bolivia), R.M. Hughes (Corvallis, Oregon, USA), E. Domínguez (Miguel Lillo, Argentina), F.-M. Gibon (Paris, France), E. Goitia (Cochabamba, Bolivia) and T. Oberdorff (Paris, France) 840
- Indicators of the impacts of development on environment: A comparison of Africa and Europe
B. Kestemont, L. Frendo and E. Zaccai (Bruxelles, Belgium) 848
- An urban university’s ecological footprint and the effect of climate change
C. Klein-Banai and T.L. Theis (Chicago, IL, USA) 857
- Fluctuating asymmetry as a biomarker of habitat fragmentation in an area-sensitive passerine, the Eurasian treecreeper (*Certhia familiaris*)
S. Helle (Turku, Finland), E. Huhta (Rovaniemi, Finland), P. Suorsa and H. Hakkarainen (Turku, Finland) ... 861
- A new methodology to infer the singing activity of an avian community: The Acoustic Complexity Index (ACI)
N. Pieretti, A. Farina and D. Morri (Urbino, Italy) 868
- A preliminary fish assemblage index for a transitional river–reservoir system in southeastern Brazil
B. de F. Terra and F.G. Araújo (Seropédica, Brazil) 874
- Bioluminescent bioreporter *Pseudomonas putida* TVA8 as a detector of water pollution. Operational conditions and selectivity of free cells sensor
G. Kuncova, J. Pazlarova, A. Hlavata (Prague, Czech Republic), S. Ripp and G.S. Sayler (Knoxville, TN, USA) ... 882
- Measurement of $\delta^{15}\text{N}$ in macroalgae stored in an environmental specimen bank for regional scale monitoring of eutrophication in coastal areas
I.G. Viana, J.A. Fernández, J.R. Aboal and A. Carballeira (Santiago de Compostela, Spain) 888

(Contents continued on page 1)

(Contents continued from Outside back cover)

Evaluation of five clustering algorithms for biodiversity surrogates R. Kent and Y. Carmel (Haifa, Israel)	896
Economic valuation of air pollution mortality: A 9-country contingent valuation survey of value of a life year (VOLY) B. Desaignes, D. Ami (Paris, France), A. Bartczak (Warsaw, Poland), M. Braun-Kohlová (Prague, Czech Republic), S. Chilton (Newcastle, UK), M. Czajkowski (Warsaw, Poland), V. Farreras (Barcelona, Spain), A. Hunt (Bath, UK), M. Hutchison (Paris, France), C. Jeanrenaud (Neuchatel, Switzerland), P. Kaderjak (Hungary), V. Máca (Prague, Czech Republic), O. Markiewicz, A. Markowska (Warsaw, Poland), H. Metcalf (Newcastle, UK), S. Navrud (Norway), J.S. Nielsen (Denmark), R. Ortiz (Bath, UK), S. Pellegrini (Neuchatel, Switzerland), A. Rabl (Paris, France), R. Riera (Barcelona, Spain), M. Scasny (Prague, Czech Republic), M.-E. Stoeckel (France), R. Szántó (Hungary) and J. Urban (Prague, Czech Republic)	902
Case Study	
Assessing trout farm pollution by biological metrics and indices based on aquatic macrophytes and benthic macroinvertebrates: A case study J.A. Camargo, C. Gonzalo and Á. Alonso (Madrid, Spain)	911
Quantifying surface gradients with a 2-band Enhanced Vegetation Index (EVI2) P. Mondal (Gainesville, FL, USA)	918
The occurrence of a <i>Euglena pascheri</i> and <i>Lepocinclis ovum</i> bloom in an oxbow lake in southern Poland under extreme environmental conditions S. Lięza (Lublin, Poland) and E. Wilk-Woźniak (Cracow, Poland)	925
Landscape metrics as indicators: Quantifying habitat network changes of a bush-cricket <i>Pholidoptera trans-sylvanica</i> in Hungary Z. Benedek (Budapest, Hungary), A. Nagy, I.A. Rácz (Debrecen, Hungary), F. Jordán (Povo, Italy) and Z. Varga (Debrecen, Hungary)	930