

CONTENTS

VOLUME 22, ISSUE No. 1

February 2011

Using non-homogeneous poisson models with multiple change-points to estimate the number of ozone exceedances in Mexico City <i>J. A. Achcar, E. R. Rodrigues and G. Tzintzun</i>	1
Assessing environmental stressors via bayesian model averaging in the presence of missing data <i>E. L. Boone, K. Ye and E. P. Smith</i>	13
Bayesian mixture model estimation of aerosol particle size distributions <i>D. Wraith, C. Alston, K. Mengersen and T. Hussein</i>	23
Recent statistical issues in multivariate receptor models <i>A. Pollice</i>	35
The institutional determinants of CO ₂ emissions: a computational modeling approach using Artificial Neural Networks and Genetic Programming <i>M. Álvarez-Díaz, G. Caballero-Miguez and M. Soliño</i>	42
Bimodal extension based on the skew-normal distribution with application to pollen data <i>H. W. Gómez, D. Elal-Olivero, H. S. Salinas and H. Bolfarine</i>	50
Multivariate quantiles in hydrological frequency analysis <i>F. Chebana and T. B. M. J. Ouarda</i>	63
Simulation and estimation of probabilities of phases of the Pacific Decadal Oscillation <i>N. Nairn-Birch, D. Diez, E. Eslami, M. M. Fauria, E. A. Johnson and F. P. Schoenberg</i>	79
Robustness of different regression modelling strategies in epidemiology: a time-series analysis of hospital admissions and air pollutants in Lisbon (1999–2004) <i>S. M. Sarmiento, T. G. Verburg, S. Marta Almeida, M. C. Freitas and H. Th. Wolterbeek</i>	86
A two-dimensional Markov renewal process <i>K. F. Davies and W. J. Braun</i>	98