

CONTENTS

Volume 159 Issue 2 2011

Review

- 351 Annual and seasonal trends of ambient ozone concentration and its impact on forest vegetation in Mercantour National Park (South-eastern France) over the 2000–2008 period**

P. Sicard, L. Dalstein-Richier, N. Vas

Trends of ambient ozone concentration, associated statistics and specific ozone-induced symptoms on conifers needles were studied in Mercantour National Park (South-eastern France) over the 2000–2008 period.

Regular Papers

- 363 Nitrogen isotope variations in camphor (*Cinnamomum Camphora*) leaves of different ages in upper and lower canopies as an indicator of atmospheric nitrogen sources**

H.-Y. Xiao, L.-H. Wu, R.-G. Zhu, Y.-L. Wang, C.-Q. Liu

Nitrogen isotope in camphor leaves indicating atmospheric nitrogen sources.

- 368 Trace metal uptake by tropical vegetables grown on soil amended with urban sewage sludge**

G. Nabulo, C.R. Black, S.D. Young

Trace metal uptake by tropical leaf vegetables can be predicted from dilute CaCl₂ extraction of soil but model parameters are genotype-specific.

- 377 Zero-valent iron and iron oxide-coated sand as a combination for removal of co-present chromate and arsenate from groundwater with humic acid**

M.S.H. Mak, P. Rao, I.M.C. Lo

Chromate and arsenate-contaminated groundwater can be better treated by using combination of zero-valent iron and iron oxide-coated sand.

- 383 Effect of an organic amendment on availability and bio-accessibility of some metals in soils of urban recreational areas**

M.d.C. Florido, F. Madrid, L. Madrid

Metal-containing amendments can deteriorate the environmental quality of soils of urban recreational areas.

- 391 Partitioning behaviour of perfluorinated alkyl contaminants between water, sediment and fish in the Orge River (nearby Paris, France)**

P. Labadie, M. Chevreuil

Sediment–water, biota–water and biota–sediment partitioning coefficients were determined for perfluorinated acids and sulfonates and were generally correlated with alkyl chain length.

- 398 The effects of the insecticide lambda-Cyhalothrin on the earthworm *Eisenia fetida* under experimental conditions of tropical and temperate regions**

M. Garcia, A. Scheffczyk, T. Garcia, J. Römbke

The effects of the insecticide lambda-cyhalothrin on earthworms did not differ considerably when performed in the same soil under different temperatures, but LC/EC₅₀ values varied by a factor of ten between OECD and tropical artificial soil.

- 401 Soil nitrogen transformations under elevated atmospheric CO₂ and O₃ during the soybean growing season**

E.I. Pujol Pereira, H. Chung, K. Scow, M.J. Sadowsky, C. van Kessel, J. Six

Our findings indicate that although elevated CO₂ increases plant biomass, N transformations were minimally affected. In contrast, elevated O₃ decreased soil mineral N likely through a reduction in plant material input and increased denitrification as indicated by the greater abundance of the denitrifier gene nosZ.

CONTENTS—Continued from outside back cover

- 408 The role of root hairs in cadmium acquisition by barley**
R. Zheng, H. Li, R. Jiang, V. Römheld, F. Zhang, F.-J. Zhao
The Cd uptake efficiency based on total root length was 8–45% lower in a barley root hairless mutant than in its wild-type, indicating an important role of root hairs in Cd acquisition.
- 416 Geographic variation in tissue accumulation of endocrine disrupting compounds (EDCs) in grazing sheep**
S.M. Rhind, C.E. Kyle, C. Mackie, K. Yates, E.I. Duff
Muscle concentrations of few of the endocrine disrupting compounds, measured in the muscle of sheep from regions exposed to greater pollution, were elevated.
- 423 Plant phenology, growth and nutritive quality of *Briza maxima*: Responses induced by enhanced ozone atmospheric levels and nitrogen enrichment**
J. Sanz, V. Bermejo, R. Muntifering, I. González-Fernández, B.S. Gimeno, S. Elvira, R. Alonso
*Forage quality and phenology are more O₃-sensitive than growth parameters in the Mediterranean annual grass *Briza maxima*.*
- 431 Nitrogen leaching and acidification during 19 years of NH₄NO₃ additions to a coniferous-forested catchment at Gårdsjön, Sweden (NITREX)**
F. Moldan, R.F. Wright
N addition has led to increased flux of N and increased C sequestration at a forested catchment in Sweden; the NO₃ released has partially offset recovery from acidification.
- 441 Bioanalytical and instrumental analysis of thyroid hormone disrupting compounds in water sources along the Yangtze River**
W. Shi, X. Wang, G. Hu, Y. Hao, X. Zhang, H. Liu, S. Wei, X. Wang, H. Yu
In vitro bioassay responses observed in Yangtze River source water extracts showed great TR antagonist activities, and DBP and DEHP were responsible.
- 449 Effects of nitrogen deposition and soil fertility on cover and physiology of *Cladonia foliacea* (Huds.) Willd., a lichen of biological soil crusts from Mediterranean Spain**
R. Ochoa-Hueso, E. Manrique
Nitrogen deposition and soil variables affect the physiology of terrestrial Mediterranean lichens.
- 458 How the redox state of tobacco 'Bel-W3' is modified in response to ozone and other environmental factors in a sub-tropical area?**
A.P.L. Dias, M. Dafré, M.C.S. Rinaldi, M. Domingos
The bioindicator efficiency of tobacco plants is not restrained under chronic doses of O₃ in the sub-tropical environment, as their redox state decreases few days after their placement in the field.
- 466 Toxic effects of oral 2-amino-4,6-dinitrotoluene in the Western fence lizard (*Sceloporus occidentalis*)**
C.A. McFarland, M.J. Quinn Jr., J. Boyce, E.M. LaFiandra, M.A. Bazar, L.G. Talent, M.S. Johnson
*The Western fence lizard (*Sceloporus occidentalis*) is a suitable reptile model for assessing the toxicity of energetic compounds and their metabolites.*
- 474 The immobilisation and retention of soluble arsenic, cadmium and zinc by biochar**
L. Beesley, M. Marmiroli
Biochar retains readily soluble cadmium and zinc from a contaminated soil, significantly reducing their potential mobility, but has limited effects on arsenic.
- 481 A rapidly equilibrating, thin film, passive water sampler for organic contaminants; characterization and field testing**
T. St. George, P. Vlahos, T. Harner, P. Helm, B. Wilford
An ethylene vinyl acetate (EVA), thin-film passive sampler for the detection of organic compounds in marine environments is calibrated and field tested.
- 487 Historical trends of polychlorinated dibenzo-p-dioxins and dibenzofurans in three dated sediment cores from Mexico**
C.-L. Yunuén, M.-Z.J. Vinicio, H.-D. Miguel Angel, L.L. Wee Kwong, J.A. Sanchez-Cabeza
Increasing concentration trends of PCDDs/Fs were observed for three sediment cores from Mexico suggesting the growing importance of local sources.
- 495 Uranium bioaccumulation and biological disorders induced in zebrafish (*Danio rerio*) after a depleted uranium waterborne exposure**
S. Barillet, C. Adam-Guillermin, O. Palluel, J.-M. Porcher, A. Devaux
This study demonstrates that U is highly bioaccumulated in fish, resulting in biological disorders such as hepatic oxidative stress as well as genotoxic and neurotoxic events.

CONTENTS—Continued from inside back cover

- 503 Modeling time-dependent toxicity to aquatic organisms from pulsed exposure of PAHs in urban road runoff**
W. Zhang, Y. Ye, Y. Tong, L. Ou, D. Hu, X. Wang
Toxicity to aquatic organisms after sequential pulsed exposure to PAHs in urban road runoff is evaluated.
- 509 Use of reporter-gene based bacteria to quantify phenanthrene biodegradation and toxicity in soil**
D. Shin, H.S. Moon, C.-C. Lin, T. Barkay, K. Nam
A cell-killing gene harboring reporter bacterium shows phenanthrene toxicity.
- 515 Mineralisation of target hydrocarbons in three contaminated soils from former refinery facilities**
M.G. Towell, J. Bellarby, G.I. Paton, F. Coulon, S.J.T. Pollard, K.T. Semple
Bioremediation strategy, native hydrocarbon concentrations and prior exposure histories of the microbial community influence hydrocarbon degradation in soil.
- 524 National survey of the levels of persistent organochlorine pesticides in the breast milk of mothers in China**
P. Zhou, Y. Wu, S. Yin, J. Li, Y. Zhao, L. Zhang, H. Chen, Y. Liu, X. Yang, X. Li
A survey of concentrations of OCPs in breast milk helps identify background concentrations in the Chinese population.
- 532 Adaptation, validation and application of the chemo-thermal oxidation method to quantify black carbon in soils**
T. Agarwal, T.D. Bucheli
Although widely used also for soils, the chemo-thermal oxidation method at 375 °C to quantify black carbon has never been properly validated for this matrix before.
- 539 Oxygen-containing polycyclic aromatic hydrocarbons (OPAHs) in urban soils of Bratislava, Slovakia: Patterns, relation to PAHs and vertical distribution**
B.A. Musa Bandowe, J. Sobocka, W. Wilcke
OPAHs and PAHs in urban soils are closely associated but OPAHs are faster translocated than PAHs.
- 550 Effects of compositional heterogeneity and nanoporosity of raw and treated biomass-generated soot on adsorption and absorption of organic contaminants**
B. Chen, W. Huang
The compositional heterogeneity and nano-structure play a regulating role in the adsorption and absorption of organic contaminants with the untreated and treated soot samples.
- 557 Nontarget analysis of polar contaminants in freshwater sediments influenced by pharmaceutical industry using ultra-high-pressure liquid chromatography–quadrupole time-of-flight mass spectrometry**
S. Terzic, M. Ahel
Comprehensive analysis of freshwater sediments by UPLC/QTOF indicated importance of pharmaceutically-derived polar contaminants.
- 567 Contamination by arsenic and other trace elements of tube-well water along the Mekong River in Lao PDR**
P. Chanpiwat, S. Sthiannopkao, K.H. Cho, K.-W. Kim, V. San, B. Suvanthon, C. Vongthavady
In the first international research on As and other trace element contamination of tube-well water in Lao PDR, concentrations of As, B, Ba, Mn, U, and Fe exceeding drinking water guidelines were found in samples taken mostly from Mekong River floodplain areas.
- 577 Atmospheric particle characterization, distribution, and deposition in Xi'an, Shaanxi Province, Central China**
Z. Cao, Y. Yang, J. Lu, C. Zhang
Atmospheric dust deposition rate is high and the levels of toxic elements associated with the settled dusts are elevated in Xi'an, Shaanxi, China.
- 585 Nutrient stoichiometry in *Sphagnum* along a nitrogen deposition gradient in highly polluted region of Central-East Europe**
M. Jiroušek, M. Hájek, L. Bragazza
*Regional climate and local forestry practices are expected to alter nutrient stoichiometry in *Sphagnum* mosses at high atmospheric N deposition in Central-East Europe.*
- 591 Effects of soil organic matter on the development of the microbial polycyclic aromatic hydrocarbons (PAHs) degradation potentials**
Y. Yang, N. Zhang, M. Xue, S.T. Lu, S. Tao
Soil organic matter regulated PAH degradation potentials through nutrition support and sequestration.
- 596 Compromised metamorphosis and thyroid hormone changes in wood frogs (*Lithobates sylvaticus*) raised on reclaimed wetlands on the Athabasca oil sands**
B.D. Hersikorn, J.E.G. Smits
This work provides guidance for reclamation of oil sands tailings and shows the usefulness of frogs and caging studies in environmental toxicology.
- 602 The relationships between air pollutants, meteorological parameters and concentration of airborne fungal spores**
A. Grinn-Gofroń, A. Strzelczak, T. Wolski
ANN models predict airspore contents from weather conditions and air pollutant.

CONTENTS—Continued from page IV

- 609 Toxicity profile of labile preservative bronopol in water: The role of more persistent and toxic transformation products**
N. Cui, X. Zhang, Q. Xie, S. Wang, J. Chen, L. Huang, X. Qiao, X. Li, X. Cai

The preservative bronopol is non-persistent and low toxic, but some transformation products can cause higher or prolonging adverse impacts.

- 616 Effects of biochar and the earthworm *Eisenia fetida* on the bioavailability of polycyclic aromatic hydrocarbons and potentially toxic elements**
J.L. Gomez-Eyles, T. Sizmur, C.D. Collins, M.E. Hodson

*Biochar decreased PAH bioavailability but was less effective at reducing PTE mobility, whilst *E. fetida* increased both PAH and PTE bioavailability.*

- 623 The lack of microbial degradation of polycyclic aromatic hydrocarbons from coal-rich soils**
C. Achten, S. Cheng, K.L. Straub, T. Hofmann

Native PAH compounds from coal-rich soils were not degraded by indigenous microorganisms in aerobic soil-slurry experiments.

- 630 Free atmospheric phosphine concentrations and fluxes in different wetland ecosystems, China**
C. Han, J. Geng, Y. Hong, R. Zhang, X. Gu, X. Wang, S. Gao, D. Glindemann

Environmental PH₃ concentrations in China are generally higher compared to other parts of the world.

- 636 Monitoring anthropogenic sewage pollution on mangrove creeks in southern Mozambique: A test of *Palaemon concinnus* Dana, 1852 (Palaemonidae) as a biological indicator**
G. Penha-Lopes, P. Torres, S. Cannicci, L. Narciso, J. Paula

*The identification of the effects of the peri-urban conditions on *P. concinnus* increases the choice of possible bioindicators in East African coastal waters.*

- 646 Assessing seasonal and spatial trends of persistent organic pollutants (POPs) in Indian agricultural regions using PUF disk passive air samplers**
K. Pozo, T. Harner, S.C. Lee, R.K. Sinha, B. Sengupta, M. Loewen, V. Geethalakshmi, K. Kannan, V. Volpi

New data on air concentrations of POPs across Indian agricultural regions is generated using cost-effective passive air samplers.

- 654 Long-chain perfluorinated chemicals in digested sewage sludges in Switzerland**
H. Sun, A.C. Gerecke, W. Giger, A.C. Alder

Concentrations of long-chain PFCs in digested sewage sludges vary substantially among different wastewater treatment plants.

Corrigendum

- 663 Corrigendum to “Metal and metalloid removal in constructed wetlands, with emphasis on the importance of plants and standardized measurements: A review” [Environ. Pollut. 158 (2010) 3447–3461]**
L. Marchand, M. Mench, D.L. Jacob, M.L. Otte