CONTENTS

Research Papers	
Potential of dry fractionation of wheat bran for the development of food ingredients, part I: Influence of ultra-fine grinding	
Y. Hemery, M. Chaurand, U. Holopainen, AM. Lampi, P. Lehtinen, V. Piironen, A. Sadoudi and X. Rouau	1
Potential of dry fractionation of wheat bran for the development of food ingredients, part II: Electrostatic separation of particles Y. Hemery, U. Holopainen, AM. Lampi, P. Lehtinen, T. Nurmi, V. Piironen, M. Edelmann and X. Rouau	9
Effects of glycerol on water properties and steaming performance of prefermented frozen dough L. Huang, J. Wan, W. Huang, P. Rayas-Duarte and G. Liu	19
Effects of roasting on barley β-glucan, thermal, textural and pasting properties P. Sharma, H.S. Gujral and C.M. Rosell	25
Water mobility, rheological and textural properties of rice starch gel S. Lu, JJ. Chen, YK. Chen, Cy. Lii, P. Lai and HH. Chen	31
Inheritance of flour paste viscosity is associated with a rice Waxy gene exon 10 SNP marker K. Traore, A.M. McClung, MH. Chen and R. Fjellstrom	37
Extraction and characterization of water-extractable and water-unextractable arabinoxylans from spelt bran: Study of the hydrolysis conditions for monosaccharides analysis E. Escarnot, M. Aguedo, R. Agneessens, B. Wathelet and M. Paquot	45
In vitro fermentation kinetics and end-products of cereal arabinoxylans and $(1,3;1,4)-\beta$ -glucans by porcine faeces B.A. Williams, D. Mikkelsen, L. le Paih and M.J. Gidley	53
Genetic variants of <i>HvGlb</i> 1 in Tibetan annual wild barley and cultivated barley and their correlation with malt quality	
X. Jin, S. Cai, Y. Han, J. Wang, K. Wei and G. Zhang	59
Impact of genetic and environmental factors on einkorn wheat (<i>Triticum monococcum</i> L. subsp. <i>monococcum</i>) polysaccharides	
A. Brandolini, A. Hidalgo, L. Plizzari and D. Erba	65
Inhibitory effect of whole oat on aberrant crypt foci formation and colon tumor growth in ICR and BALB/c mice HC. Wang, CH. Hung, JD. Hsu, MY. Yang, SJ. Wang and CJ. Wang	73
Magnetically modified spent grain for dye removal I. Safarik, K. Horska and M. Safarikova	78
Sensory texture of cooked rice is rather linked to chemical than to physical characteristics of raw grain C. Mestres, F. Ribeyre, B. Pons, V. Fallet and F. Matencio	81
Polyphenol composition and "in vitro" antiradical activity of single and multigrain breads A. Angioloni and C. Collar	90
Staling of frozen partly and fully baked breads. Study of the combined effect of amylopectin recrystallization and	
water content on bread firmness F. Ronda, P.A. Caballero, J. Quilez and Y.H. Roos	97
Waxy endosperm accompanies increased fat and saccharide contents in bread wheat (Triticum aestivum L.) grain	
T. Yasui and K. Ashida	104

(contents continued from	outside back	cover)
--------------------------	--------------	--------

Puroindolines, Pin alleles, hordoindolines and grain softness proteins are sources of bactericidal and fungicidal peptides	
R.L. Phillips, E.A. Palombo, J.F. Panozzo and M. Bhave	112
Improved nitrogen status enhances zinc and iron concentrations both in the whole grain and the endosperm fraction of wheat	
U.B. Kutman, B. Yildiz and I. Cakmak	118
Experimental data and modeling of the thermodynamic properties of bread dough at refrigeration and freezing temperatures T.G. Matuda, P.A. Pessôa Filho and C.C. Tadini	106
A novel method to prepare gluten-free dough using a meso-structured whey protein particle system L.E. van Riemsdijk, P.J.M. Pelgrom, A.J. van der Goot, R.M. Boom and R.J. Hamer	126 133
Rheological and thermal properties of masa as related to changes in corn protein during nixtamalization A.Q. Guzmán, M.E. Jaramillo Flores, J.S. Feria, M.G. Méndez Montealvo and YJ. Wang	139
Calendar	I