

CONTENTS

Aquatic environment**Simultaneous removal of hydrogen sulfide and toluene in a bioreactor: Performance and characteristics of microbial community**

Min Gao, Lin Li, Junxin Liu 353

The addition of H₂S could enhance the removal efficiency of toluene, because the change of the microenvironment of microbial growth resulted in an evolution of dominant microorganisms.

Reaction kinetic model for a recent co-produced water treatment technology

Abdulwahab M Ali Tuwati, Maohong Fan, Mark A. Bentley 360

This work provides a decarbonation model for water treatment industry.

Nitrification performance of nitrifying bacteria immobilized in waterborne polyurethane at low ammonia nitrogen concentrations

Yamei Dong, Zhenjia Zhang, Yongwei Jin, Zhirong Li, Jian Lu 366

A novel method for removing low concentration of ammonia nitrogen by nitrifying bacteria immobilized in waterborne polyurethane was developed.

Simultaneous removal of chromium and arsenate from contaminated groundwater by ferrous sulfate: Batch uptake behavior

Xiaohong Guan, Haoran Dong, Jun Ma, Irene M. C. Lo 372

Groundwater contaminated by both chromate and arsenate can be effectively remediated by Fe(II).

Removal of bentazone from micro-polluted water using MIEX resin: Kinetics, equilibrium, and mechanism

Zequn Liu, Xiaomin Yan, Mary Drikas, Danong Zhou, Dongsheng Wang, Min Yang, Jiuhui Qu 381

Using MIEX resin as the sorbent to remove bentazone contamination was an effective method in water supply treatment, and ion exchange was the dominant mechanism for bentazone removal by MIEX resin.

Treatment and remediation of a wastewater lagoon using microelectrolysis and modified DAT/IAT methods

Qiong Wu, Tao Hua, Qixing Zhou 388

The cost-effective microelectrolysis and modified demand aeration tank/intermittent aeration tank method was proposed to treat the wastewater lagoon with high toxicity and poor biodegradability.

Biodegradation of acrylamide by *Enterobacter aerogenes* isolated from wastewater in Thailand

Kanokhathai Buranasilp, Jittima Charoenpanich 396

As a novel acrylamide-degrading bacterium isolated from domestic wastewater, *Enterobacter aerogenes* can degrade acrylamide to acrylic acid and further to lower polarity compounds.

Removal of copper ions from aqueous solution by calcium alginate immobilized kaolin

Yanhui Li, Bing Xia, Quansheng Zhao, Fuqiang Liu, Pan Zhang, Qiuju Du, Dechang Wang, Da Li, Zonghua Wang, Yanzhi Xia 404

A novel environmental friendly material, calcium alginate immobilized kaolin, was prepared and used to remove heavy metal ions from aqueous solutions.

Atmospheric environment**Characterization of condensed phase nitric acid particles formed in the gas phase**

Long Jia, Yongfu Xu 412

The formation of nitric acid hydrates during the dark reaction of NO₂ with O₃ has been observed and characterized in a chamber, which is the first time to demonstrate the formation of condensed nitric acid can take place in the gas phase at room temperature.

On-road pollutant emission and fuel consumption characteristics of buses in Beijing

Aijuan Wang, Yunshan Ge, Jianwei Tan, Mingliang Fu, Asad Naeem Shah, Yan Ding, Hong Zhao, Bin Liang 419

Based on approximately 28,700 groups of instantaneous data obtained in Beijing using a portable emissions measurement system, on-road emission and fuel consumption characteristics of Euro III and IV buses fueled on diesel

and compressed natural gas were evaluated.

Polybrominated diphenyl ethers in atmosphere and soil of a production area in China: Levels and partitioning

Jun Jin, Ying Wang, Weizhi Liu, Congqiao Yang, Jicheng Hu, Jian Cui 427

Environmental behaviors of PBEDs were investigated in atmosphere and soil at a PBDE production source area of Laizhou Bay in China.

Terrestrial environment

Effect of moisture regime on the redistribution of heavy metals in paddy soil

Shunan Zheng, Mingkui Zhang 434

Soil moisture regime did not change the direction and pathways of transformation of metal speciation, however, it significantly affected the transformation rate.

Selective uptake of major and trace elements in *Erica andevalensis*, an endemic species to extreme habitats in the Iberian Pyrite Belt

Fabrizio Monaci, Eduardo O. Leidi, Maria Dolores Mingorance, Benito Valdés, Sabina Rossini Oliva, Roberto Bargagli 444

Erica andevalensis has an extraordinary capability to adapt to the harshest edaphic condition of Riotinto mining districts (SW Spain) and it is a priority species for promoting phytostabilization of pyrite mine tailings.

Plant coexistence can enhance phytoextraction of cadmium by tobacco (*Nicotiana tabacum* L.) in contaminated soil

Ling Liu, Yuefang Li, Jianjun Tang, Liangliang Hu, Xin Chen 453

Co-planting tobacco with Japanese clover can effectively increase phytoextraction of cadmium in Cd-contaminated soil.

Remediation of copper polluted red soils with clay materials

Gangya Zhang, Yunqing Lin, Mingkuang Wang 461

The bioavailability of Cu in heavy metal polluted red soils was effectively reduced with the remediation of montmorillonite.

Nitrogen sink in a small forested watershed of subtropical China

Laiming Huang, Jinling Yang, Ganlin Zhang 468

The typical forest ecosystem in subtropical China remains as nitrogen sink in contrast to nitrogen saturated ecosystems reported in many parts of Europe and North America.

Diversity of free-living nitrogen-fixing microorganisms in wastelands of copper mine tailings during the process of natural ecological restoration

Jing Zhan, Qingye Sun 476

The dominant plant, colonizing on wastelands of copper mine tailings during the process of natural ecological restoration, affected the diversity and structure of free-living diazotrophs by ameliorating the physico-chemical conditions of wastelands, and Proteobacteria and Cyanobacteria were the dominating groups in mine wastelands.

Factors affecting trace element content in periurban market garden subsoil in Yunnan Province, China

Yanqun Zu, Laurent Bock, Christian Schvartz, Gilles Colinet, Yuan Li 488

Relationships between Pb, Cd, Cu, and Zn content and physico-chemical characteristics in subsoil were investigated.

Environmental biology

Microbial population dynamics and changes in main nutrients during the acidification process of pig manures

Dongdong Zhang, Xufeng Yuan, Peng Guo, Yali Suo, Xiaofen Wang, Weidong Wang, Zongjun Cui 497

This study evaluated the metabolic transformations occurred during slurry acidification by using molecular methods.

Effect of chromium on accumulation and antioxidants in *Cucumis utillissimus* L.: Response under enhanced bioavailability condition

Geetgovind Sinam, Sarita Sinha, Shekhar Mallick 506

The accumulation of Cr(VI) and biochemical changes in *Cucumis utillissimus* L. in Cr spiked soil was studied and bioavailability conditions for forecasting the tolerance mechanism under high concentration were enhanced.

Environmental catalysis and materials

Effects of UV/Ag-TiO₂/O₃ advanced oxidation on unicellular green alga *Dunaliella salina*: Implications for removal of invasive species from ballast water

Donghai Wu, Hong You, Jiaxuan Du, Chuan Chen, Darui Jin 513

The UV/Ag-TiO₂/O₃ process has a potential for ballast water treatment, which was proven to be efficient for *Dunaliella salina* inactivation.

Conversion of the refractory ammonia and acetic acid in catalytic wet air oxidation of animal byproducts

Virginie Fontanier, Sofiane Zalouk, Stéphane Barbaty..... 520

The application of catalytic wet air oxidation to complex and highly inhomogeneous mixture could be envisaged as an alternative outlet of animal byproducts.

Serial parameter: CN 11-2629/X*1989*m*176*en*P*22*2011-3