

Special Issue: Risk indication of Genetically Modified

Up-scaling ecological effects of genetically modified plants in agriculture B. Breckling and H. Reuter (Bremen, Germany)	935
Risk indication of genetically modified organisms (GMO): Modelling environmental exposure and dispersal across different scales. Oilseed rape in Northern Germany as an integrated case study B. Breckling, H. Reuter (Bremen, Germany), U. Middelhoff (Berlin, Germany), M. Glemnitz, A. Wurbs (Müncheberg, Germany), G. Schmidt, W. Schröder (Vechta, Germany) and W. Windhorst (Kiel, Germany)	936
Remote sensing as a data source to analyse regional implications of genetically modified plants in agriculture—Oilseed rape (<i>Brassica napus</i>) in Northern Germany B. Breckling, H. Laue and H. Pehlke (Bremen, Germany)	942
Regionalisation of climate variability used for modelling the dispersal of genetically modified oil seed rape in Northern Germany G. Schmidt and W. Schröder (Vechta, Germany)	951
Derivation of regional crop sequences as an indicator for potential GMO dispersal on large spatial scales M. Glemnitz, A. Wurbs and R. Roth (Müncheberg, Germany)	964
GeneTraMP, a spatio-temporal model of the dispersal and persistence of transgenes in feral, volunteer and crop plants of oilseed rape and related species U. Middelhoff, H. Reuter and B. Breckling (Bremen, Germany)	974
Regional distribution of genetically modified organisms (GMOs)—Up-scaling the dispersal and persistence potential of herbicide resistant oilseed rape (<i>Brassica napus</i>) H. Reuter (Bremen, Germany), G. Schmidt, W. Schröder (Vechta, Germany), U. Middelhoff, H. Pehlke and B. Breckling (Bremen, Germany)	989
An integrative methodology to predict dispersal of genetically modified genotypes in oilseed rape at landscape-level—A study for the region of Schleswig-Holstein, Germany U. Middelhoff, E.-W. Reiche and W. Windhorst (Kiel, Germany)	1000