

## Original Articles

Differences in hoarding behaviors among six sympatric rodent species on seeds of oil tea ( <i>Camellia oleifera</i> ) in Southwest China <i>G. Chang and Z. Zhang</i> .....	165
Comparing distribution models for small samples of overdispersed counts of freshwater fish <i>L. Vaudor, N. Lamouroux and J.-M. Olivier</i> .....	170
Spatio-temporal change in the relationship between habitat heterogeneity and species diversity <i>A. González-Megías, J.M. Gómez and F. Sánchez-Piñero</i> .....	179
Evaluating and interpreting cross-taxon congruence: Potential pitfalls and solutions <i>M. Gioria, G. Bacaro and J. Feehan</i> .....	187
Plant species richness drives the density and diversity of Collembola in temperate grassland <i>A.C.W. Sabais, S. Scheu and N. Eisenhauer</i> .....	195
Long-term scenarios of the invasive black cherry in pine-oak forest: Impact of regeneration success <i>M. Vanhellemont, L. Baeten, H. Verbeeck, M. Hermy and K. Verheyen</i> .....	203
Niche overlap and species assemblage dynamics in an ageing pasture gradient in north-western France <i>T. Decaëns, P. Margerie, J. Renault, F. Bureau, M. Aubert and M. Hedde</i> .....	212
Differential response of bird functional traits to post-fire salvage logging in a boreal forest ecosystem <i>E.T. Azeria, J. Ibarzabal, C. Hébert, J. Boucher, L. Imbeau and J.-P.L. Savard</i> .....	220
Grazing effects on species composition in different vegetation types (La Palma, Canary Islands) <i>J.R. Arévalo, L. de Nascimento, S. Fernández-Lugo, J. Mata and L. Bermejo</i> .....	230
An integrated approach for the conservation of threatened plants: The case of <i>Arabis kennedyae</i> (Brassicaceae) <i>M. Andreou, P. Delipetrou, C. Kadis, G. Tsiamis, K. Bourtzis and K. Georghiou</i> .....	239
Feral livestock threatens landscapes dominated by columnar cacti <i>J.E. Malo, P. Acebes, S.M. Giannoni and J. Traba</i> .....	249

Contents continued from outside back cover

Seed germination of montane forest species in response to ash, smoke and heat shock in Mexico <i>S. Zuloaga-Aguilar, O. Briones and A. Orozco-Segovia</i> . . . . .	256
Diet selection variation of a large herbivore in a feeding experiment with increasing species numbers and different plant functional group combinations <i>L. Wang, D. Wang, J. Liu, Y. Huang and K.C. Hodgkinson</i> . . . . .	263
Exploring local borders of distribution in the shrub <i>Daphne laureola</i> : Individual and populations traits <i>A.R. Castilla, C. Alonso and C.M. Herrera</i> . . . . .	269
Photosynthesis, growth and foliar herbivory of four <i>Ardisia</i> species (Myrsinaceae) <i>J. Zhao and J. Chen</i> . . . . .	277
Belowground zone of influence in a tussock grass species <i>C. Armas and F.I. Pugnaire</i> . . . . .	284
Grazing-induced vegetation patchiness controls net N mineralization rate in a semi-natural grassland <i>N. Rossignol, A. Bonis and J.-B. Bouzillé</i> . . . . .	290