

## Insights

- 209 *E Müller, P Jud & W Nentwig* Artificial infection of *Cirsium arvense* with the rust pathogen *Puccinia punctiformis* by imitation of natural spore transfer by the weevil *Ceratopion onopordi*

## Review Papers

- 214 *C Andreasen & J C Streibig* Evaluation of changes in weed flora in arable fields of Nordic countries – based on Danish long-term surveys
- 227 *D R Clements & A Ditommaso* Climate change and weed adaptation: can evolution of invasive plants lead to greater range expansion than forecasted?

## Research Papers

- 241 *S L Poggio & C M Ghersa* Species richness and evenness as a function of biomass in arable plant communities
- 250 *B De Cauwer, T D'Hose, M Cougnon, B Leroy, R Bulcke & D Reheul* Impact of the quality of organic amendments on size and composition of the weed seed bank
- 261 *K A Kirkby, J E Pratley, D E Hume, M J Faville, M An & H Wu* Incidence of endophyte *Neotyphodium occultans* in *Lolium rigidum* from Australia
- 273 *N-O Bertholdsson* Use of multivariate statistics to separate allelopathic and competitive factors influencing weed suppression ability in winter wheat
- 284 *S S Kaundun, S-J Hutchings, R P Dale, G C Bailly & P Glanfield* Syngenta 'RISQ' test: a novel in-season method for detecting resistance to post-emergence ACCase and ALS inhibitor herbicides in grass weeds
- 294 *D Massa, B Krenz & R Gerhards* Target-site resistance to ALS-inhibiting herbicides in *Apera spica-venti* populations is conferred by documented and previously unknown mutations
- 304 *J L Gonzalez-Andujar, C Fernandez-Quintanilla, F Bastida, R Calvo, J Izquierdo & J A Lezaun* Assessment of a decision support system for chemical control of annual ryegrass (*Lolium rigidum*) in winter cereals
- 310 *D S Kim, E J P Marshall, P Brain & J C Caseley* Effects of crop canopy structure on herbicide deposition and performance