

CONTENTS

RESEARCH REVIEW

Progress in research on dry afterripening[†] 69–80

Iglesias-Fernández, R., del Carmen Rodríguez-Gacio, M. & Matilla, A.J.

RESEARCH OPINION

Granivore seed-size preferences 81–83

Radtke, T.M.

RESEARCH PAPERS

Tight translational control by the initiation factors eIF4E and eIF(iso)4E is required for maize seed germination 85–93

Dinkova, T.D., Márquez-Velázquez, N.A., Aguilar, R., Lázaro-Mixteco, P.E. & de Jiménez, E.S.

Rice seed quality development and temperature during late development and maturation 95–101

Ellis, R.H.

Timing of seed dispersal and dormancy, rather than persistent soil seed-banks, control seedling recruitment of woody plants in Neotropical savannas 103–116

Salazar, A., Goldstein, G., Franco, A.C. & Miralles-Wilhelm, F.

The germination niches of grassland species targeted for restoration: effects of seed pre-treatments 117–131

Wagner, M., Pywell, R.F., Knopp, T., Bullock, J.M. & Heard, M.S.

Environmental control of dormancy in quinoa (*Chenopodium quinoa*) seeds: two potential genetic resources for pre-harvest sprouting tolerance 133–141

Ceccato, D.V., Bertero, H.D. & Batlla, D.

TECHNICAL UPDATE

Visualization of molecular processes associated with seed dormancy and germination using MapMan 143–152

Joosen, R.V.L., Ligterink, W., Dekkers, B.J.W. & Hilhorst, H.W.M.

SHORT COMMUNICATIONS

Corrections for interferences and extraction conditions make a difference: use of the TBARS assay for lipid peroxidation of orthodox *Spartina pectinata* and recalcitrant *Spartina alterniflora* seeds during desiccation 153–158

Chappell, J.H., Jr & Cohn, M.A.

Using size-class structure to monitor growth of underdeveloped embryos in seeds of three *Aristolochia* species: implications for seed ecology 159–164

Adams, C.A., Baskin, J.M. & Baskin, C.C.