

Review Articles

Prospects for sustainable bioenergy production in selected former communist countries T. Srebotnjak (Berlin, Germany) and P. Hardi (Budapest, Hungary)	1009
Methodology to select a set of urban sustainability indicators to measure the state of the city, and performance assessment N. Munier (Valencia, Spain)	1020
Deadwood as a surrogate for forest biodiversity: Meta-analysis of correlations between deadwood volume and species richness of saproxylic organisms A. Lassauce (Nogent-sur-Vernisson, France and Angers, France), Y. Paillet (Nogent-sur-Vernisson, France), H. Jactel (Cestas, France) and C. Bouget (Nogent-sur-Vernisson, France)	1027

Original Articles

Are butterflies and moths suitable ecological indicator systems for restoration measures of semi-natural calcareous grassland habitats? L. Rákósy (Cluj, Romania) and T. Schmitt (Trier, Germany)	1040
Combining geostatistical models and remotely sensed data to improve tropical tree richness mapping J.L. Hernández-Stefanoni (Mérida, Mexico), J. Alberto Gallardo-Cruz, J.A. Meave (México, Mexico) and J.M. Dupuy (Mérida, Mexico)	1046
Non-biting midges in biodiversity conservation and environmental assessment: Findings from boreal freshwater ecosystems J. Raunio (Kouvola, Finland), J. Heino (Oulu, Finland) and L. Paasivirta (Salo, Finland)	1057
Constraints in the identification and interpretation of ozone as a significant predictor of effects on the supersensitive indicator <i>Nicotiana tabacum</i> Bel-W3 in biomonitoring studies F. Cristofolini, A. Cristofori, E. Gottardini (San Michele all' Adige, Italy), S. Maccherini and M. Ferretti (Siena, Italy)	1065
Indicators for sustainable land use management in Santiago de Chile U. Weiland, A. Kindler, E. Banzhaf, A. Ebert (Leipzig, Germany) and S. Reyes-Paecke (Santiago de Chile, Germany)	1074
Analyses of interventions and structural breaks in marine and fisheries time series: Detection of shifts using iterative methods J.P. Gröger (Hamburg, Germany and Rostock, Germany), M. Missong (Bremen, Germany) and R.A. Rountree (Waquoit, MA, USA)	1084
Land use disturbance indicators and water quality variability in the Biscayne Bay Watershed, Florida R.O. Carey, K.W. Migliaccio, Y. Li, B. Schaffer (Homestead, FL, USA), G.A. Kiker and M.T. Brown (Gainesville, FL, USA)	1093
Rhizomes and fronds of <i>Athyrium filix-femina</i> as possible bioindicators of chemical elements from soils over different parent materials in southwest Poland A. Samecka-Cymerman, K. Kolon, A. Stankiewicz, J. Kaszewska, L. Mróz (Wroclaw, Poland) and A.J. Kempers (Nijmegen, The Netherlands)	1105
Evaluating the influence of off-shore cage aquaculture on the benthic ecosystem in Alghero Bay (Sardinia, Italy) using AMBI and M-AMBI A. Forchino (Varese, Italy), A. Borja (Portualdea, Spain), F. Brambilla (Varese, Italy), J.G. Rodríguez, I. Muxika (Portualdea, Spain), G. Terova and M. Saroglia (Varese, Italy)	1112

Assessment of soil microbial diversity measurements as indicators of soil functioning in organic and conventional horticulture systems R. Romaniuk, L. Giuffr�, A. Costantini (Buenos Aires, Argentina) and P. Nannipieri (Firenze, Italy)	1345
Functional convergence of fish assemblages in urban streams of Brazil and the United States A.M. Cunico (Paran�, Brazil), J.D. Allan (Michigan, MI, USA) and A.A. Agostinho (Paran�, Brazil)	1354
Analysis of spatial and temporal variation in the levels of microbial fecal indicators in the major rivers flowing into the Aegean Sea, Turkey A. Kacar (Inciralti, Turkey)	1360
Forest proportion as indicator of ecological integrity in streams using Plecoptera as a proxy J. T�rnblom (Skinnskatteberg, Sweden), E. Degerman (�rebro, Sweden) and P. Angelstam (Skinnskatteberg, Sweden)	1366
Historical perspectives of material use in Czechoslovakia in 1855–2007 J. Kovanda and T. Hak (Prague, Czech Republic)	1375
Lessons from seven sustainability indicator programs in developing countries of Asia S. Krank and H. Wallbaum (Zurich, Switzerland)	1385
A proposal of framework to obtain an integrated biodiversity indicator for agricultural ponds incorporating the simultaneous effects of multiple pressures T. Kadoya, M. Akasaka (Ibaraki, Japan), T. Aoki (Kobe, Japan) and N. Takamura (Ibaraki, Japan)	1396
Using chlorophyll <i>a</i> and cyanobacteria in the ecological classification of lakes M. S�ndergaard (Silkeborg, Denmark and Beijing, China), S.E. Larsen, T.B. J�rgensen (Silkeborg, Denmark) and E. Jeppesen (Silkeborg, Denmark and Beijing, China)	1403
Application of detectability in the use of indicator species: A case study with birds J.E. Quinn, J.R. Brandle (Lincoln, NE, USA), R.J. Johnson (Clemson, SC, USA) and A.J. Tyre (Lincoln, NE, USA)	1413
An emergy analysis-based methodology for eco-efficiency evaluation of building manufacturing D. Li, J. Zhu (Nanjing, PR China), E.C.M. Hui, B.Y.P. Leung (Kowloon, Hong Kong) and Q. Li (Nanjing, PR China)	1419
<i>Donax trunculus</i> (Bivalvia: Donacidae) as a potential biological indicator of grain-size variations in beach sediment P. La Valle, L. Nicoletti, M.G. Finoina and G.D. Ardizzone (Rome, Italy)	1426
Forest transition in an ecologically important region: Patterns and causes for landscape dynamics in the Niger Delta A.O. Onojeghuo and G.A. Blackburn (Lancaster, UK)	1437
Towards a global criteria based framework for the sustainability assessment of bioethanol supply chains. Application to the Swiss dilemma: Is local produced bioethanol more sustainable than bioethanol imported from Brazil? T. Corbi�re-Nicollier (Lausanne, Switzerland), I. Blanc (France) and S. Erkman (Lausanne, Switzerland)	1447
Improving national habitat specific biodiversity indicators using relative habitat use for common birds J.L. Larsen, H. Heldbjerg and A. Eskildsen (Copenhagen, Denmark)	1459
Case Study	
Fragmentation in the Legal Amazon, Brazil: Can landscape metrics indicate agricultural policy differences? F. Colson (Wilrijk, Belgium), J. Bogaert (Brussels, Belgium) and R. Ceulemans (Wilrijk, Belgium)	1467
A multiple watershed field test of hydrogeomorphic functional assessment of headwater streams—Variability in field measurements between independent teams J. Berkowitz, A.F. Casper and C. Noble (Vicksburg, MS, USA)	1472
Remote sensing data predict indicators of soil functioning in semi-arid steppes, central Spain M. Garc�a-G�mez (Madrid, Spain and M�stoles, Spain) and F.T. Maestre (M�stoles, Spain)	1476
Application of the index of marsh bird community integrity to coastal wetlands of Georgian Bay and Lake Ontario, Canada L.A. Smith-Cartwright and P. Chow-Fraser (Hamilton, Canada)	1482
Use of sex ratio of bream (<i>Abramis brama</i> L.) as an indicator of endocrine effects: Results from the German Environmental Specimen Bank D. Teubner, K. Tarricone, M. Veith, G. Wagner, M. Bartel-Steinbach, R. Klein (Trier, Germany), A. K�rner (Berlin, Germany), M. Quack and M. Paulus (Trier, Germany)	1487
Letter to the Editor	
Extrapolation and biodiversity indicators: Handle with caution! J.-P. Rossi (Cestas, France)	1490
The Emperor’s new clothes? A comment on Besse-Lototskaya et al. (2011) M. Kelly (Durham, UK)	1492
Erratum	
Corrigendum to “Implementing landscape indices to predict stream water quality in an agricultural setting: An assessment of the Lake and River Enhancement (LARE) protocol in the Mississinewa River watershed, East-Central Indiana” by Shiels [Ecol. Indic. 10(6) (2010) 1102–1110] D.R. Shiels (Upland, IN, USA)	1495

(Contents continued from Outside back cover)

Connectivity between estuaries and marine environment: Integrating metrics to assess estuarine nursery function	
R.P. Vasconcelos, P. Reis-Santos, M.J. Costa and H.N. Cabral (Lisboa, Portugal)	1123
Korcak dimension as a novel indicator of landscape fragmentation and re-forestation	
A.R. Imre, D. Cseh (Budapest, Hungary), M. Neteler and D. Rocchini (Michele all'Adige, Italy)	1134
Examination of the bioindicator potential of <i>Oncholaimus campyloceroides</i> (Oncholaimidae, Nematoda) from Bizerte bay (Tunisia)	
F. Boufahja, A. Hedfi, J. Amorri, P. Aïssa, H. Beyrem and E. Mahmoudi (Zarzouna, Tunisia)	1139
Monitoring changes in lichen resources for range management purposes in reindeer husbandry	
A. Olofsson, Ö. Danell, P. Forslund and B. Åhman (Uppsala, Sweden)	1149
Using a sub-pixel mapping model to improve the accuracy of landscape pattern indices	
X. Li, Y. Du, F. Ling, S. Wu and Q. Feng (Wuhan, China)	1160
Concordance among assemblages of upland Amazonian lakes and the structuring role of spatial and environmental factors	
P.M. Lopes, A. Caliman, L.S. Carneiro (Rio de Janeiro, Brazil), L.M. Bini (Goiania, Brazil), F.A. Esteves, V. Farjalla (Rio de Janeiro, Brazil and Goiania, Brazil) and R.L. Bozelli (Rio de Janeiro, Brazil)	1171
How to choose a biodiversity indicator – Redundancy and complementarity of biodiversity metrics in a freshwater ecosystem	
B. Gallardo (Zaragoza, Spain), S. Gascón, X. Quintana (Girona, Spain) and F.A. Comín (Zaragoza, Spain)	1177
Energy-based complexity measures in natural and social systems	
S. Ulgiati, M. Ascione, A. Zucaro (Napoli, Italy) and L. Campanella (Roma, Italy)	1185
Spatial factors contribute to benthic diatom structure in streams across spatial scales: Considerations for biomonitoring	
N.J. Smucker and M.L. Vis (Athens, OH, USA)	1191
Development of metrics based on fish body size and species traits to assess European coldwater streams	
M. Logez (Aix-en-Provence, France) and D. Pont (Antony, France)	1204
Towards a process-based evaluation of land vulnerability to soil degradation in Italy	
L. Salvati, S. Bajocco, T. Ceccarelli, M. Zitti and L. Perini (Rome, Italy)	1216
An approach to identifying potential surrogates of periphytic ciliate communities for monitoring water quality of coastal waters	
H. Xu, W. Zhang, Y. Jiang (Qingdao, China), G.-S. Min and J.-K. Choi (Incheon, Republic of Korea)	1228
Can biotic indices detect mild organic enrichment of the seafloor?	
L. Sampaio, A.M. Rodrigues and V. Quintino (Aveiro, Portugal)	1235
Physiological and isotopic signals in epilithic mosses for indicating anthropogenic sulfur on the urban-rural scale	
X.-Y. Liu, H.-Y. Xiao and C.-Q. Liu (Guiyang, China)	1245
Functional Group Density as an index for assessing habitat quality in tallgrass prairie	
V.A. Sivicek and J.B. Taft (Champaign, IL, USA)	1251
Identifying grasshopper bioindicators for habitat quality assessment of ecological networks	
C.S. Bazelet and M.J. Samways (Matieland, South Africa)	1259
Effects of forest continuity and tree age on epiphytic lichen biota in coniferous forests in Estonia	
L. Marmor, T. Tõrra, L. Saag and T. Randlane (Tartu, Estonia)	1270
Indicators to support environmental sustainability of bioenergy systems	
A.C. McBride, V.H. Dale, L.M. Baskaran, M.E. Downing, L.M. Eaton, R.A. Efroymson, C.T. Garten Jr., K.L. Kline, H.I. Jager, P.J. Mulholland, E.S. Parish, P.E. Schweizer and J.M. Storey (Oak Ridge, TN, USA)	1277
Evaluation of coastal perturbations: A new mathematical procedure to detect changes in the reference state of coastal systems	
E. Goberville (Wimereux, France and Arcachon, France), G. Beaugrand (Wimereux, France), B. Sautour (Arcachon, France) and P. Tréguer (Plouzané, France)	1290
Contribution of habitat patches to network connectivity: Redundancy and uniqueness of topological indices	
G. Baranyi (Budapest, Hungary), S. Saura (Madrid, Spain), J. Podani (Budapest, Hungary) and F. Jordán (Trento, Italy)	1301
Diversity of plant-animal interactions: Possibilities for a new plant defense indicator value?	
F. Borchard (Lünen, Germany), H.-J. Berger, M. Bunzel-Drüke (Bad Sassendorf, Germany) and T. Fartmann (Münster, Germany)	1311
Corticolous lichens as environmental indicators in urban areas in southern Brazil	
M.I. Käffer, S.M. de Azevedo Martins, C. Alves, V.C. Pereira, J. Fachel and V.M.F. Vargas (Porto Alegre, Brazil)	1319
Modeling and dynamic assessment of urban economy-resource-environment system with a coupled system dynamics – geographic information system model	
D. Guan (Saga, Japan), W. Gao (Kitakyushu, Japan), W. Su (Chongqing, China and Guiyang, China), H. Li and K. Hokao (Saga, Japan)	1333