

## CONTENTS

## RESEARCH PAPERS

- Remote estimation of chlorophyll on two wheat cultivars in two rainfed environments  
*Daive Cammarano, Glenn Fitzgerald, Bruno Basso, Deli Chen, Peter Grace, Garry O'Leary* 269
- The 1BL/1RS translocation decreases grain yield of spring wheat germplasm in low yield environments of north-eastern Australia  
*Allan S. Peake, Arthur Gilmour, Mark Cooper* 276
- Modelling yield response of a traditional and a modern barley cultivar to different water and nitrogen levels in two contrasting soil types  
*L. Gabriela Abeledo, Daniel F. Calderini, Gustavo A. Slafer* 289
- Nitrogen and sulfur uptake and remobilisation in canola genotypes with varied N- and S-use efficiency differ at vegetative and maturity stages  
*Tatjana Balint, Zdenko Rengel* 299
- Indirect selection using reference and probe genotype performance in multi-environment trials  
*Ky L. Mathews, Richard Trethowan, Andrew W. Milgate, Thomas Payne, Maarten van Ginkel, Jose Crossa, Ian DeLacy, Mark Cooper, Scott C. Chapman* 313
- Diagnosis and simulation: a suitable combination to support farming systems design  
*Guillaume Martin, Jean-Pierre Theau, Olivier Therond, Roger Martin-Clouaire, Michel Duru* 328
- Soil organic carbon and total nitrogen under *Leucaena leucocephala* pastures in Queensland  
*Alejandro Radrizzani, H. Max Shelton, Scott A. Dalzell, Gunnar Kirchhof* 337
- Sward and tiller growth dynamics of *Lolium perenne* L. as affected by defoliation frequency during autumn  
*Jonathan A. Poff, Oscar A. Balocchi, Ignacio F. López* 346
- The use of conjoint analysis to determine the relative importance of specific traits as selection criteria for the improvement of perennial pasture species in Australia  
*K. F. Smith, P. F. Fennessy* 355