

CONTENTS

Aquatic environment

Chemical cleaning of fouled PVC membrane during ultra-filtration of algal-rich water

Yan Zhang, Jiayu Tian, Heng Liang, Jun Nan, Zhonglin Chen, Guibai Li 529

Among the tested cleaning reagents (NaOH, HCl, EDTA, and NaClO), NaClO exhibited the best performance in removing the irreversible fouling resistance, while the negligible membrane permeability restoration was achieved after NaOH cleaning.

Effects of activated sludge flocs and pellets seeds on aerobic granule properties

Huacheng Xu, Pinjing He, Guanzhao Wang, Liming Shao 537

Extracellular polymeric substance-free pellets have more advantages than activated sludge floc as the seed for aerobic granulation.

Destruction of methylphosphonic acid in a supercritical water oxidation bench-scale double wall reactor

Bambang Veriansyah, Eun-Seok Song, Jae-Duck Kim 545

The destruction efficiency and reaction rate of methylphosphonic acid in a supercritical water oxidation bench-scale double wall reactor were determined.

Petroleum hydrocarbons and their effects on fishery species in the Bohai Sea, North China

Sisi Xu, Jinming Song, Huamao Yuan, Xuegang Li, Ning Li, Liqin Duan, Yu Yu 553

The semiquantitative effects of PHs on population growth rates of several common fishery species in the Bohai Sea were assessed.

Coupling of zero valent iron and biobarriers for remediation of trichloroethylene in groundwater

Mullika Teerakun, Alissara Reungsang, Chien-Jung Lin, Chih-Hsiang Liao 560

Permeable reactive barriers series consisting of iron filling base barrier followed by anaerobic bioreactive barrier and aerobic bioreactive barrier, respectively, could remove high concentrations of TCE as well as its metabolites.

Inhibition and recovery of nitrification in treating real coal gasification wastewater with moving bed biofilm reactor

Huiqiang Li, Hongjun Han, Maoan Du, Wei Wang 568

The factors inhibiting nitrification in treating coal gasification wastewater using moving bed biofilm reactor were determined and the feasible recovery methods were also evaluated.

Cyanobacteria-/cyanotoxin-contaminations and eutrophication status before Wuxi Drinking Water Crisis in Lake Taihu, China

Yongmei Liu, Wei Chen, Dunhai Li, Zebo Huang, Yinwu Shen, Yongding Liu 575

The present work has put forward the viewpoint of comparison study in eutrophic lakes and will lead to a better understanding of the eutrophication status and the potential risks before "Wuxi Drinking Water Crisis" in Lake Taihu.

Changes in different organic matter fractions during conventional treatment and advanced treatment

Chao Chen, Xiaojian Zhang, Lingxia Zhu, Wenjie He, Hongda Han 582

One step XAD-8 resin isolation which divided organic matter into the hydrophobic and hydrophilic fractions helped to understand the mechanism of NOM removal in conventional and advanced water treatment processes.

Evaluation of reusing alum sludge for the coagulation of industrial wastewater containing mixed anionic surfactants

Siriprapha Jangkorn, Sinchai Kuhakaew, Suwapee Theantano, Harit Klinla-or, Tongchai Sriwiriyarat 587

The alum sludge was reused as a coagulant or coagulation aid to minimize the fresh alum dosage or enhance the removal efficiency in coagulating the industrial wastewater containing anionic surfactants.

Development and application of a water pollution emergency response system for the Three Gorges Reservoir in the Yangtze River, China

Qiang He, Shujuan Peng, Jun Zhai, Haiwen Xiao 595

An integrated geographic information system-based water pollution management information system for the Three Gorges Reservoir Area (WPMS ER TGRA) was developed to simulate the transfer and diffusion of accidental pollutants in the river, and quickly identify the affected area and how it will change over time within a few minutes of an accident occurring

Terrestrial environment

Arsenic uptake, accumulation and phytofiltration by duckweed (*Spirodela polyrhiza* L.)

Xin Zhang, Ying Hu, Yunxia Liu, Baodong Chen 601

Duckweed *Spirodela polyrhiza* L. was able to survive in high concentration of arsenate solution and showed the potential for arsenic phytofiltration.

Mineral materials as feasible amendments to stabilize heavy metals in polluted urban soils

Mingkui Zhang, Jincheng Pu 607

Calcium magnesium phosphate and palygorskite treatments were superior to agricultural limestone and rock phosphate for stabilizing heavy metals in polluted urban soils.

Impacts of wastewater sludge amendments in restoring nitrogen cycle in *p*-nitrophenol contaminated soil

F. Olcay Topac Sagban 616

The study evaluated the impacts of wastewater sludge amendments in restoring nitrogen-related processes, such as urease activity, nitrification potential and arginine ammonification rate, in *p*-nitrophenol contaminated soil.

Speciation and bioaccessibility of lead and cadmium in soil treated with metal-enriched Indian mustard leaves

Yanshan Cui, Jin Fu, Xiaochen Chen 624

Lead and cadmium risk in metal-enriched Indian mustard leaves should be considered in phytoremediation system in which metal might be brought back to soil and change over time.

Effect of water regimes and organic matters on transport of arsenic in summer rice (*Oryza sativa* L.)

Sefaur Rahaman, Ashim Chandra Sinha, Dibyendu Mukhopadhyay 633

Saturation of paddy field instead of continuous ponding alone or in combination with organic matters could minimize arsenic load in summer rice and corresponding soils.

Effect of pH on transport of Pb^{2+} , Mn^{2+} , Zn^{2+} , and Ni^{2+} through lateritic soil: Column experiments and transport modeling

Srilert Chotpantararat, Say Kee Ong, Chakkaphan Sutthirat, Khemarath Osathaphan 640

The effects of pH on the sorption and transport of Pb^{2+} , Mn^{2+} , Zn^{2+} , and Ni^{2+} through lateritic aquifer were determined and modeled using equilibrium and chemical non-equilibrium models.

Environmental biology

Characterization of the effective cellulose degrading strain CTL-6

Yucai Lü, Xiaofen Wang, Ning Li, Xiaojuan Wang, Masaharu Ishii, Yasuo Igarashi, Zongjun Cui 649

A thermophilic anaerobic bacterium CTL-6 belonged to the genus *Clostridium thermocellum* was characterized; and its striking capability to degrade cellulose and high efficiency value of the endocellulase were observed.

Competition of bloom-forming marine phytoplankton at low nutrient concentrations

Hanhua Hu, Jun Zhang, Weidong Chen 656

Growth competition of three kinds of bloom-forming marine phytoplankton was observed in laboratory conditions and it was shown that diatom was a poor competitor compared to the dinoflagellates at limited phosphate levels.

Environmental health and toxicology

Differential response of multiple zebrafish hepatic F-box protein genes to 17α -ethinylestradiol treatment

Hongping Zheng, Shifeng Li, Zhili Wu, Yunbin Zhang, Shengnan Hu, Yuanchang Yan, Yiping Li 664

The expression of hepatic mRNA concentration of FBP genes were quantified based on the expression of the optimal internal control gene in male zebrafish exposed to EE2.

In vitro thyroid disrupting effects of organic extracts from WWTPs in Beijing

Na Li, Mei Ma, Kaifeng Rao, Zijian Wang 671

A two-hybrid yeast assay was used to evaluate the thyroid receptor (TR) mediated effects in three WWTPs, and TR antagonistic activities were found in all processes from the studied WWTPs.

Individual and combined toxic effects of cypermethrin and chlorpyrifos on earthworm

Shiping Zhou, Changqun Duan, Wong Hang Gi Michelle, Fazhong Yang, Xuehua Wang 676

The toxic effects of combined application of chlorpyrifos and cypermethrin on earthworm were investigated and the results demonstrated that the mixture of the two pesticide caused stronger effects on endpoints, such as growth and reproduction, than the individual application.

Environmental catalysis and materials

Experimental and density functional theory study of the adsorption of N₂O on ion-exchanged ZSM-5: Part II. The adsorption of N₂O on main-group ion-exchanged ZSM-5

Bo Zhang, Yongan Lu, Hong He, Jianguo Wang, Changbin Zhang, Yunbo Yu, Li Xue 681

The adsorption and desorption behaviors of N₂O on main group ion-exchanged ZSM-5 were evaluated, and the results showed that alkali earth cations are more adsorptive to N₂O than alkali cations since alkali earth cations with higher valence and larger radius lead to stronger interaction between metal cations and N₂O

Effect of platinum on the photocatalytic degradation of chlorinated organic compound

Chih Ming Ma, Ya Wen Lee, Gui Bing Hong, Te Li Su, Je Lueng Shie, Chang Tang Chang 687

Different Pt-modified TiO₂ films coated on quartz substrates were prepared and their photocatalytic activities with their DCM removal performance were compared.

Environmental analytical methods

Improved method for analyzing the degradation of estrogens in water by solid-phase extraction coupled with ultra performance liquid chromatography-ultraviolet detection

Minggang Zheng, Ling Wang, Yuandui Bi, Feng Liu 693

The novel method was established to separate and detect four estrogens by SPE-UPLC-UV and the method was applied to examine 17β-estradiol and 17α-ethynylestradiol changes in simulated natural water to determine the kinetic processes and decomposition products.

Municipal solid waste and green chemistry

Catalytic oxidation of gas-phase elemental mercury by nano-Fe₂O₃

Fanhai Kong, Jianrong Qiu, Hao Liu, Ran Zhao, Zhihui Ai · 699

The study focused on the heterogeneous interactions between nano-Fe₂O₃ and Hg⁰ in simulated flue gas. The effects of oxygen concentration, bed temperature, water vapour concentration and particle size were also discussed.

Serial parameter: CN 11-2629/X*1989*m*176*en*P*25*2011-4