

■ **Review**

Armstrong DP, Wittmer HU

Incorporating Allee effects into reintroduction strategies 687

■ **Original Articles**

Xiang W, Liu S, Deng X, Shen A, Lei X, Tian D, Zhao M, Peng C

General allometric equations and biomass allocation of *Pinus massoniana* trees on a regional scale in southern China 697

Ge Q, Dai J, Zheng J, Bai J, Zhong S, Wang H, Wang W-C

Advances in first bloom dates and increased occurrences of yearly second blooms in eastern China since the 1960s: further phenological evidence of climate warming 713

Ishikawa H

Effects of dung beetles on seedling emergence from herbaceous seeds in the dung of sika deer (*Cervus nippon*) in a temperate Japanese grassland ecosystem 725

Zhang H, Zhou X, Lu F, Pang J, Feng Z, Liu W, Ouyang Z, Wang X

Seasonal dynamics of soil CO₂ efflux in a conventional tilled wheat field of the Loess Plateau, China 735

Luiselli L, Madsen T, Capizzi D, Rugiero L, Pacini N, Capula M

Long-term population dynamics in a Mediterranean aquatic snake 745

Gu J, Yu S, Sun Y, Wang Z, Guo D

Influence of root structure on root survivorship: an analysis of 18 tree species using a minirhizotron method 755

Beckmann M, Bruelheide H, Erfmeier A

Germination responses of three grassland species differ between native and invasive origins 763

Schmitt FG, Devreker D, Dur G, Souissi S

Direct evidence of tidally oriented behavior of the copepod *Eurytemora affinis* in the Seine estuary 773

Tinoco Torres R, Carvalho JC, Panzacchi M, Linnell JDC, Fonseca C

Comparative use of forest habitats by roe deer and moose in a human-modified landscape in southeastern Norway during winter 781

Bekku YS, Sakata T, Tanaka T, Nakano T

Midday depression of tree root respiration in relation to leaf transpiration 791

Akamatsu F, Toda H

Flow regime alters body size but not the use of aquatic subsidies in a riparian predatory arthropod 801

Hata H, Umezawa Y

Food habits of the farmer damselfish *Stegastes nigricans* inferred by stomach content, stable isotope, and fatty acid composition analyses 809

Zhang W-P, Jia X, Bai Y-Y, Wang G-X

The difference between above- and below-ground self-thinning lines in forest communities 819

Lescano MN, Farji-Brener AG

Exotic thistles increase native ant abundance through the maintenance of enhanced aphid populations 827

Chikaraishi Y, Ogawa NO, Doi H, Ohkouchi N

¹⁵N/¹⁴N ratios of amino acids as a tool for studying terrestrial food webs: a case study of terrestrial insects (bees, wasps, and hornets) 835

Castillo DF, Lucherini M, Casanave EB

Denning ecology of Molina's hog-nosed skunk in a farmland area in the Pampas grassland of Argentina 845

Magrath A, Guitián J, Larrinaga AR, Guitián J

Land-use and edge effects unbalance seed dispersal and predation interactions under habitat fragmentation 851