

CONTENTS

Volume 159 Issue 1 2011

Review

- 1 Potential for childhood lead poisoning in the inner cities of Australia due to exposure to lead in soil dust**
M.A.S. Laidlaw, M.P. Taylor

Previous use of Pb in gasoline and Pb in exterior paints in Australia has contaminated urban soils in the older inner suburbs of large cities and the risks remain unconstrained.

Regular Papers

- 10 Habitat-specific bioaccumulation of methylmercury in invertebrates of small mid-latitude lakes in North America**
J. Chételat, M. Amyot, E. Garcia

Methylmercury levels in dominant invertebrate prey for fish differ between littoral and pelagic habitats within a lake.

- 18 Use of the MicroResp™ method to assess pollution-induced community tolerance to metals for lotic biofilms**
A. Tlili, M. Marechal, B. Montuelle, B. Volat, U. Dorigo, A. Bérard

A modified MicroResp™ technique as a tool for measuring induced tolerance to heavy metals of a microbial biofilm community.

- 25 Influence of arbuscular mycorrhizal fungi on the growth and nutrient status of bermudagrass grown in alkaline bauxite processing residue**
A. Giridhar Babu, M. Sudhakara Reddy

Inoculation of red mud tolerant AM fungi enhanced the growth and nutrient status of bermudagrass and the physico-chemical properties of the bauxite residues amended with gypsum or sewage sludge.

- 30 Root porosity and radial oxygen loss related to arsenic tolerance and uptake in wetland plants**
H. Li, Z.H. Ye, Z.J. Wei, M.H. Wong

Wetland plants with high rates of ROL tended to form more Fe plaque on root surfaces and possess higher As tolerance.

- 38 Molecular effects and bioaccumulation of levonorgestrel in the non-target organism *Dreissena polymorpha***
V. Contardo-Jara, C. Lorenz, S. Pflugmacher, G. Nützmann, W. Kloas, C. Wiegand

Fundamental cell processes as biotransformation, elimination and prevention from oxidative stress are influenced by exposure of the contraceptive levonorgestrel in non-target organisms.

- 45 The physiological, transcriptional and genetic responses of an ozone-sensitive and an ozone tolerant poplar and selected extremes of their F₂ progeny**
N.R. Street, T.M. James, T. James, B. Mikael, K. Jaakko, B. Mark, G. Taylor

*Ozone-responsive transcriptional changes and genetic control were studied in *Populus* plants with contrasting ozone sensitivity.*

- 55 Responses of evergreen and deciduous *Quercus* species to enhanced ozone levels**
V. Calatayud, J. Cerveró, E. Calvo, F.-J. García-Breijo, J. Reig-Armiñana, M.J. Sanz

*Ozone tolerance overlapped with leaf traits in four *Quercus* species.*

- 64 Is black carbon a better predictor of polycyclic aromatic hydrocarbon distribution in soils than total organic carbon?**
T. Agarwal, T.D. Bucheli

Light PAH correlated with TOC in background soils, whereas heavy PAH were associated with BC close to emission sources.

- 71 Accumulation and maternal transfer of polychlorinated biphenyls in Steller Sea Lions (*Eumetopias jubatus*) from Prince William Sound and the Bering Sea, Alaska**
J. Wang, K. Hülck, S.-M. Hong, S. Atkinson, Q.X. Li

PCBs at median concentrations of 1.2–3.7 µg/g lipid weight in different tissues of the western stock of Steller sea lions have physiological effects.

Continued on inside back cover

ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 78 Enhanced multi-metal extraction with EDDS of deficient and excess dosages under the influence of dissolved and soil organic matter**
D.Y.S. Yan, I.M.C. Lo
This study focuses on the effect of dissolved and soil organic matter on EDDS enhanced metal extraction under both deficient and excess dosage conditions.
- 84 The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants**
F. Zeng, S. Ali, H. Zhang, Y. Ouyang, B. Qiu, F. Wu, G. Zhang
Soil pH and organic matter content significantly affect heavy metal availability and accumulation in rice plants.
- 92 Eighty-year sedimentary record of heavy metal inputs in the intertidal sediments from the Nanliu River estuary, Beibu Gulf of South China Sea**
P. Xia, X. Meng, P. Yin, Z. Cao, X. Wang
The enrichment factor was successfully used to identify the different sources of heavy metals derived from natural, maricultural and agricultural inputs.
- 100 Transcriptional response of stress genes to metal exposure in zebra mussel larvae and adults**
A. Navarro, M. Faria, C. Barata, B. Piña
Coordinated expression of stress genes in zebra mussel.
- 108 Sediment quality guidelines for copper and zinc for filter-feeding estuarine oysters?**
G.F. Birch, T.D. Hogg
A strong correlation between oyster tissue Cu and Zn concentrations and fine-fraction surficial sediment digested in 1M HCl provided a sedimentary guideline which predicted tissue metal concentrations in oysters and established a level for protecting oysters from exceeding human consumption levels for these metals. The guidelines for fine surficial sediment developed in the present work accurately predicted oyster tissue concentrations in the field.
- 116 Assessing the nature of the combined effects of copper and zinc on estuarine infaunal communities**
A. Fukunaga, M.J. Anderson, J.G. Webster-Brown
The nature of the combined effects of copper and zinc depends on species and the particular response variable being examined.
- 125 Waste water treatment plants as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air**
I. Weinberg, A. Dreyer, R. Ebinghaus
Waste water treatment plants can be regarded as sources of musk fragrances, polyfluorinated compounds (PFCs) and polybrominated diphenyl ethers (PBDEs) to the atmosphere
- 133 Transport of polycyclic aromatic hydrocarbons in highly vulnerable karst systems**
K. Schwarz, T. Gocht, P. Grathwohl
Particle-associated transport is the dominant process for groundwater pollution with polycyclic aromatic hydrocarbons in remote areas.
- 140 Assessing the recovery potential of alpine moss-sedge heath: Reciprocal transplants along a nitrogen deposition gradient**
H.F. Armitage, A.J. Britton, S.J. Woodin, R. van der Wal
*Alpine moss-sedge heath, dominated by *Racomitrium lanuginosum*, from across the UK has the potential to recover from the impacts of N pollution; however, this is constrained by persistence of enhanced moss tissue N contents.*
- 148 Assessing estrogenic activity in surface water and sediment of the Liao River system in northeast China using combined chemical and biological tools**
L. Wang, G.-G. Ying, J.-L. Zhao, S. Liu, B. Yang, L.-J. Zhou, R. Tao, H.-C. Su
Estrogenic risks to aquatic organisms were assessed by using combined chemical analysis and in vitro bioassay.
- 157 Plants' use of different nitrogen forms in response to crude oil contamination**
M. Nie, M. Lu, Q. Yang, X.-D. Zhang, M. Xiao, L.-F. Jiang, J. Yang, C.-M. Fang, J.-K. Chen, B. Li
Plant strategies of utilizing nitrogen in crude oil-contaminated soils.
- 164 Antagonism of phenanthrene cytotoxicity for human embryo lung fibroblast cell line HFL-I by green tea polyphenols**
X. Mei, Y.-Y. Wu, X. Mao, Y.-Y. Tu
Green tea polyphenols antagonised cytotoxicity of a low-ring PAH phenanthrene.
- 169 Distribution of perfluorochemicals between sera and milk from the same mothers and implications for prenatal and postnatal exposures**
S.-K. Kim, K.T. Lee, C.S. Kang, L. Tao, K. Kannan, K.-R. Kim, C.-K. Kim, J.S. Lee, P.S. Park, Y.W. Yoo, J.Y. Ha, Y.-S. Shin, J.-H. Lee
Functional group and -CF₂ chain-length dependent transfer of PFCs from maternal serum to cord serum and to breast milk was clearly observed.
- 175 Metal and metalloid contamination in roadside soil and wild rats around a Pb-Zn mine in Kabwe, Zambia**
S.M.M. Nakayama, Y. Ikenaka, K. Hamada, K. Muzandu, K. Choongo, H. Teraoka, N. Mizuno, M. Ishizuka
The area around Kabwe, Zambia is highly polluted with metals and As. Wild rats from this area had high tissue concentrations of Pb and decreased body weight.
- 182 Plant communities in relation to flooding and soil contamination in a lowland Rhine River floodplain**
A.M. Schipper, K. Lotterman, R.S.E.W. Leuven, A.M.J. Ragas, H. de Kroon, A.J. Hendriks
Multiple contaminants and periodic flooding may pose cumulative stress to plants in lowland floodplains.

ENVIRONMENTAL POLLUTION

CONTENTS—Continued from inside back cover

- 190 Body metal concentrations and glycogen reserves in earthworms (*Dendrobaena octaedra*) from contaminated and uncontaminated forest soil**
M. Holmstrup, J.G. Sørensen, J. Overgaard, M. Bayley, A.-M. Bindesbøl, S. Slotsbo, K.V. Fisker, K. Maraldo, D. Waagner, R. Labouriau, G. Asmund
Detoxification and accumulation of cadmium and lead by earthworms carries little energetic expenses whereas strict internal regulation of aluminium and nickel has energetic costs.
- 198 Effects of C₆₀ nanoparticle exposure on earthworms (*Lumbricus rubellus*) and implications for population dynamics**
M.J.C. van der Ploeg, J.M. Baveco, A. van der Hout, R. Bakker, I.M.C.M. Rietjens, N.W. van den Brink
*C₆₀ nanoparticle exposure can affect *Lumbricus rubellus* populations.*
- 204 Behaviour of arsenic in forested catchments following a high-pollution period**
M. Novak, L. Erbanova, D. Fottova, P. Cudlin, A. Kubena
Following a period of high atmospheric As deposition, a considerable proportion of old industrial arsenic is flushed out of soil and exported from forested catchments.
- 212 Bisphenol A induces a rapid activation of Erk1/2 through GPR30 in human breast cancer cells**
S. Dong, S. Terasaka, R. Kiyama
We showed here that the mechanism by which BPA induces the activation of Erk1/2 is distinguishable from the mechanism of ER α -mediated signaling in human breast cancer cells.
- 219 History of mercury use and environmental contamination at the Oak Ridge Y-12 Plant**
S.C. Brooks, G.R. Southworth
Mercury discharges from an industrial plant have created a legacy contamination problem exhibiting complex and at times counter-intuitive patterns in Hg cycling.
- 229 Dietary exposure of juvenile common sole (*Solea solea* L.) to polybrominated diphenyl ethers (PBDEs): Part 1. Bioaccumulation and elimination kinetics of individual congeners and their debrominated metabolites**
C. Munsch, K. Héas-Moisan, C. Tixier, N. Olivier, O. Gastineau, N. Le Bayon, V. Buchet
Juvenile sole exposed to artificially-contaminated food accumulate PBDEs, including the higher brominated congeners, and have a debromination capacity.
- 238 Plant uptake and dissipation of PBDEs in the soils of electronic waste recycling sites**
H. Huang, S. Zhang, P. Christie
Weathered PBDEs can be taken up by plants and lower brominated PBDEs are associated with higher bioavailability to plants and a higher dissipation in soils.
- 244 The functional and physiological status of *Gammarus fossarum* (Crustacea; Amphipoda) exposed to secondary treated wastewater**
M. Bundschuh, J.P. Zubrod, R. Schulz
Micropollutants in wastewater cause functional and physiological alteration in a leaf-shredding amphipod.
- 250 Environmental quality of mussel farms in the Vigo estuary: Pollution by PAHs, origin and effects on reproduction**
Y. Ruiz, P. Suarez, A. Alonso, E. Longo, A. Villaverde, F. San Juan
Physiological parameters of sentinel organisms provides information about potentially negative effects of pollutants on aquaculture farms productivity and human food safety.
- 266 Synthesis of isotopically modified ZnO nanoparticles and their potential as nanotoxicity tracers**
A.D. Dybowska, M.-N. Croteau, S.K. Misra, D. Berhanu, S.N. Luoma, P. Christian, P. O'Brien, E. Valsami-Jones
ZnO nanoparticles with distinct isotopic composition can be tailor synthesized to be used as tracers of environmental fate and uptake by organisms.
- 274 Tissue-specific incorporation and genotoxicity of different forms of tritium in the marine mussel, *Mytilus edulis***
B.C. Jaeschke, G.E. Millward, A.J. Moody, A.N. Jha
Tritium from tritiated glycine demonstrates greater accumulation and persistence in tissues and enhanced genotoxicity in haemocytes of marine mussels, compared to tritium from tritiated water.
- 281 Sensitivity of *Bidens laevis* L. to mutagenic compounds. Use of chromosomal aberrations as biomarkers of genotoxicity**
D.J. Pérez, G. Lukaszewicz, M.L. Menone, E.L. Camadro
*The wetland macrophyte *Bidens laevis* is sensitive to genotoxic compounds similarly to terrestrial standardized species.*
- 287 Spatial distribution and seasonal variation of atmospheric bulk deposition of polycyclic aromatic hydrocarbons in Beijing–Tianjin region, North China**
W. Wang, S.L. Massey Simonich, B. Giri, M. Xue, J. Zhao, S. Chen, H. Shen, G. Shen, R. Wang, J. Cao, S. Tao
The spatial distribution and seasonal variation of PAHs deposition in Beijing–Tianjin region were studied and quantitatively related to PAHs emission density and ambient air concentration.
- 294 Carbon isotope signature of dissolved inorganic carbon (DIC) in precipitation and atmospheric CO₂**
M. Górka, P.E. Sauer, D. Lewicka-Szczepak, M.-O. Jędrysek
Carbon isotopic composition of DIC in precipitation is not in equilibrium with atmospheric CO₂ in an urban area.

CONTENTS – Continued from page IV

302 The definition of species richness used by species sensitivity distributions approximates observed effects of salinity on stream macroinvertebrates

B.J. Kefford, R. Marchant, R.B. Schäfer, L. Metzeling, J.E. Dunlop, S.C. Choy, P. Goonan

Definitions of species richness inherit in SSDs and biomonitoring are for salinity in south-east Australia commensurable.

311 New tabletop SEM-EDS-based approach for cost-efficient monitoring of airborne particulate matter

K. Wilkinson, J. Lundkvist, G. Seisenbaeva, V. Kessler

Demonstration of method for novel scanning electron microscopy in environmental monitoring of chemical constituents of particulate airborne impurities.

319 Acetylcholinesterase activity in the terrestrial snail *Xeropicta derbentina* transplanted in apple orchards with different pesticide management strategies

C. Mazzia, Y. Capowiez, J.C. Sanchez-Hernandez, H.-R. Köhler, R. Triebskorn, M. Rault

Snails as sentinel species to evaluate insecticide impacts in apple orchards.

324 Compartmentation of metals in foliage of *Populus tremula* grown on soils with mixed contamination. I. From the tree crown to leaf cell level

P. Vollenweider, T. Menard, M.S. Günthardt-Goerg

Within foliage of poplars growing on contaminated soils, Zinc was stored at metabolically safe as well as sensitive subcellular sites, ensuring sizable bioaccumulation but also causing injuries.

337 Compartmentation of metals in foliage of *Populus tremula* grown on soils with mixed contamination. II. Zinc binding inside leaf cell organelles

P. Vollenweider, P. Bernasconi, H.-P. Gautschi, T. Menard, B. Frey, M.S. Günthardt-Goerg

Zinc contaminants translocated to symplast of aged leaves were detoxified by phytic acid ligands.

Corrigendum

348 Corrigendum to “Effects of temperature and fertilization on total vs. active bacterial communities exposed to crude and diesel oil pollution in NW Mediterranean Sea” [Environ. Pollut. 158 (2010) 663–673]

A. Rodríguez-Blanco, V. Antoine, E. Pelletier, D. Delille, J.-F. Ghiglione