

Contents

(Abstracts/contents list published in *Biological Abstracts, Biological & Agricultural Index, Current Advances in Biological Sciences, Current Contents AB & ES, Ecological Abstracts, EMBiology, Environmental Abstracts, Field Crop Abstracts, Geo Abstracts, Geobase, Geographical Abstracts–Economic Geography, TROPAG and RURAL database*). Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®.

Review

| | |
|--|-----|
| Variance–covariance structure and its influence on variety assessment in regional crop trials X. Hu and J. Spilke (China, Germany) | 1 |
| Short-season cotton (<i>Gossypium hirsutum</i>) may be a suitable response to late planting in sub-Saharan regions T.-V. Cao, P. Oumarou, G. Gawrysiak, C. Klassou and B. Hau (France, Cameroon) | 9 |
| Genotype-by-environment interaction of barley DH lines infected with <i>Fusarium culmorum</i> (W.G.Sm.) Sacc. T. Warzecha, T. Adamski, Z. Kaczmarek, M. Surma, J. Chełkowski, H. Wiśniewska, K. Krystkowiak and A. Kuczyńska (Poland) | 21 |
| Quantification of fructan concentration in grasses using NIR spectroscopy and PLSR N. Shetty and R. Gislum (Denmark) | 31 |
| Sweet sorghum productivity for biofuels under increased soil salinity and reduced irrigation I. Vasilakoglou, K. Dhima, N. Karagiannidis and T. Gatsis (Greece) | 38 |
| Dryland maize yields and water use efficiency in response to tillage/crop stubble and nutrient management practices in China X. Wang, K. Dai, D. Zhang, X. Zhang, Y. Wang, Q. Zhao, D. Cai, W.B. Hoogmoed and O. Oenema (China, The Netherlands) ... | 47 |
| The effect of tillage, crop rotation and residue management on maize and wheat growth and development evaluated with an optical sensor N. Verhulst, B. Govaerts, V. Nelissen, K.D. Sayre, J. Crossa, D. Raes and J. Deckers (Mexico, Belgium) | 58 |
| Evaluation of yield performance in rice near-isogenic lines with increased spikelet number A. Ohsumi, T. Takai, M. Ida, T. Yamamoto, Y. Arai-Sanoh, M. Yano, T. Ando and M. Kondo (Japan) | 68 |
| Increased productivity through integrated soil fertility management in cassava–legume intercropping systems in the highlands of Sud-Kivu, DR Congo P. Pypers, J.-M. Sanginga, B. Kasereka, M. Walangululu and B. Vanlauwe (Kenya, Democratic Republic of Congo) | 76 |
| Combining ability analysis for <i>Phaeosphaeria</i> leaf spot resistance and grain yield in tropical advanced maize inbred lines J. Sibiyi, P. Tongoona, J. Derera, N. van Rij and I. Makanda (South Africa, Kenya) | 86 |
| Improving in-season nitrogen recommendations for maize using an active sensor J. Schmidt, D. Beegle, Q. Zhu and R. Sripada (USA) | 94 |
| Potential and cost of carbon sequestration in Indian agriculture: Estimates from long-term field experiments H. Pathak, K. Byjesh, B. Chakrabarti and P.K. Aggarwal (India) | 102 |
| Effect of water management on dry seeded and puddled transplanted rice. Part 1: Crop performance Sudhir-Yadav, G. Gill, E. Humphreys, S.S. Kukal and U.S. Walia (Australia, Philippines, India) | 112 |

(Contents continued)

| | |
|---|-----|
| Effect of water management on dry seeded and puddled transplanted rice. Part 2: Water balance and water productivity Sudhir-Yadav, E. Humphreys, S.S. Kukal, G. Gill and R. Rangarajan (Australia, Philippines, India) | 123 |
| High-yield irrigated maize in the Western U.S. Corn Belt: II. Irrigation management and crop water productivity P. Grassini, H. Yang, S. Irmak, J. Thorburn, C. Burr and K.G. Cassman (USA) | 133 |
| High-yield irrigated maize in the Western U.S. Corn Belt: I. On-farm yield, yield potential, and impact of agronomic practices P. Grassini, J. Thorburn, C. Burr and K.G. Cassman (USA) | 142 |
| Post-flowering leaflet removals increase pod initiation in soybean canopies A. Quijano and E.N. Morandi (Argentina) | 151 |
| The potential contribution of wild barley (<i>Hordeum vulgare</i> ssp. <i>spontaneum</i>) germplasm to drought tolerance of cultivated barley (<i>H. vulgare</i> ssp. <i>vulgare</i>) B. Lakew, J. Eglinton, R.J. Henry, M. Baum, S. Grando and S. Ceccarelli (Ethiopia, Australia, Syria) | 161 |
| Determinants of barley grain yield in a wide range of Mediterranean environments E. Francia, A. Tondelli, F. Rizza, F.W. Badeck, O. Li Destri Nicosia, T. Akar, S. Grando, A. Al-Yassin, A. Benbelkacem, W.T.B. Thomas, F. van Eeuwijk, I. Romagosa, A.M. Stanca and N. Pecchioni (Italy, Germany, Turkey, Syria, Jordan, Algeria, UK, The Netherlands, Spain) | 169 |
| Using CERES-Wheat to simulate durum wheat production and phenology in Southern Sardinia, Italy M. Dettori, C. Cesaraccio, A. Motroni, D. Spano and P. Duce (Italy) | 179 |
| Tillage system effect on nitrogen rhizodeposited by faba bean and chickpea L. López-Bellido, J. Benítez-Vega, P. García, R. Redondo and R.J. López-Bellido (Spain) | 189 |
| Ear rot, aflatoxin accumulation, and fungal biomass in maize after inoculation with <i>Aspergillus flavus</i> W.P. Williams, S. Ozkan, A. Ankala and G.L. Windham (USA) | 196 |
| Erratum | |
| Erratum to “The effect of various dynamics of N availability on winter pea–wheat intercrops: Crop growth, N partitioning and symbiotic N ₂ fixation” [Field Crops Res. 119 (2010) 2–11] C. Naudin, G. Corre-Hellou, S. Pineau, Y. Crozat and M.-H. Jeuffroy (France) | 201 |