

Food Hydrocolloids

CONTENTS

Volume 25, Issue 3, 2011

- 275 Combined microscopic and dynamic rheological methods for studying the structural breakdown properties of whey protein gels and emulsion filled gels
A. R. ABHYANKAR, D. M. MULVIHILL AND M. A. E. AUTY (Ireland)
- 283 Determination of the upper temperature limit of water loss by food systems
YU. I. MATVEEV (Russian Federation)
- 286 Molecular characteristics of a novel water-soluble polysaccharide from the New Zealand black tree fern (*Cyathea medullaris*)
K. K. THA GOH, L. MATIA-MERINO, D. N. PINDER, C. SAAVEDRA AND H. SINGH (New Zealand)
- 293 Rheological properties of gum arabic solution: From Newtonianism to thixotropy
X. LI (China), Y. FANG, H. ZHANG (UK), K. NISHINARI (Japan), S. AL-ASSAF AND G. O. PHILLIPS (UK)
- 299 Rheological behavior of blueberry (*Vaccinium ashei*) purees containing xanthan gum and fructose as ingredients
C. P. KECHINSKI, A. B. SCHUMACHER, L. D. F. MARCZAK, I. C. TESSARO AND N. S. M. CARDOZO (Brazil)
- 307 Encapsulation systems based on ovalbumin fibrils and high methoxyl pectin
K. N. P. HUMBLET-HUA, G. SCHELTENS, E. VAN DER LINDEN AND L. M. C. SAGIS (The Netherlands)
- 315 A comparative study of physicochemical and conformational properties in three vicilins from *Phaseolus* legumes: Implications for the structure–function relationship
C.-H. TANG AND X. SUN (PR China)
- 325 The determining role of bolus rheology in triggering a swallowing
J. CHEN AND L. LOLIVRET (UK)
- 333 New insights on the formation of colloidal whey protein particles
L. E. VAN RIEMSDIJK, J. P. M. SNOEREN, A. J. VAN DER GOOT, R. M. BOOM AND R. J. HAMER (The Netherlands)
- 340 Rheology and breadmaking performance of rice-buckwheat batters supplemented with hydrocolloids
D. PERESSINI, M. PIN AND A. SENSIDONI (Italy)
- 350 Structure formation and phase-separation behaviour of aqueous casein–alginate emulsions in the presence of strong polyelectrolyte
Y. A. ANTONOV (Russia) AND P. MOLDENAERS (Belgium)
- 361 Micellar properties of OSA starch and interaction with xanthan gum in aqueous solution
V. KRSTONOŠIĆ, L. DOKIĆ AND J. MILANOVIĆ (Serbia)
- 368 Study on the properties of gelatins from skin of carp (*Cyprinus carpio*) caught in winter and summer season
R. DUAN, J. ZHANG, F. XING, K. KONNO (Japan) AND B. XU (China)
- 374 Partitioning of α -lactalbumin and β -lactoglobulin in whey protein concentrate/hydroxypropylmethylcellulose aqueous two-phase systems
F. JARA AND A. M. R. PILOSOF (Argentina)

Contents continued on BM II

- 381 Characteristics of gelatin from the skin of unicorn leatherjacket (*Aluterus monoceros*) as influenced by acid pretreatment and extraction time
M. AHMAD AND S. BENJAKUL (Thailand)
- 389 β -Conglycinin and glycinin soybean protein emulsions treated by combined temperature–high-pressure treatment
M. C. PUPPO (Argentina), V. BEAUMAL (France), F. SPERONI (Argentina), M. DE LAMBALLERIE (France), M. C. AÑÓN (Argentina) AND M. ANTON (France)
- 398 Freeze-thaw stability of oil-in-water emulsions prepared with native and thermally-denatured soybean isolates
G. G. PALAZOLO, P. A. SOBRAL AND J. R. WAGNER (Argentina)
- 410 Processing stability of cross-linked starches in acid sauce applications and identification of some of the molecular factors involved
P. A. M. STEENEKEN, A. J. J. WOORTMAN AND A. A. C. M. (LIZETTE) OUDHUIS (The Netherlands)
- 419 Charge domain of modified pectins influence interaction with acidified caseins
Y. KIM (South Korea) AND L. WICKER (USA)
- 426 Roles of lipid oxidation and pH on properties and yellow discolouration during storage of film from red tilapia (*Oreochromis niloticus*) muscle protein
P. TONGNUANCHAN, S. BENJAKUL AND T. PRODPRAN (Thailand)
- 434 Quantifying structural characteristics of partially de-esterified pectins
D. E. NGOUÉMAZONG, F. F. TENGWEH, T. DUVETTER, I. FRAEYE, A. VAN LOEY, P. MOLDENAERS AND M. HENDRICKX (Belgium)
- 444 Rheological properties of extracted malva nut gum (*Scaphium scaphigerum*) in different conditions of solvent
A. SRICHAMROEN AND V. CHAVASIT (Thailand)
- 451 Steady shear flow properties of wild sage (*Salvia macrosiphon*) seed gum as a function of concentration and temperature
S. M. A. RAZAVI, H. TAHERI AND L. A. QUINCHIA (Iran, Spain)
- 459 The effect of protein types and low molecular weight surfactants on spray drying of sugar-rich foods
M. JAYASUNDERA, B. ADHIKARI, R. ADHIKARI AND P. ALDRED (Australia)
- 470 Evaluation of edible film-forming ability of pumpkin oil cake; effect of pH and temperature
S. POPOVIĆ, D. PERIČIN, Ž. VAŠTAG, L. POPOVIĆ AND V. LAZIĆ (Serbia)
- 477 Long- and short-range structural changes of recrystallised cassava starch subjected to *in vitro* digestion
C. MUTUNGI, C. ONYANGO, T. DOERT, S. PAASCH, S. THIELE, S. MACHILL, D. JAROS AND H. ROHM (Germany)
- 486 Effects of molecular characteristics of tea polysaccharide in green tea on glass transitions of potato amylose, amylopectin and their mixtures
L. GUO, Q. LIANG AND X. DU (China)
- 495 Droplet deformation and break-up under shear: Hydrocolloid solution vs. suspension of starch granules
M. DESSE, J. MITCHELL, B. WOLF AND T. BUDTOVA (France, UK)
- 503 On quantifying the dissolution behaviour of milk protein concentrate
Y. FANG, C. SELOMULYA, S. AINSWORTH, M. PALMER AND X. D. CHEN (Australia)
- 511 Study of gelatin renaturation in aqueous solution by AFIFFF–MALS: Influence of a thermal pre-treatment applied on gelatin
K. RBII, O. SUREL, N. BRAMBATI, A.-M. BUCHERT AND F. VIOLLEAU (France)
- 515 Effect of succinylation on the physicochemical, rheological, thermal and retrogradation properties of red and white sorghum starches
O. O. OLAYINKA, B. I. OLU-OWOLABI AND K. O. ADEBOWALE (Nigeria)
- 521 Characteristics of native and enzymatically hydrolyzed *Zea mays* L., *Fritillaria ussuriensis* Maxim. and *Dioscorea opposita* Thunb. starches
Q. JIANG, W. GAO, X. LI AND J. ZHANG (China)
- 529 Antioxidant capacity of sulfated polysaccharides from seaweeds. A kinetic approach
T. BARAHONA, N. P. CHANDÍA, M. V. ENCINAS, B. MATSUHIRO AND E. A. ZÚÑIGA (Chile)
- 536 Structure–physicochemical function relationships of 7S globulins (vicilins) from red bean (*Phaseolus angularis*) with different polypeptide constituents
C.-H. TANG AND X. SUN (PR China)
- 545 Enhancement of anti-radical activity of pectin from apple pomace by hydroxamation
H. J. RHA, I. Y. BAE, S. LEE, S.-H. YOO, P.-S. CHANG AND H. G. LEE (Republic of Korea)
- 549 Bibliography
- 555 Calendar