

■ Data Papers

Takamura N, Nakagawa M

Phytoplankton species abundance in Lake Kasumigaura (Japan) monitored monthly or biweekly since 1978 837

Takamura N, Nakagawa M

The densities of bacteria, picophytoplankton, heterotrophic nanoflagellates and ciliates in Lake Kasumigaura (Japan) monitored monthly since 1996 839

■ Original Articles

Malíková L, Mudrák O, Klimešová J

Adventitious sprouting enables the invasive annual herb *Euphorbia geniculata* to regenerate after severe injury 841

Kobayashi K, Sato Y, Kaji K

Increased brown bear predation on sika deer fawns following a deer population irruption in eastern Hokkaido, Japan 849

Abou Jaoudé R, de Dato G, De Angelis P

Photosynthetic and wood anatomical responses of *Tamarix africana* Poiret to water level reduction after short-term fresh- and saline-water flooding 857

Zheng Y-L, Feng Y-L, Wang R-F, Shi X-D, Lei Y-B, Han L-H

Invasive *Eupatorium adenophorum* suffers lower enemy impact on carbon assimilation than native congeners 867

Bezemer TM, Jones TH

The effects of CO₂ and nutrient enrichment on photosynthesis and growth of *Poa annua* in two consecutive generations 873

Cao J, Wang X, Tian Y, Wen Z, Zha T

Pattern of carbon allocation across three different stages of stand development of a Chinese pine (*Pinus tabulaeformis*) forest 883

Suzuki SN, Kachi N, Suzuki J-I

Variability of local spatial structure in a wave-regenerated *Abies* forest 893

Pirk GI, Farji-Brener AG

Foliar herbivory and its effects on plant growth in native and exotic species in the Patagonian steppe 903

Lucas-Borja ME, Fonseca TF, Lousada JL, Silva-Santos P, Garcia EM, Abellán MA

Natural regeneration of Spanish black pine [*Pinus nigra* Arn. ssp. *salzmannii* (Dunal) Franco] at contrasting altitudes in a Mediterranean mountain area 913

Chen F-S, Feng X, Liang C

Endogenous versus exogenous nutrient affects C, N, and P dynamics in decomposing litters in mid-subtropical forests of China 923

Bontempo e Silva EA, Hasegawa SF, Ono K, Sumida A, Uemura S, Hara T

Differential photosynthetic characteristics between seedlings and saplings of *Abies sachalinensis* and *Picea glehnii*, in the field 933

Jonsson M, Strasevicius D, Malmqvist B

Influences of river regulation and environmental variables on upland bird assemblages in northern Sweden 945

Lu T, Zeng H, Luo Y, Wang Q, Shi F, Sun G, Wu Y, Wu N

Monitoring vegetation recovery after China's May 2008 Wenchuan earthquake using Landsat TM time-series data: a case study in Mao County 955

De Crop E, Brys R, Hoffmann M

The impact of habitat fragmentation on the interaction between *Centaurium erythraea* (Gentianaceae) and its specialized seed predator *Stenoptilia zophodactylus* (Pterophoridae, Lepidoptera) 967

Cárdenas RE, Dangles O

Do canopy herbivores mechanically facilitate subsequent litter decomposition in soil? A pilot study from a Neotropical cloud forest 975

■ Note and Comment

Striker GG

Time is on our side: the importance of considering a recovery period when assessing flooding tolerance in plants 983