

CONTENTS

Volume 171 December 2012

- 1 Temporal and spatial trends of perfluoroalkyl substances in liver of roe deer (*Capreolus capreolus*)**
S. Falk, H. Brunn, C. Schröter-Kermani, K. Failing, S. Georgii, K. Tarricone, T. Stahl
This is the first study to investigate temporal and spatial trends of PFASs in a terrestrial herbivore mammal.
- 9 Development of a distributed air pollutant dry deposition modeling framework**
S. Hirabayashi, C.N. Kroll, D.J. Nowak
Employing nationally available datasets and a GIS, this study will provide urban forest managers in U.S. cities a framework to quantify and visualize urban forest structure and its air pollution removal effect.
- 18 Environmental contextualisation of potential toxic elements and polycyclic aromatic hydrocarbons in biochar**
A. Freddo, C. Cai, B.J. Reid
Environmental impacts attributable to metals, metalloids and PAHs associated with biochar following its application to soil are likely to be minimal.
- 25 Physiological response of the epiphytic lichen *Evernia prunastri* (L.) Ach. to ecologically relevant nitrogen concentrations**
S. Munzi, L. Paoli, E. Fiorini, S. Loppi
*Physiological parameters of the lichen *Evernia prunastri* respond to ecologically relevant concentrations of nitrogen compounds.*
- 30 The long-term impact of urbanization on nitrogen patterns and dynamics in Shanghai, China**
B. Gu, X. Dong, C. Peng, W. Luo, J. Chang, Y. Ge
Major nitrogen emission source shifts from agriculture to industry and human, and decoupling of nitrogen with GDP and urban expansion arises alongside urbanization.
- 38 Summer–winter concentrations and gas-particle partitioning of short chain chlorinated paraffins in the atmosphere of an urban setting**
T. Wang, S. Han, B. Yuan, L. Zeng, Y. Li, Y. Wang, G. Jiang
Atmospheric levels and gas-particle partitioning of SCCPs in Beijing, China.
- 46 Physiological and molecular responses of the earthworm (*Eisenia fetida*) to soil chlortetracycline contamination**
D. Lin, Q. Zhou, Y. Xu, C. Chen, Y. Li
Chlortetracycline in soil could induce physiological responses and genotoxicity on earthworms at realistic environmental concentrations.
- 52 Assessing the significance of *Ruditapes philippinarum* as a sentinel for sediment pollution: Bioaccumulation and biomarker responses**
V. Moschino, E. Delaney, L. Da Ros
*Biological and chemical results on *Ruditapes philippinarum* used as a bioindicator suggested its cautious use in biomonitoring transitional environments.*
- 61 Lipid and fatty acid composition, and persistent organic pollutant levels in tissues of migrating Atlantic bluefin tuna (*Thunnus thynnus*, L.) broodstock**
M. Sprague, J.R. Dick, A. Medina, D.R. Tocher, J.G. Bell, G. Mourente
Lipids, fatty acids (particularly omega-3) and POPs were determined in Atlantic bluefin tuna to examine tissue distribution, and assess the risk-benefit to human health.
- 72 Distributions and impact factors of antimony in topsoils and moss in Ny-Ålesund, Arctic**
N. Jia, L. Sun, X. He, K. You, X. Zhou, N. Long
The distribution of Sb in Ny-Ålesund is partially associated with traffic and historical mining activities, and moss can be a useful tool for monitoring Sb in both highly and lightly polluted areas.

Continued on inside back cover

CONTENTS—Continued from outside back cover

- 78 Distribution of PCDD/Fs and organometallic compounds in sewage sludge of wastewater treatment plants in China**
M. Lu, X.-J. Wu, D.-C. Zeng, Y. Liao
This is the first study on the survey of the distributions of POPs and organometallic compounds in sewage sludge in China on a national scale.
- 85 Enrichment behavior and transport mechanism of soil-bound PAHs during rainfall-runoff events**
Y. Zheng, X. Luo, W. Zhang, B. Wu, F. Han, Z. Lin, X. Wang
The traditional enrichment theory is not fully valid for PAHs, and soil organic matters have a significant impact on the transport of PAHs during rainfall-runoff events.
- 93 Fate and bioavailability of ¹⁴C-pyrene and ¹⁴C-lindane in sterile natural and artificial soils and the influence of aging**
K. Šmídová, J. Hofman, A.E. Ite, K.T. Semple
Sterilization reduced the differences in POPs fate and bioavailability between artificial and natural soils with the same TOC content and eliminated the influence of soil contact time.
- 99 Ecological impact assessment of sediment remediation in a metal-contaminated lowland river using translocated zebra mussels and resident macroinvertebrates**
M. De Jonge, C. Belpaire, C. Geeraerts, W. De Cooman, R. Blust, L. Bervoets
Metal accumulation in selected aquatic invertebrates can be used as an integrated measure of metal bioavailability and to predict ecological effects of metal toxicity.
- 109 Were mercury emission factors for Chinese non-ferrous metal smelters overestimated? Evidence from onsite measurements in six smelters**
L. Zhang, S. Wang, Q. Wu, Y. Meng, H. Yang, F. Wang, J. Hao
Acid plants in smelters have high mercury removal efficiency, and thus mercury emission factors for Chinese non-ferrous metal smelters were probably overestimated.
- 118 Effects of elevated O₃ concentration on winter wheat and rice yields in the Yangtze River Delta, China**
X. Wang, Q. Zhang, F. Zheng, Q. Zheng, F. Yao, Z. Chen, W. Zhang, P. Hou, Z. Feng, W. Song, Z. Feng, F. Lu
The dose–response relationships derived from field fumigation experiments over 5 years can be used to accurately estimate crop losses in China.
- 126 CO₂ emissions from farm inputs “Case study of wheat production in Canterbury, New Zealand”**
M. Šafa, S. Samarasinghe
Total CO₂ emissions were 1032 kg CO₂/ha in wheat production. Around 52% of the total CO₂ emissions were released from fertilizer.
- 133 Ecotoxicity of the veterinary pharmaceutical ivermectin tested in a soil multi-species (SMS) system**
J. Jensen, J.J. Scott-Fordsmand
Hazardous substances are shown more toxic to soil dwelling invertebrates when exposed in co-existence with other species including a predator.
- 140 Airborne psychotropic substances in eight Italian big cities: Burdens and behaviours**
A. Cecinato, C. Balducci, P. Romagnoli, M. Perilli
Both licit and illicit psychotropic substances modulate in space and time along Italy, resulting influenced by both the environmental and human contours.
- 148 Factors regulating the bioavailability of methylmercury to breeding rusty blackbirds in northeastern wetlands**
S.T. Edmonds, N.J. O’Driscoll, N.K. Hillier, J.L. Atwood, D.C. Evers
Elevated mercury concentrations in the disappearing Acadian population of breeding rusty blackbirds are due to a diet of aquatic macroinvertebrates in an environment with high methylmercury bioavailability.
- 155 Polybrominated diphenyl ethers (PBDEs) in sediments of the coastal East China Sea: Occurrence, distribution and mass inventory**
Y. Li, T. Lin, Y. Chen, L. Hu, Z. Guo, G. Zhang
The levels, distribution, potential sources and mass inventory of PBDEs in sediments of the coastal East China Sea were revealed.
- 162 Morphology changes in human lung epithelial cells after exposure to diesel exhaust micron sub particles (PM_{1.0}) and pollen allergens**
V. Esposito, A. Lucariello, L. Savarese, M.P. Cinelli, F. Ferraraccio, A. Bianco, A. De Luca, G. Mazzarella
The urban environment with the combination of inhalable air pollution and particulate can damage the acinar lung units and activate cells of the immune system.
- 168 Temperature dependence of ofloxacin fluorescence quenching and complexation by Cu(II)**
B. Pan, X. Han, M. Wu, H. Peng, D. Zhang, H. Li, B. Xing
Cu(II) ions complex OFL molecules with the molar ratio of 1:1 to 1:2 on carboxylic and ketonic groups.
- 174 Evaluation through column leaching tests of metal release from contaminated estuarine sediment subject to CO₂ leakages from Carbon Capture and Storage sites**
M.C. Payán, B. Galan, A. Coz, C. Vandecasteele, J.R. Viguri
Column metal release from CO₂ acidified seawater leakages in contact with estuarine contaminated sediment in CCS sites

CONTENTS—Continued from inside back cover

185 Sorption kinetics of ofloxacin in soils and mineral particles

B. Pan, P. Wang, M. Wu, J. Li, D. Zhang, D. Xiao

Ofloxacin sorption in soils was not a diffusion process and could be modeled using a two-compartment first-order kinetics model.

191 Hexabromocyclododecane in terrestrial passerine birds from e-waste, urban and rural locations in the Pearl River Delta, South China: Levels, biomagnification, diastereoisomer- and enantiomer-specific accumulation

Y.-x. Sun, X.-j. Luo, L. Mo, M.-j. He, Q. Zhang, S.-j. Chen, F.-s. Zou, B.-x. Mai

Enantiomer-specific accumulation of hexabromocyclododecane in terrestrial passerine birds in South China was demonstrated.

199 Warm season chloride concentrations in stream habitats of freshwater mussel species at risk

A.K. Todd, M.G. Kaltenecker

Warm season chloride concentrations increased in southern Ontario streams with road salt use, such that reproduction of freshwater mussel species at risk may be affected.

207 Methylmercury in water, sediment, and invertebrates in created wetlands of Rouge Park, Toronto, Canada

K.A. Sinclair, Q. Xie, C.P.J. Mitchell

Sediment methylmercury concentrations and bioaccumulation in many invertebrate taxa are significantly elevated in newly created wetlands.

216 AhR-mediated potency of sediments and soils in estuarine and coastal areas of the Yellow Sea region: A comparison between Korea and China

S. Hong, J.S. Khim, J.E. Naile, J. Park, B.-O. Kwon, T. Wang, Y. Lu, W.J. Shim, P.D. Jones, J.P. Giesy

AhR-mediated potencies of sediments and soils in estuarine and coastal areas of the West Sea of Korea and north Bohai Sea of China was due known and unknown AhR agonists.

226 Integrated assessment on groundwater nitrate by unsaturated zone probing and aquifer sampling with environmental tracers

L. Yuan, Z. Pang, T. Huang

Denitrification loss has been evaluated in the unsaturated zone and saturated zone.

234 Summer atmospheric polybrominated diphenyl ethers in urban and rural areas of northern China

C. Wang, W. Li, J. Chen, H. Wang, T. Li, G. Shen, H. Shen, Y. Huang, R. Wang, B. Wang, Y. Zhang, J. Tang, W. Liu, X. Wang, S. Tao

In northern China, PBDEs in air in rural residential areas were significantly higher than those in open fields.

241 PBDE and PCB accumulation in benthos near marine wastewater outfalls: The role of sediment organic carbon

P.M. Dinn, S.C. Johannessen, P.S. Ross, R.W. Macdonald, M.J. Whitticar, C.J. Lowe, A. van Roodselaar

The physical receiving environment affects the accumulation of PBDEs by benthic invertebrates near submarine municipal outfalls, and uptake of PBDE congeners is governed by different properties than for PCB congeners.

249 Short-term molecular-level effects of silver nanoparticle exposure on the earthworm, *Eisenia fetida*

O.V. Tsyusko, S.S. Hardas, W.A. Shoults-Wilson, C.P. Starnes, G. Joice, D.A. Butterfield, J.M. Unrine

*Similarity in molecular-level responses between silver nanoparticles (Ag-NPs) and ions suggests that ions are primarily responsible for short-term toxicity of Ag-NPs to *Eisenia fetida*.*

256 Ecodynamics of PAHs at a peri-urban site of the French Mediterranean Sea

F. Charles, C. Nozais, A.M. Pruski, S. Bourgeois, L. Méjanelle, G. Vétion, B. Rivière, J. Coston-Guarini

We consider the nutritional quality of sinking particles and surficial sediments in connection with the flux of selected PAHs at a nearshore site of the northwestern Mediterranean Sea.

265 Modelling lead bioaccessibility in urban topsoils based on data from Glasgow, London, Northampton and Swansea, UK

J.D. Appleton, M.R. Cave, J. Wragg

Total Pb is the only significant predictor for bioaccessible Pb in topsoils from four urban areas in the UK.