

ORIGINAL ARTICLES

- J.-L. HEMPTINNE, A. MAGRO, C. SALADIN, A. F. G. DIXON
Role of intraguild predation in aphidophagous guilds 161–170
- P. REMPOULAKIS, D. NESTEL
Dispersal ability of marked, irradiated olive fruit flies [*Bactrocera oleae* (Rossi)
(Diptera: Tephritidae)] in arid regions 171–180
- M. T. MARTINEZ-FERRER, J. M. CAMPOS, J. M. FIBLA
Field efficacy of *Ceratitis capitata* (Diptera: Tephritidae) mass trapping technique on
clementine groves in Spain 181–190
- Z. L. JIANG, Y. AKHTAR, X. ZHANG, R. BRADBURY, M. B. ISMAN
Insecticidal and feeding deterrent activities of essential oils in the cabbage looper,
Trichoplusia ni (Lepidoptera: Noctuidae) 191–202
- B. KALINOVÁ, R. R. DO NASCIMENTO, M. HOSKOVEC, A. L. MENDONÇA, E. L. SILVA,
M. R. T. DE FREITAS, C. R. CABRAL-JR, C. E. SILVA, A. E. G. SANT'ANA, A. SVATOŠ
Identification of two components of the female sex pheromone of the sugarcane-borer
Diatraea flavipennella (Lepidoptera: Crambidae) 203–211
- B. WERMELINGER, C. EPPER, M. KENIS, S. GHOSH, O. HOLDENRIEDER
Emergence patterns of univoltine and bivoltine *Ips typographus* (L.) populations and
associated natural enemies 212–224
- M. KNAPP, P. SASKA
The effects of habitat, density, gender and duration on overwintering success in *Bembidion*
lampros (Coleoptera: Carabidae) 225–233
- SHORT COMMUNICATIONS**
- P. KEHRLI, N. DELABAYS
Controlling 'bois noir' disease on grapevine: does the timing of herbicide application affect
vector emergence? 234–237
- T. E. SHELLY
Mate choice by wild and mass-reared females of the Mediterranean fruit fly 238–240