

ORIGINAL ARTICLES

U. NILSSON, L.-M. RÄNNBÄCK, P. ANDERSON, B. RÄMERT

Herbivore response to habitat manipulation with floral resources: a study of the cabbage root fly

481–489

R. MOERKENS, H. LEIRS, G. PEUSENS, T. BELIËN, B. GOBIN

Natural and human causes of earwig mortality during winter: temperature, parasitoids and soil tillage

490–500

C. ROMEU-DALMAU, X. ESPADALER, J. PIÑOL

Abundance, interannual variation and potential pest predator role of two co-occurring earwig species in citrus canopies

501–509

M. Z. AHMED, P. J. DE BARRO, A. OLLEKA, S.-X. REN, N. S. MANDOUR, J. M. GREEFF, B.-L. QIU

Use of consensus sequences and genetic networks to identify members of the *Bemisia tabaci* cryptic species complex in Egypt and Syria

510–519

A. CHOW, A. CHAU, K. M. HEINZ

Reducing fertilization: a management tactic against western flower thrips on roses

520–529

Y. GÓMEZ-BONILLA, M. LÓPEZ-FERBER, P. CABALLERO, X. LÉRY, D. MUÑOZ

Costa Rican soils contain highly insecticidal granulovirus strains against *Phthorimaea operculella* and *Tecia solanivora*

530–538

Z.-X. YANG, Q.-J. WU, S.-L. WANG, X.-L. CHANG, J.-H. WANG, Z.-J. GUO, Y.-Y. LEI, B.-Y. XU, Y.-J. ZHANG

Expression of cadherin, aminopeptidase N and alkaline phosphatase genes in Cry1Ac-susceptible and Cry1Ac-resistant strains of *Plutella xylostella* (L.)

539–548

W. L. YEE

Detection of *Rhagoletis indifferens* (Dipt., Tephritidae) larvae using brown sugar flotation and hot water methods

549–560