

- 1 Use of chitosan-based edible coatings in combination with other natural compounds, to control *Rhizopus stolonifer* and *Escherichia coli* DH5 α in fresh tomatoes
M. Ramos-García, E. Bosquez-Molina, J. Hernández-Romano, G. Zavala-Padilla, E. Terrés-Rojas, I. Alía-Tejacal, L. Barrera-Necha, M. Hernández-López and S. Bautista-Baños
- 7 Effect of neemarin on life table indices of *Plutella xylostella* (L.)
N. Ahmad, M.S. Ansari and Nazrussalam
- 15 Weed age affects chemical control of *Conyza bonariensis* in fallows
S. Walker, L. Boucher, T. Cook, B. Davidson, A. McLean and M. Widderick
- 21 Shade is conducive to coffee rust as compared to full sun exposure under standardized fruit load conditions
D.F. López-Bravo, E.d.M. Virginio-Filho and J. Avelino
- 30 Effectiveness of 1,3-dichloropropene as an alternative to methyl bromide in rotations of tomato (*Solanum lycopersicum*) and cucumber (*Cucumis sativus*) in China
K. Qiao, S. Dong, H. Wang, X. Xia, X. Ji and K. Wang
- 35 Multiple Pro-197 substitutions in the acetolactate synthase of rigid ryegrass (*Lolium rigidum*) and their impact on chlorsulfuron activity and plant growth
N.S. Kaloumenos, V.C. Tsioni, E.G. Daliani, S.E. Papavassileiou, A.G. Vassileiou, P.N. Laoutidou and I.G. Eleftherohorinos
- 44 Evaluation of potato tuber moth mortality following postharvest cold storage of potatoes
G. Saour, A. Al-Daoude and H. Ismail
- 49 Effect of hail protection nets on apple scab in 'Royal Gala' and 'Fuji' apple cultivars
A. Bogo, R.T. Casa, L. Agostineto, M.J. Gonçalves and L. Rufato
- 53 The effects of water management, timing and the rate of several herbicides on the growth of *Murdannia keisak* (Hassk.) Handel-Mazz
A. Ferrero, F. Tesio, M. Tabacchi and F. Vidotto
- 57 Ecology and management of weeds under conservation agriculture: A review
B.S. Chauhan, R.G. Singh and G. Mahajan

- 66 Simple and efficient method for detection and identification of *Pyrenochaeta terrestris* on maize root
J. Lević, S. Stanković, V. Krnjaja and A. Bočarov-Stančić
- 72 Species specific primers for identification of *Alternaria solani*, in combination with analysis of the F129L substitution associates with loss of sensitivity toward strobilurins
E. Edin
- 74 When and where a seed mix refuge makes sense for managing insect resistance to Bt plants
M.W. Carroll, G. Head and M. Caprio
- 80 Control of *Botrytis cinerea*, *Alternaria alternata* and *Pyrenochaeta lycopersici* on tomato with whey compost-tea applications
C. Pane, G. Celano, D. Vilecco and M. Zaccardelli
- 87 Mass-rearing of insects for pest management: Challenges, synergies and advances from evolutionary physiology
J.G. Sørensen, M.F. Addison and J.S. Terblanche
- 95 Investigation of the predation potential of different fish species on brown planthopper (*Nilaparvata lugens* (Stål)) in experimental rice-fish aquariums and tanks
N. Cao Quoc, N. Vromant, B. Tran Thanh and F. Ollevier
- 103 The effect of ridge planting and earliness of durum wheat varieties on sunn pest (*Eurygaster* spp.) damage and grain yield
S. Gürsoy, Ç. Mutlu, M. Urğun, B. Kolay, V. Karaca and M. Duman
- 108 IAPPS Newsletter
- 111 Calendar