

ORIGINAL-ARTICLES

- M. UEFUNE, Y. CHOH, J. ABE, K. SHIOJIRI, K. SANO, J. TAKABAYASHI**
Application of synthetic herbivore-induced plant volatiles causes increased parasitism of herbivores in the field **561–567**
- J. RAZMJOU, C. VORBURGER, M. MOHAMMADI, M. HASSANPOUR**
Influence of vermicompost and cucumber cultivar on population growth of *Aphis gossypii* Glover **568–575**
- C. P. KEATHLEY, D. A. POTTER**
Arthropod abundance in tall fescue, *Lolium arundinaceum*, pastures containing novel 'safe' endophytes **576–587**
- B. L. ENZMANN, K. M. KAPHEIM, T. B. WANG, P. NONACS**
Giving them what they want: manipulating Argentine ant activity patterns with water **588–595**
- E. STOIANOVA, T. WILLIAMS, J. CISNEROS, D. MUÑOZ, R. MURILLO, E. TASHEVA, P. CABALLERO**
Interactions between an ectoparasitoid and a nucleopolyhedrovirus when simultaneously attacking *Spodoptera exigua* (Lepidoptera: Noctuidae) **596–604**
- R. P. BLACKSHAW, J. P. MOORE**
Within-generation dynamics of leatherjackets (*Tipula paludosa* Meig.) **605–613**
- T.-H. JUN, M. A. R. MIAN, K. FREEWALT, O. MITTAPALLI, A. P. MICHEL**
Development of genic-SSRs markers from soybean aphid sequences generated by high-throughput sequencing of cDNA library **614–625**
- K. T. TSOUMANI, K. D. MATHIOPOULOS**
Genome size estimation with quantitative real-time PCR in two Tephritidae species: *Ceratitis capitata* and *Bactrocera oleae* **626–631**
- SHORT COMMUNICATIONS**
- A. ZHANG, X. LI, H. ZHANG, H. WANG, L. MIAO, J. ZHANG, Q. QIN**
A new cell line from *Spodoptera exigua* (Lepidoptera: Noctuidae) and its differentially expressed genes **632–637**
- U. BERNARDO, R. SASSO, M. GEBIOLA, G. VIGGIANI**
First record of a walnut shield bearer *Coptodisca* (Lepidoptera: Heliozelidae) in Europe **638–640**