

CONTENTS

Volume 170 November 2012

- 1 Polychlorinated naphthalenes (PCNs) in the surface soils of the Pearl River Delta, South China: Distribution, sources, and air-soil exchange**
Y. Wang, Z. Cheng, J. Li, C. Luo, Y. Xu, Q. Li, X. Liu, G. Zhang
As a potential source during the hot period, low-chlorinated PCNs released from contaminated soils may transport to nearby regions.
- 8 Dynamics of CH₄ oxidation in landfill biocover soil: Effect of O₂/CH₄ ratio on CH₄ metabolism**
Z.-F. Chi, W.-J. Lu, H. Li, H.-T. Wang
A dual-substrates model of CH₄ oxidation is developed to provide a new model to control landfill CH₄ pollution to the environment.
- 15 Biogeochemical malfunctioning in sediments beneath a deep-water fish farm**
T. Valdemarsen, R.J. Bannister, P.K. Hansen, M. Holmer, A. Ervik
Siting fish farms at deep water farming locations is not a universal solution for alleviating benthic impacts.
- 26 PFOS and PFOA in influents, effluents, and biosolids of Chinese wastewater treatment plants and effluent-receiving marine environments**
H. Chen, C. Zhang, J. Han, Y. Yu, P. Zhang
Levels of PFOS and PFOA in influent, effluent and sludge samples from Chinese WWTPs were examined and found much higher than those in receiving seawater.
- 32 Protozoan predation in soil slurries compromises determination of contaminant mineralization potential**
N. Badawi, A.R. Johnsen, K.K. Brandt, J. Sørensen, J. Aamand
Protozoan predation may strongly bias the quantification of mineralization potential when performed in soil slurries, especially when the initial density of degrader bacteria is low such as in subsoil or very dilute topsoil slurries.
- 39 To what extent do molecular collisions arising from water vapour efflux impede stomatal O₃ influx?**
J. Uddling, R. Matyssek, J.B.C. Pettersson, G. Wieser
Negligence of molecular collisions arising from transpiration causes overestimation of stomatal ozone flux that increase with leaf-to-air mole fraction difference of water vapour.
- 43 Relevance of aerosol size spectrum analysis as support to qualitative source apportionment studies**
M. Manigrasso, A. Febo, F. Guglielmi, V. Ciambottini, P. Avino
Long range transport episodes are clearly demonstrated by time and size resolved aerosol and natural radioactivity measurements without any chemical analysis.
- 52 Use of geostatistics for remediation planning to transcend urban political boundaries**
T.M. Milillo, G. Sinha, J.A. Gardella Jr.
Using geostatistical interpolation and mapping results to involve the affected community can substantially improve remediation planning and promote its long-term effectiveness.
- 63 High levels of DDT in breast milk: Intake, risk, lactation duration, and involvement of gender**
H. Bouwman, H. Kylin, B. Sereda, R. Bornman
High levels of DDT in breast milk from a DDT-sprayed area exceeded known limits in milk, but did not affect duration lactation. Infant gender may affect levels of ΣDDT in breast milk.
- 71 Distribution and long-range transport of polyfluoroalkyl substances in the Arctic, Atlantic Ocean and Antarctic coast**
Z. Zhao, Z. Xie, A. Möller, R. Sturm, J. Tang, G. Zhang, R. Ebinghaus
Polyfluoroalkyl substances (PFASs) have been reported for the Arctic, Atlantic and the Southern Ocean, which improves understanding the fate of PFASs in the global oceans.

Continued on inside back cover

CONTENTS—Continued from outside back cover

- 78 Occupational exposure to airborne particles and other pollutants in an aviation base**
G. Buonanno, M. Bernabei, P. Avino, L. Stabile
A negligible impact of a high performance jet engine airport, in terms of airborne particles and other pollutants, was measured through an experimental campaign at three spatial scales.
- 88 Application of XAD-2 resin-based passive samplers and SPME–GC–MS/MS analysis for the monitoring of spatial and temporal variations of atmospheric pesticides in Luxembourg**
C. Schummer, L. Tuduri, O. Briand, B.M. Appenzeller, M. Millet
XAD-2 passive sampling of current-use pesticides in the atmosphere.
- 95 Mechanistic insights into the role of river sediment in the attenuation of the herbicide isoproturon**
S.B. Trinh, K.M. Hiscock, B.J. Reid
Catabolic activity with respect to IPU had a positive log-linear relationship with IPU concentration above a threshold concentration of between $0.1 \mu\text{g L}^{-1}$ and $1 \mu\text{g L}^{-1}$.
- 102 Proteomic analysis of Sydney Rock oysters (*Saccostrea glomerata*) exposed to metal contamination in the field**
E.L. Thompson, D.A. Taylor, S.V. Nair, G. Birch, G.C. Hose, D.A. Raftos
*This study used proteomics to analyse impacts of metal contamination on Sydney Rock oyster (*Saccostrea glomerata*) haemolymph in multiple field trials.*
- 113 Modeling and validation of on-road CO₂ emissions inventories at the urban regional scale**
M.N. Brondfield, L.R. Hutyrá, C.K. Gately, S.M. Raciti, S.A. Peterson
This work presents a new, reproducible methodology for downscaling and validating on-road CO₂ emissions estimates.
- 124 High nitrogen and elevated [CO₂] effects on the growth, defense and photosynthetic performance of two eucalypt species**
E. Novriyanti, M. Watanabe, M. Kitao, H. Utsugi, A. Uemura, T. Koike
The two eucalypts differently allocate assimilates in response to high N; while elevated [CO₂] tended to increase condensed tannin and to decrease total phenolics.
- 131 Using magnetic and chemical measurements to detect atmospherically-derived metal pollution in artificial soils and metal uptake in plants**
B. Sapkota, M.T. Cioppa
Variations in atmospheric particulate levels are measurable using magnetic and chemical techniques on soils and plant biomass, and suggest pollutant levels may be higher than previously recognized.
- 145 Persistent organic pollutants in the Tibetan surface soil: Spatial distribution, air–soil exchange and implications for global cycling**
X.-p. Wang, J.-j. Sheng, P. Gong, Y.-g. Xue, T.-d. Yao, K.C. Jones
Tibetan soils may be potential “secondary sources” of the HCB, low molecular weight PCBs and HCHs that are observed in air.
- 152 Natural and anthropogenically-produced brominated compounds in endemic dolphins from Western South Atlantic: Another risk to a vulnerable species**
M.B. Alonso, E. Eljarrat, M. Gorga, E.R. Secchi, M. Bassoi, L. Barbosa, C.P. Bertozzi, J. Marigo, M. Cremer, C. Domit, A.F. Azevedo, P.R. Dorneles, J.P.M. Torres, J. Lailson-Brito, O. Malm, D. Barceló
Concentrations and accumulation profiles of PBDEs, MeO-PBDEs and emerging brominated compounds in livers of dolphins from South Atlantic.
- 161 Diversities of phthalate esters in suburban agricultural soils and wasteland soil appeared with urbanization in China**
S. Kong, Y. Ji, L. Liu, L. Chen, X. Zhao, J. Wang, Z. Bai, Z. Sun
(1) Vegetable soil and wasteland soil exhibited higher PAEs concentrations; (2) PAEs in wasteland soils from suburban area of cities in China should be paid attention.
- 169 Ecological relationship analysis of the urban metabolic system of Beijing, China**
S. Li, Y. Zhang, Z. Yang, H. Liu, J. Zhang
Using Beijing as an example of an urban superorganism, we used ecological network analysis to describe the ecological relationships among the urban metabolic system's compartments.
- 177 Relationships between aging of PAHs and soil properties**
L. Luo, S. Lin, H. Huang, S. Zhang
Hard organic carbon and meso- and micropore distribution can act as better predictors of PAH sequestration in soils than total organic carbon and surface area, respectively.
- 183 The fractional patterns of polybrominated diphenyl ethers in the soil of the central Tibetan Plateau, China: The influence of soil components**
G.-L. Yuan, W. Xie, X.-C. Che, P. Han, C. Liu, G.-H. Wang
Certain minerals, such as clays, have a greater impact on the adsorption of POPs than SOC in certain areas.
- 190 Fate and occurrence of steroids in swine and dairy cattle farms with different farming scales and wastes disposal systems**
S. Liu, G.-G. Ying, R.-Q. Zhang, L.-J. Zhou, H.-J. Lai, Z.-F. Chen
The swine and cattle farms contribute higher steroids masses to the environment than the human sources.

CONTENTS—Continued from inside back cover

- 202 Physiological responses of lichens to factorial fumigations with nitric acid and ozone**
J. Riddell, P.E. Padgett, T.H. Nash III
Nitric acid can be highly toxic to lichens through several physiological mechanisms. Ozone is relatively non-toxic to fumigated lichens.
- 211 The effect of nitrogen addition on biomass production and competition in three expansive tall grasses**
P. Holub, I. Tůma, K. Fiala
Competition abilities of Calamagrostis and Bromus were greater than Brachypodium and the competitive superiority of Calamagrostis to Bromus was in N rich substrate.
- 217 The effect of airborne particles and weather conditions on pediatric respiratory infections in Cordoba, Argentine**
A.C. Amarillo, H.A. Carreras
The concentration of airborne particles as well as low socio-economic conditions and low education levels are significant risk factors for upper and lower respiratory infections in children from Cordoba, Argentine.
- 222 Sulfur-aided phytoextraction of Cd and Zn by *Salix smithiana* combined with in situ metal immobilization by gravel sludge and red mud**
M. Iqbal, M. Puschenreiter, E. Oburger, J. Santner, W.W. Wenzel
*Magnified Cd and Zn availability in S-amended rhizosphere of *Salix smithiana* enhances shoot Cd/Zn removal while metal leaching from the root zone can be reduced by immobilizing soil amendments.*
- 232 Bottom trawling resuspends sediment and releases bioavailable contaminants in a polluted fjord**
C. Bradshaw, I. Tjensvoll, M. Sköld, I.J. Allan, J. Molvaer, J. Magnusson, K. Naes, H.C. Nilsson
Bottom trawling in a Norwegian fjord resuspends tonnes of sediment per trawl pass and releases into the water contaminants (PCDD/Fs) that are taken up by blue mussels.
- 242 Exploratory study using proton induced X-ray emission analysis and histopathological techniques to determine the toxic burden of environmental pollutants**
S. Ejaz, M. Ashraf, L. Shakir, N. Ahmad
Toxic burden of environmental pollutants on wild animals
- 254 Bees, honey and pollen as sentinels for lead environmental contamination**
O. Lambert, M. Piroux, S. Puyo, C. Thorin, M. Larhantec, F. Delbac, H. Pouliquen
The aim of this study was to investigate the contamination of 18 apiaries by lead through analyses on 3 different matrices, foraging bees, honey and pollen over 2 beekeeping seasons and in 4 different landscape contexts.
- 260 Changing redox potential by controlling soil moisture and addition of inorganic oxidants to dissipate pentachlorophenol in different soils**
J. Lin, Y. He, J. Xu
The addition of inorganic oxidants limited the decrease in redox potential and inhibited the reductive dechlorination of pentachlorophenol.
- 268 Improved reproductive success in otters (*Lutra lutra*), grey seals (*Halichoerus grypus*) and sea eagles (*Haliaeetus albicilla*) from Sweden in relation to concentrations of organochlorine contaminants**
A.M. Roos, B.-M.V.M. Bäcklin, B.O. Helander, F.F. Rigét, U.C. Eriksson
The reproductive success in otters, grey seals and white-tailed sea eagles has increased as the concentrations of PCBs and Σ DDT have decreased supporting a causative relationship.
- 276 Spatio-temporal patterns of Cu contamination in mosses using geostatistical estimation**
A. Martins, R. Figueira, A.J. Sousa, C. Sérgio
The contamination of copper in mosses was estimated by spatio-temporal kriging, with determination of uncertainty classes in the temporal variation.