

Grass and Forage Science

Contents

- 305 J. Schellberg and L. da. S. Pontes
Plant functional traits and nutrient gradients on grassland
Review
- 320 L. Sulas, G. Piluzza, J. J. Rochon, J. P. Goby, J. M. Greef, U. Sölter, D. Headon and D. Scholefield
Assessing the potential for nutrient leaching from beneath grazed leguminous swards at four European sites
- 337 R. B. Valencia-Gica, R. S. Yost and G. Porter
Biomass production and nutrient removal by tropical grasses subsurface drip-irrigated with dairy effluent
- 350 D. R. LeCain, J. A. Morgan, G. L. Hutchinson, J. D. Reeder and F. A. Dijkstra
Interactions between elevated atmospheric CO₂ and defoliation on North American rangeland plant species at low and high N availability
- 361 S. J. Krizsan, Å. T. Randby and F. Westad
Effect of acetic acid, caproic acid and tryptamine on voluntary intake of grass silage by growing cattle
- 369 A. Jonker, M. Y. Gruber, Y. Wang, B. Coulman, J. J. McKinnon, D. A. Christensen and P. Yu
Foam stability of leaves from anthocyanidin- accumulating *Lc*-alfalfa and relation to molecular structures detected by fourier-transformed infrared-vibration spectroscopy
- 382 L. J. Iannone, A. D. Pinget, P. Nagabhyru, C. L. Schardl and J. P. De Battista
Beneficial effects of *Neotyphodium tembladera* and *Neotyphodium pampeanum* on a wild forage grass
- 391 J. L. Moir, B. J. Malcolm, K. C. Cameron and H. J. Di
The effect of dicyandiamide on pasture nitrate concentration, yield and N offtake under high N loading in winter and spring
- 403 K. Čámská and H. Skálová
Effect of low-dose N application and early mowing on plant species composition of mesophilous meadow grassland (*Arrhenatherion*) in Central Europe
- 411 M. C. Raeside, M. A. Friend, R. Behrendt, A. R. Lawson and S. G. Clark
Evaluation of tall fescue (*Festuca arundinacea*) as a forage for sheep in the temperate high-rainfall zone of south-eastern Australia
- 426 O. Scheneiter and C. Améndola
Tiller demography in tall fescue (*Festuca arundinacea*) swards as influenced by nitrogen fertilization, sowing method and grazing management
- 437 X. Z. Sun and G. C. Waghorn
Improving *in sacco* incubation technique to evaluate fresh forage for selecting fast-degrading perennial ryegrass (*Lolium perenne* L.)
- 446 T. Sasaki, T. Ohkuro, U. Jamsran and K. Takeuchi
Changes in the herbage nutritive value and yield associated with threshold responses of vegetation to grazing in Mongolian rangelands
- 456 E. Tommasino, S. Griffa, K. Grunberg, A. Ribotta, E. López Colomba, E. Carloni, M. Quiroga and C. M. Luna
Malondialdehyde content as a potential biochemical indicator of tolerant *Cenchrus ciliaris* L. genotypes under heat stress treatment
Research Note