

## MINI REVIEW

- A. M. ESTES, D. NESTEL, A. BELCARI, A. JESSUP, P. REMPOULAKIS, A. P. ECONOMOPOULOS  
A basis for the renewal of sterile insect technique for the olive fly, *Bactrocera oleae* (Rossi) 1-16

## ORIGINAL ARTICLES

- V. GIROLAMI, M. MARZARO, L. VIVAN, L. MAZZON, M. GREATTI, C. GIORIO, D. MARTON,  
A. TAPPARO  
Fatal powdering of bees in flight with particulates of neonicotinoids seed coating and  
humidity implication 17-26
- A. RAYBOULD, G. GRASER, K. HILL, K. WARD  
Ecological risk assessments for transgenic crops with combined insect-resistance traits: the  
example of Bt11 × MIR604 maize 27-37
- M. KRUGER, J. B. J. VAN RENSBURG, J. VAN DEN BERG  
Transgenic Bt maize: farmers' perceptions, refuge compliance and reports of stem borer  
resistance in South Africa 38-50
- I. BIGLER, W. NENTWIG, A. LINDFELD  
Food preference in the woodlouse *Porcellio scaber* (Isopoda) in a choice test with fungicidal  
GM wheat 51-59
- R. NDEMAH, F. SCHULTHESS, A. ABANG, R. T. GHOGOMU, N. NTONIFOR, S. DUPAS, B. LERÜ  
Suitability of cereal stemborers in Cameroon to Kenyan populations of the braconid larval  
parasitoid *Cotesia sesamiae* 60-69
- C. LÖFSTEDT, G. P. SVENSSON, E. V. JIRLE, O. ROSENBERG, A. ROQUES, J. G. MILLAR  
(3Z,6Z,9Z,12Z,15Z)-pentacosapentaene and (9Z,11E)-tetradecadienyl acetate: sex pheromone of  
the spruce coneworm *Dioryctria abietella* (Lepidoptera: Pyralidae) 70-78
- A. L. KNIGHT, L. L. STELINSKI, V. HEBERT, L. GUT, D. LIGHT, J. BRUNNER  
Evaluation of novel semiochemical dispensers simultaneously releasing pear ester and sex  
pheromone for mating disruption of codling moth (Lepidoptera: Tortricidae) 79-86
- R. S. MANN, S. TIWARI, J. M. SMOOT, R. L. ROUSEFF, L. L. STELINSKI  
Repellency and toxicity of plant-based essential oils and their constituents against *Diaphorina  
citri* Kuwayama (Hemiptera: Psyllidae) 87-96
- R. MUSUNDIRE, A. CHABI-OLAYE, K. KRÜGER  
Host plant effects on morphometric characteristics of *Liriomyza huidobrensis*, *L. sativae* and  
*L. trifolii* (Diptera: Agromyzidae) 97-108
- K. BIDINGER, S. LÖTTERS, D. RÖDDER, M. VEITH  
Species distribution models for the alien invasive Asian Harlequin ladybird (*Harmonia axyridis*) 109-123
- W. L. YEE  
Behavioural responses by *Rhagoletis indifferens* (Dipt., Tephritidae) to sweet cherry treated  
with kaolin- and limestone-based products 124-132
- M. UEFUNE, S. KUGIMIYA, K. SANO, J. TAKABAYASHI  
Herbivore-induced plant volatiles enhance the ability of parasitic wasps to find hosts on a  
plant 133-138
- G. CALABRIA, J. MÁCA, G. BÄCHLI, L. SERRA, M. PASCUAL  
First records of the potential pest species *Drosophila suzukii* (Diptera: Drosophilidae) in Europe 139-147
- P. J. LANDOLT, T. ADAMS, H. ROGG  
Trapping spotted wing drosophila, *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae),  
with combinations of vinegar and wine, and acetic acid and ethanol 148-154
- R. PENG, K. CHRISTIAN, K. GIBB  
The best time of day to monitor and manipulate weaver ant colonies in biological control 155-160