

Participatory generation of sustainability indicators in a natural protected area of Mexico N. Santana-Medina, S. Franco-Maass, E. Sánchez-Vera (Toluca, Mexico), J. Imbernon (Montpellier, France) and G. Nava-Bernal (Toluca, Mexico).....	1
Response of intertidal macrobenthic communities and primary producers to mitigation measures in a temperate estuary L. Marques, A. Carriço, F. Bessa, R. Gaspar, J.M. Neto and J. Patrício (Coimbra, Portugal).....	10
Do population indicators work? Investigating correlated responses of bird populations in relation to predator management J.M. Hoare (Christchurch, New Zealand), A. Monks (Dunedin, New Zealand) and C.F.J. O'Donnell (Christchurch, New Zealand).....	23
Soil and tree chemistry reflected the cumulative impact of acid deposition in <i>Pinus banksiana</i> and <i>Populus tremuloides</i> stands in the Athabasca oil sands region in western Canada K. Jung and S.X. Chang (Edmonton, Canada).....	35
Defining quantitative stream disturbance gradients and the additive role of habitat variation to explain macroinvertebrate taxa richness R. Ligeiro (Belo Horizonte, Brazil), R.M. Hughes, P.R. Kaufmann (Corvallis, OR, USA), D.R. Macedo, K.R. Firmiano, W.R. Ferreira, D. Oliveira (Belo Horizonte, Brazil), A.S. Melo (Goiânia, Brazil) and M. Callisto (Belo Horizonte, Brazil).....	45
Using Auchenorrhyncha (Insecta: Hemiptera), to develop a new insect index in measuring North American tallgrass prairie quality A.M. Wallner, B. Molano-Flores and C.H. Dietrich (Champaign, IL, USA).....	58
Response of fish-based metrics to anthropogenic pressures in temperate rocky reefs S. Henriques, M.P. Pais, M.I. Batista, M.J. Costa and H.N. Cabral (Lisboa, Portugal).....	65
Ranking municipal solid waste treatment alternatives based on ecological footprint and multi-criteria analysis M. Herva and E. Roca (Santiago de Compostela, Spain).....	77
Ecological manifestation of slope instability, its application in identification of areas of potential hill slope movement in Indian Himalaya P.S. Negi (Dehra Dun, India).....	85
Diversity, occurrence and feeding traits of caddisfly larvae as indicators for ecological integrity of river- floodplain habitats along a connectivity gradient F.W.B. Van den Brink (Maastricht, The Netherlands), G. Van der Velde (Nijmegen, The Netherlands and Leiden, The Netherlands) and S. Wijnhoven (Yerseke, The Netherlands).....	92
Ecological indicators for protection impact assessment at two scales in the Bozin and Marakhil protected area, Iran A. Henareh Khalyani, A.L. Mayer, C.R. Webster and M.J. Falkowski (Houghton, MI, USA).....	99
Characterizing and evaluating successional pathways of fen degradation and restoration J. Schrautzer (Kiel, Germany), F. Sival (Wageningen, The Netherlands), M. Breuer (Kiel, Germany), H. Runhaar (Nieuwegein, The Netherlands) and A. Fichtner (Kiel, Germany).....	108
Development of a framework for landscape assessment of Taiwanese wetlands T.-S. Chen (Nantou, Taiwan and Taichung, Taiwan) and H.-J. Lin (Taichung, Taiwan and Taipei, Taiwan).....	121

Identifying keystone trophic groups in benthic ecosystems: Implications for fisheries management M. Ortiz (Antofagasta, Chile), R. Levins (Boston, MA, USA), L. Campos, F. Berrios, F. Campos (Antofagasta, Chile), F. Jordán (Rovereto, Italy), B. Hermosillo, J. Gonzalez (Antofagasta, Chile) and F. Rodriguez (Zapopan, Mexico)	133
Development of an Angiosperm Quality Assessment Index (AQuA-Index), for ecological quality evaluation of Portuguese water bodies-A multi-metric approach I. Caçador (Lisbon, Portugal), J.M. Neto (Coimbra, Portugal), B. Duarte (Lisbon, Portugal), D.V. Barroso (Coimbra, Portugal), M. Pinto (Lisbon, Portugal) and J.C. Marques (Coimbra, Portugal)	141
Importance of scalar and riparian habitat effects for assessment of ecological status using littoral diatoms M.A. Snell (Lancaster, UK) and K.A. Irvine (Delft, The Netherlands)	149
Are aquatic insect species sensitive to banana plant cultivation? J.J. Corbi, P. Kleine and S. Trivinho-Strixino (São Carlos, Brazil)	156
Derivation of dynamic reference values for the classification of contaminant concentrations in bream (<i>Abramis brama</i>), of German rivers and lakes using data of the Environmental Specimen Bank D. Teubner, P. Müller and M. Paulus (Trier, Germany)	162
An indicator-based method for quantifying farm multifunctionality P.S. Andersen, H. Vejre (Frederiksberg C, Denmark), T. Dalgaard (Tjele, Denmark) and J. Brandt (Roskilde, Denmark)	166
Application of analytic hierarchy process in selecting a biological indicator for a river flow restoration W. Huang (Beijing, China and Fuku oka, Japan), S. Yano (Fukuoka, Japan), J. Zhang and Y. Wang (Chengdu, China)	180
Detecting the structural and functional impacts of fine sediment on stream invertebrates C. Buendia (Aberdeen, UK and Lleida, Spain), C.N. Gibbins (Aberdeen, UK), D. Vericat (Lleida, Spain and UK), R.J. Batalla (Catalonia, Spain and Wales, UK) and A. Douglas (Aberdeen, UK)	184
Radioecological indexes of fallout measurements from the Fukushima nuclear accident M. Manolopoulou, S. Stoulos, A. Ioannidou, E. Vagena and C. Papastefanou (Thessaloniki, Greece)	197
Lake Macrophyte Nutrient Index of standing waters in Serbia (LIMNIS), D. Laketić, S. Radulović, M. Živković, T. Jurca and M.H. Alford	200
Assessing the relationship between the Lake Habitat Survey and littoral macroinvertebrate communities in European lakes E. McGoff (Uppsala, Sweden), J. Aroviita (Finland), F. Pilotto (Berlin, Germany and Rome, Italy), O. Miler (Berlin, Germany), A.G. Solimini (Rome, Italy), G. Porst (Berlin, Germany), T. Jurca, L. Donohue (Dublin, Ireland) and L. Sandin (Silkeborg, Denmark)	205
Bivalve condition index as an indicator of aquaculture intensity: A meta-analysis R. Filgueira (Halifax, Canada and Moncton, Canada), L.A. Comeau, T. Landry (Moncton, Canada), J. Grant (Halifax, Canada), T. Guyondet (Moncton, Canada) and A. Mallet (Dartmouth, Canada)	215
Ecological assessment of groundwater trophic status by using artificial substrates to monitor biofilm growth and activity F. Mermillod-Blondin (Villeurbanne, France), A. Foulquier (Villeurbanne, France and Lyon Cedex, France), C. Maazouzi, S. Navel, Y. Negrutiu, A. Vienney, L. Simon and P. Marmonier (Villeurbanne, France)	230
Methodology for determining the ecological footprint of the construction of residential buildings in Andalusia (Spain), J. Solís-Guzmán, M. Marrero and A. Ramírez-de-Arellano (Seville, Spain)	239
Improving the "chain and tape" method: A combined topography index for marine fish ecology studies M.P. Pais, S. Henriques, M.J. Costa and H.N. Cabral (Lisboa, Portugal)	250
Soil and tree ring chemistry of <i>Pinus banksiana</i> and <i>Populus tremuloides</i> stands as indicators of changes in atmospheric environments in the oil sands region of Alberta, Canada K. Jung (Alberta, Canada and Suwon, Republic of Korea), W.-J. Choi (Gwangju, Republic of Korea), S.X. Chang and M.A. Arshad (Alberta, Canada)	256
An ecological quality status assessment procedure for soft-sediment benthic habitats: Weighing alternative approaches G. Van Hoey (Oostende, Belgium), D.C. Permuy (Oostende, Belgium and Ghent, Belgium), S. Vandendriessche (Oostende, Belgium), M. Vincx (Ghent, Belgium) and K. Hostens (Oostende, Belgium)	266
Assessing biodiversity with sound: Do acoustic diversity indices reflect phylogenetic and functional diversities of bird communities? A. Gasc, J. Sueur, F. Jiguet (Paris, France), V. Devictor (Montpellier, France), P. Grandcolas, C. Burrow, M. Depraetere (Paris, France) and S. Pavoine (Paris, France and Oxford, UK)	279