

CONTENTS

| | |
|--|---------|
| C. BUSSI and D. PLENET: Effect of three different irrigation regimes on young apricot (<i>Prunus armeniaca</i> L. Batsch) trees | 519–524 |
| FRANCESCO PANARA, DESPOINA PETOUMENOU, ORNELLA CALDERINI, FRANCESCA DINI, CLAUDIO D'ONOFRIO, LAURA BEDINI and ALBERTO PALLIOTTI: Ampelographic and genetic characterisation of ancestral grapevine (<i>Vitis vinifera</i> L.) accessions present in the Umbria Region (Central Italy) | 525–530 |
| R. L. BHARDWAJ: Effects of nine different propagation media on seed germination and the initial performance of papaya (<i>Carica papaya</i> L.) seedlings | 531–536 |
| BARBARA MENIN, ANDREA MOGLIA, CINZIA COMINO, JOHANNA CHRISTINA HAKKERT, SERGIO LANTERI and JULES BEEKWILDER: <i>In vitro</i> callus-induction in globe artichoke (<i>Cynara cardunculus</i> L. var. <i>scolymus</i>) as a system for the production of caffeoylquinic acids | 537–542 |
| O. A. ODUWAYE, D. K. OJO and C. O. ALAKE: Using genotype plus genotype × environment (GGE) bi-plot analysis to identify promising amaranth accessions in Western Nigeria | 543–550 |
| P. D. GAMES, C. R. KOSCKY-PAIER, H. O. ALMEIDA-SOUZA, M. O. BARBOSA, P. W. P. ANTUNES, L. C. CARRIJO, P. R. G. PEREIRA and M. C. BARACAT-PEREIRA: <i>In vitro</i> anti-bacterial and anti-fungal activities of hydrophilic plant defence compounds obtained from the leaves of bell pepper (<i>Capsicum annuum</i> L.). | 551–558 |
| A. A. WAMAN, P. BOHRA, B. N. SATHYANARAYANA, S. C. CHANDRASHEKAR and R. THILAKA RANI: Are bananas (<i>Musa</i> spp.) really safe from their aesthetic relatives? Screening potential alternative hosts of <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> | 559–562 |
| GUOPENG LI, HUIJUAN JIA, RUIYUAN WU and YUANWEN TENG: Changes in volatile organic compound composition during the ripening of 'Nanguoli' pears (<i>Pyrus ussuriensis</i> Maxim) harvested at different growing locations | 563–570 |
| M. PAGTER, K. K. PETERSEN and K. H. KJAER: Direct and indirect effects of shoot-and/or root-chilling stress on growth, photosynthesis, and osmotic root water uptake in <i>Kalanchoe blossfeldiana</i> Poelln. 'Molly' | 571–579 |
| S. K. KAUSHIK, R. SHARMA, I. D. GARG, B. P. SINGH, S. K. CHAKRABARTI, V. BHARDWAJ and S. K. PANDEY: Development of a triplex (YYYy) parental potato line with extreme resistance to potato virus Y using marker assisted selection | 580–584 |
| MINERVA GHANI, SURINDER KUMAR and MANISHA THAKUR: Induction of novel variants through physical and chemical mutagenesis in Barbeton daisy (<i>Gerbera jamesonii</i> Hook.) | 585–590 |
| A. SØNSTEBY, J. A. STAVANG and O. M. HEIDE: Production of high-yielding raspberry long canes: The way to 3 kg of fruit per cane | 591–599 |
| J. HAJILOU and S. FAKHIMREZAEI: Effects of post-harvest calcium chloride or salicylic acid treatments on the shelf-life and quality of apricot fruit | 600–604 |
| K. V. RAVISHANKAR, K. P. RAGHAVENDRA, V. ATHANI, A. REKHA, K. SUDEEPA, D. BHAVYA, V. SRINIVAS and L. ANANAD: Development and characterisation of microsatellite markers for wild banana (<i>Musa balbisiana</i>) | 605–609 |
| NASIM BASTAM, BAHRAM BANINASAB and CYRUS GHOBADI: Interactive effects of ascorbic acid and salinity stress on the growth and photosynthetic capacity of pistachio (<i>Pistacia vera</i> L.) seedlings | 610–616 |
| R. L. SU, H. T. HU, M. WU, W. D. GUO, J. H. CHEN, C. C. WANG and L. YANG: Isolation and characterisation of a UDP-glucose pyrophosphorylase gene associated with fruit ripening in autumn olive (<i>Elaeagnus umbellata</i> Thunb.) | 617–623 |
| C. BARCELÓ-VIDAL, J. BONANY, J. A. MARTÍN-FERNÁNDEZ and J. CARBÓ: Modelling of weather parameters to predict russet on 'Golden Delicious' apple | 624–630 |
| QING-JIE FAN, SI-CHENG ZHENG, FENG-XIA YAN, BING-XEE ZHANG, GUANG QIAO and XIAO-PENG WEN: Efficient regeneration of dragon fruit (<i>Hylocereus undatus</i>) and an assessment of the genetic fidelity of <i>in vitro</i> -derived plants using ISSR markers | 631–637 |
| H. M. I. HERATH: <i>In vitro</i> propagation of <i>Chirita moonii</i> Gardn. (Gesneriaceae), a potential ornamental plant endemic to Sri Lanka | 638–642 |
| G. MADAAN, S. K. GOSAL, S. S. GOSAL, G. S. SAROA and M. I. S. GILL: Effect of microbial inoculants on the growth and yield of micropropagated banana (<i>Musa indica</i>) cv. Grand Naine | 643–649 |
| C. MA, W. N. ZHANG, X. W. DUAN, Y. LU and T. Z. LI: Comparative analysis of the efficiencies of siRNA and amiRNA gene silencing of the tobacco lycopene cyclase gene (<i>NsCrtL-1</i>) | 650–656 |
| T. MORI, T. UMEDA, T. HONDA, K. ZUSHI, T. WAJIMA and N. MATSUZOE: Varietal differences in the chlorogenic acid, anthocyanin, soluble sugar, organic acid, and amino acid concentrations of eggplant fruit | 657–663 |
| V. RADHIKA, D. C. L. REDDY, R. VEEREGOWDA and C. ASWATH: <i>In silico</i> identification and validation of microsatellite markers from onion EST sequences | 664–670 |
| SESHADRI SHIVASHANKAR, JAYA JOSHI and MANOHARAN SUMATHI: The role of seed viability in the development of corky tissue in sapota (<i>Manilkara achras</i>) fruit in India | 671–677 |