

# Contents

The white collar complex is essential for sexual reproduction but dispensable for conidiation and invasive growth in <i>Fusarium verticillioides</i> Á. M. Bodor, V. Stubnya, A. L. Ádám, M. Ládai and L. Hornok .....	1
<i>Poaceae</i> : A rich source of endophytic fusaria Á. Szécsi, D. Magyar, S. Tóth and C. Szőke .....	19
Tomato spotted wilt virus (TSWV) in birthwort ( <i>Aristolochia clematitis</i> L.) in Hungary E. Cseh, M. Apró, G. Bese, L. Krizbai, K. Bóka, R. Gáborjányi and A. P. Takács .....	33
Resistance gene analogues in mango against mango malformation Saboki Ebrahim, T. R. Sharma, K. Usha and Bhupinder Singh .....	39
Evaluation of black gram [ <i>Vigna mungo</i> (L.) Hepper] genotypes against whitefly, <i>Bemisia tabaci</i> (Gennadius) under screen-house conditions G. K. Taggar, R. S. Gill and J. S. Sandhu.....	53
Seasonal abundance of sucking insect pests on transgenic <i>Bt</i> cotton vis-à-vis weather parameters in Punjab, India P. S. Shera, Vijay Kumar and Anand Aneja .....	63
Seasonal flight of the potato tuber moth, <i>Phthorimaea operculella</i> (Zeller) (Lepidoptera: Gelechiidae) in three regions in Bulgaria established by pheromone traps M. Subchev, T. B. Toshova, D. I. Atanasova, V. D. Petrova and M. Tóth.....	75
Catches of target and non-target scarab pests in floral-chemical baited color traps in Bulgaria M. A. Subchev, T. B. Toshova, R. A. Andreev, V. D. Petrova, V. D. Maneva, T. S. Spasova, N. T. Marinova, P. M. Minkov and D. I. Velchev .....	87
Intra-guild vs extra-guild prey: effect on development, predation and preference of <i>Typhlodromus negevi</i> Swirski and Amitai and <i>Typhlodromips swirskii</i> (Athias-Henriot) (Acari: Phytoseiidae) F. M. Momen, H. Hussein and A. S. Reda .....	95
An annotated list of the encyrtids of Egypt (Hymenoptera: Chalcidoidea: Encyrtidae) G. A. Evans and S. Abd-Rabou .....	107
A new species of cecidomyiid (Diptera: Cecidomyiidae) predator of scale insect (Hemiptera: Coccoidea) pests of pineapple M. P. Culik and J. A. Ventura .....	129
Morphological characteristics of pupal stages of Thripidae collected in Poland (Thysanoptera, Terebrantia) H. Kucharczyk and M. Kucharczyk .....	135

Redescription of <i>Leipothrix polygalae</i> (Farkas, 1968) n. comb. from Hungary (Acari: Prostigmata: Eriophyoidea)	
<i>G. Ripka</i> .....	143
New data on the mesostigmatid mite fauna of Hungary (Acari: Mesostigmata)	
<i>Á. Szabó, G. Ripka, Zs. Hajdu, B. Tempfli, M. Varga, I. Mészáros, Cs. Kutasi, T. Németh and B. Péntzes</i> .....	149
A study on the ants (Hymenoptera: Vespoidea: Formicidae) from Western Iran	
<i>H. Ghahari and C. A. Collingwood</i> .....	155
Insecticidal effectiveness of certain bio-insecticides, inert dusts and modified atmospheres against <i>Sitophilus oryzae</i> (L.) (Coleoptera: Curculionidae) on stored wheat	
<i>Nour El-Hoda A. Zidan</i> .....	165
Repellent, antifeedent and toxic effects of three essential oils on the two spotted spider mite, <i>Tetranychus urticae</i> Koch (Acari: Tetranychidae)	
<i>H. Hussein, A. S. Reda and F. M. Momen</i> .....	177