

CONTENTS

Regular Paper

- MIYAUCHI M, HIRAI C and NAKAJIMA H: The Solar Exposure Time Required for Vitamin D₃ Synthesis in the Human Body Estimated by Numerical Simulation and Observation in Japan. 257–263
- MARUYAMA C, KIKUCHI N, MASUYA Y, HIROTA S, ARAKI R and MARUYAMA T: Effects of Green-Leafy Vegetable Intake on Postprandial Glycemic and Lipidemic Responses and α -Tocopherol Concentration in Normal Weight and Obese Men. 264–271
- CHEN Y-Y, LAI M-H, HUNG H-Y and LIU J-F: Sweet Potato [*Ipomoea batatas* (L.) Lam. “Tainong 57”] Starch Improves Insulin Sensitivity in High-Fructose Diet-Fed Rats by Ameliorating Adipocytokine Levels, Pro-Inflammatory Status, and Insulin Signaling. 272–280
- IMAEDA N, KURIKI K, FUJIWARA N, GOTO C, TOKUDOME Y and TOKUDOME S: Usual Dietary Intakes of Selected Trace Elements (Zn, Cu, Mn, I, Se, Cr, and Mo) and Biotin Revealed by a Survey of Four-Season 7-Consecutive Day Weighed Dietary Records in Middle-Aged Japanese Dietitians. 281–288
- MIZUNOYA W, SAWANO S, IWAMOTO Y, SATO Y, TATSUMI R and IKEUCHI Y: Effect of 48-h Food Deprivation on the Expressions of Myosin Heavy-Chain Isoforms and Fiber Type-Related Factors in Rats 289–298
- KANEKO K, ITO C, KOIZUMI K, WATANABE S, UMEDA Y and ISHIKAWA-TAKATA K: Resting Energy Expenditure (REE) in Six- to Seventeen-Year-Old Japanese Children and Adolescents 299–309
- TSUBOTA-UTSUGI M, IMAI E, NAKADE M, MATSUMOTO T, TSUBOYAMA-KASAOKA N, NISHI N and TSUBONO Y: Evaluation of the Prevalence of Iodine Intakes above the Tolerable Upper Intake Level from Four 3-Day Dietary Records in a Japanese Population 310–316
- HIRASAKA K, MAEDA T, IKEDA C, HARUNA M, KOHNO S, ABE T, OCHI A, MUKAI R, OARADA M, TESHIMA-KONDO S, OHNO A, OKUMURA Y, TERAJIMA J and NIKAWA T: Isoflavones Derived from Soy Beans Prevent MuRF1-Mediated Muscle Atrophy in C2C12 Myotubes through SIRT1 Activation 317–324
- SONE Y, KIDO T, AINUKI T, SONODA M, ICHI I, KODAMA S, SONE H, KONDO K, MORITA Y, EGAWA S, KAWAHARA K, OTSUKA Y and FUJIWARA Y: Genetic Variants of the Fatty Acid Desaturase Gene Cluster Are Associated with Plasma LDL Cholesterol Levels in Japanese Males 325–335
- NI Q, WANG Z, XU G, GAO Q, YANG D, MORIMATSU F and ZHANG Y: Altitudinal Variation of Antioxidant Components and Capability in *Indocalamus latifolius* (Keng) McClure Leaf 336–342

Note

- SHIBATA K, TACHIKI A, MUKAEDA K, FUKUWATARI T, SASAKI S and JINNO Y: Changes in Plasma Pyridoxal 5'-Phosphate Concentration during Pregnancy Stages in Japanese Women. 343–346
- QIAO Y, FENG J, YANG J and GU G: The Relationship between Dietary Vitamin A Intake and the Levels of Sialic Acid in the Breast Milk of Lactating Women 347–351
- KISHIMOTO Y, KANAHORI S, SAKANO K and EBIHARA S: The Maximum Single Dose of Resistant Maltodextrin That Does Not Cause Diarrhea in Humans. 352–357
- ITO Y, MITANI T, HARADA N, ISAYAMA A, TANIMORI S, TAKENAKA S, NAKANO Y, INUI H and YAMAJI R: Identification of Carbonyl Reductase 1 as a Resveratrol-Binding Protein by Affinity Chromatography Using 4'-Amino-3,5-dihydroxy-*trans*-stilbene 358–364

Letter to the Editor

- KAPOOR S: Vitamin B₆ and Its Effect on Systemic Carcinogenesis 365