

**Fernandez-Gonzalez V, Aguado-Giménez F,  
Gairin JI, Sanchez-Jerez P**

Exploring patterns of variation in amphipod assemblages at multiple spatial scales: natural variability versus coastal aquaculture effect 93–105

**Serra-Llinares RM, Nilsen R, Uglem I,  
Arechavala-Lopez P, Bjørn PA, Noble C**

Post-escape dispersal of juvenile Atlantic cod *Gadus morhua* from Norwegian fish farms and their potential for recapture 107–116

**Jones B, Fuller S, Carton AG**

Earthy-muddy tainting of cultured barramundi linked to geosmin in tropical northern Australia 117–124

**Bui S, Oppedal F, Korsøen ØJ,  
Dempster T**

Modifying Atlantic salmon behaviour with light or feed stimuli may improve parasite control techniques 125–133

**Somarakis S, Pavlidis M, Saapoglou C,  
Tsigenopoulos CS, Dempster T**

Evidence for 'escape through spawning' in large gilt-head sea bream *Sparus aurata* reared in commercial sea-cages 135–152

**Arechavala-Lopez P, Sanchez-Jerez P,  
Bayle-Sempere JT, Uglem I, Mladineo I**

*AS WE SEE IT*: Reared fish, farmed escapees and wild fish stocks—a triangle of pathogen transmission of concern to Mediterranean aquaculture management 153–161

**Doubleday ZA, Clarke SM, Li X, Pecl GT,  
Ward TM, Battaglione S, Frusher S, Gibbs PJ,  
Hobday AJ, Hutchinson N, Jennings SM,  
Stoklosa R**

Assessing the risk of climate change to aquaculture: a case study from south-east Australia 163–175

**Xia B, Gao QF, Li H, Dong SL, Wang F**

Turnover and fractionation of nitrogen stable isotope in tissues of grass carp *Ctenopharyngodon idellus* 177–186