

CONTENTS

Articles

- KANZAKI, N., MASUYA, H., TAKI, H., OKABE, K. & CHEN, C.-Y.— Description of *Ruehmaphelenchus formosanus* n. sp. (Tylenchina: Aphelenchoididae) isolated from *Euwallacea fornicates* from Taiwan 895-906
- TOMALAK, M., WORRALL, J. & FILIPIAK, A.— *Bursaphelenchus masseyi* sp. n. (Nematoda: Parasitaphelenchinae) – a nematode associate of the bark beetle, *Trypophloeus populi* Hopkins (Coleoptera: Curculionidae: Scolytinae), in aspen, *Populus tremuloides* Michx. affected by sudden aspen decline in Colorado 907-921
- TOYOTA, K., GOTO, K. & ITO, D.— Effect of bean sprout residue on the hatching and density of the soybean cyst nematode *Heterodera glycines* in soil 923-929
- SATO, E., OKUMURA, I., KOKATSU, Y. & TOYOTA, K.— Effects of the density of root-lesion nematode (*Pratylenchus penetrans*), soil chemical and microbial properties on the damage to Japanese radish 931-938
- AYUB, F., STRAUCH, O., SEYCHELLES, L. & EHLERS, R.-U.— Influence of temperature on life history traits of the free-living, bacterial-feeding nematode *Panagrolaimus* sp. strain NFS-24 939-946
- MCSORLEY, R.— Effect of forest and field habitat features on abundance of genera in nematode assemblages 947-956
- KANZAKI, N., GIBLIN-DAVIS, R.M., YE, W., HERRE, E.A. & CENTER, B.J.— Description of *Parasitodiplogaster pharmaconema* n. sp. and redescription of *P. maxinema* from *Ficus maxima* Mill. (Moraceae) 957-974
- KANZAKI, N. & TANAKA, R.— *Sheraphelenchus sucus* n. sp. (Tylenchina: Aphelenchoididae) isolated from sap flow of *Quercus serrata* in Japan 975-990
- NGUYEN, X.H., NAING, K.W., LEE, Y.S., JUNG, W.J., ANEES, M. & KIM, K.Y.— Antagonistic potential of *Paenibacillus elgii* HOA73 against the root-knot nematode, *Meloidogyne incognita* 991-1000
- HERRMANN, M., RAGSDALE, E.J., KANZAKI, N. & SOMMER, R.J.— *Sudhausia aristotokia* n. gen., n. sp. and *S. crassa* n. gen., n. sp. (Nematoda: Diplogastridae): viviparous new species with precocious gonad development 1001-1020