

MINI-REVIEW

- C. M. OLIVEIRA, A. M. AUAD, S. M. MENDES, M. R. FRIZZAS
Economic impact of exotic insect pests in Brazilian agriculture 1–15

ORIGINAL ARTICLE

- A. COCCO, S. DELIPERI, G. DELRIO
Control of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in greenhouse tomato crops using the mating disruption technique 16–28

SHORT COMMUNICATION

- V. H. P. BUENO, J. C. VAN LENTEREN, J. C. LINS JR, A. M. CALIXTO, F. C. MONTES, D. B. SILVA, L. D. SANTIAGO, L. M. PÉREZ
New records of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) predation by Brazilian Hemipteran predatory bugs 29–34

ORIGINAL ARTICLES

- V. GIROLAMI, M. MARZARO, L. VIVAN, L. MAZZON, C. GIORIO, D. MARTON, A. TAPPARO
Aerial powdering of bees inside mobile cages and the extent of neonicotinoid cloud surrounding corn drillers 35–44

- M. OBOPILE, R. B. HAMMOND, P. R. THOMISON
Interaction among planting dates, transgenic maize, seed treatment, corn rootworm damage and grain yield 45–55

- Z. SVOBODOVÁ, O. HABUŠTOVÁ, F. SEHNAL, M. HOLEC, H. M. HUSSEIN
Epigeic spiders are not affected by the genetically modified maize MON 88017 56–67

- C. HANNA, C. HANNA
The lethal and sublethal effects of three pesticides on the striped lynx spider (*Oxyopes salticus* Hentz) 68–76

- M. J. DOMINGUE, J. P. LELITO, I. FRASER, V. C. MASTRO, J. H. TUMLINSON, T. C. BAKER
Visual and chemical cues affecting the detection rate of the emerald ash borer in sticky traps 77–87

- S. ANBESSE, R.-U. EHLERS
Heterorhabditis sp. not attracted to synthetic (*E*)- β -caryophyllene, a volatile emitted by roots upon feeding by corn rootworm 88–96

- M. HORI, S. ENYA
Attractiveness of synthetic volatile blends of flowering rice panicles to *Trigonotylus caelestialium* (Kirkaldy) (Heteroptera: Miridae) 97–103

- N. S. MANDOUR, Y. KAINOH, R. OZAWA, M. UEFUNE, J. TAKABAYASHI
Effects of prohydrojasmon-treated corn plants on attractiveness to parasitoids and the performance of their hosts 104–112

- J. M. MARUBAYASHI, V. A. YUKI, K. C. G. ROCHA, T. MITUTI, F. M. PELEGRINOTTI, F. Z. FERREIRA, M. F. MOURA, J. NAVAS-CASTILLO, E. MORIONES, M. A. PAVAN, R. KRAUSE-SAKATE
At least two indigenous species of the *Bemisia tabaci* complex are present in Brazil 113–121

- L. L. ESTERHUIZEN, K. G. MABASA, S. W. VAN HEERDEN, H. CZOSNEK, J. K. BROWN, H. VAN HEERDEN, M. E. C. REY
Genetic identification of members of the *Bemisia tabaci* cryptic species complex from South Africa reveals native and introduced haplotypes 122–135

- T. J. ASLAM, S. N. JOHNSON, A. J. KARLEY
Plant-mediated effects of drought on aphid population structure and parasitoid attack 136–145

- S. NIEDERMAYER, E. OBERMAIER, J. L. M. STEIDLE
Some like it hot, some not: influence of extreme temperatures on *Lariophagus distinguendus* and *Anisopteromalus calandrae* 146–152

- S. SCHULZE-BOPP, J. A. JEHL
Development of a direct test of baculovirus resistance in wild codling moth populations 153–160