

CONTENTS

Variation of energy flow and greenhouse gas emissions in vineyards located in Natura 2000 sites V.D. Litskas, C.S. Karaolis, G.C. Menexes, A.P. Mamolos, T.M. Koutsos and K.L. Kalburtji (Thessaloniki, Greece)	1
<i>Myriophyllum aquaticum</i> as a biomonitor of water heavy metal input related to agricultural activities in the Xanaes River (Córdoba, Argentina) C.A. Harguinteguy (Córdoba, Argentina), R. Schreiber (Leipzig, Germany) and M.L. Pignata (Córdoba, Argentina).....	8
Ecosystem functional units characterized by satellite observed phenology and productivity gradients: A case study for Europe E. Ivits, M. Cherlet, W. Mehl and S. Sommer (Ispra, Italy).....	17
Comparing stressor-specific indices and general measures of taxonomic composition for assessing the status of boreal lacustrine macrophyte communities A. Kanninen (Kuopio, Finland), S. Hellsten (Finland) and H. Hämäläinen (Finland)	29
Uncertainties in biotic indicators and a corrigendum to Ponader et al. (2007): Implications for biomonitoring B.O.L. Demars (Aberdeen, UK).....	44
Is coarse taxonomy sufficient for detecting macroinvertebrate patterns in floodplain lakes? X. Jiang (Wuhan, China and Beijing, China), J. Xiong (Wuhan, China), Z. Song (Wuhan, China and Beijing, China), J.C. Morse (Clemson, SC, USA), F.C. Jones (Dorset, Canada) and Z. Xie (Wuhan, China).....	48
The collembola <i>Lobella sokamensis</i> juvenile as a new soil quality indicator of heavy metal pollution Y.-J. An, S.W. Kim and W.-M. Lee (Seoul, Republic of Korea)	56
The grazing fingerprint: Modelling species responses and trait patterns along grazing gradients in semi-arid Namibian rangelands D. Wesuls, M. Pellowski, S. Suchrow, J. Oldeland (Hamburg, Germany), F. Jansen (Greifswald, Germany) and J. Dengler (Hamburg, Germany).....	61
Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua L. Rousseau (Bondy Cedex, France), S.J. Fonte (Cali, Colombia), O. Téllez, R. van der Hoek (Managua, Nicaragua) and P. Lavelle (Bondy Cedex, France and Cali, Colombia)	71
Stressor prioritisation in riverine ecosystems: Which environmental factors shape benthic invertebrate assemblage metrics? A. Sundermann, M. Gerhardt (Gelnhausen, Germany and Frankfurt am Main, Germany), H. Kappes (RA Leiden, The Netherlands) and P. Haase (Gelnhausen, Germany and Frankfurt am Main, Germany).....	83
Quantifying associations of large vertebrates with salmon in riparian areas of British Columbia streams by means of camera-traps, bait stations, and hair samples T.F. Shardlow and K.D. Hyatt (Nanaimo, Canada).....	97
Assessing air quality with regards to its effect on human health in the European Union through air quality indices K. Dimitriou, A.K. Paschalidou and P.A. Kassomenos (Ioannina, Greece)	108

Contents continued from outside back cover

Strong isolation-by-distance in the absence of genetic population structure in the eelpout (<i>Zoarces viviparus</i> , Linnaeus 1758) T. Kinitz, M. Quack, M. Paulus, M. Veith (Trier, Germany), S. Bergek (Drottningholm, Sweden), J. Strand (Roskilde, Denmark), A. Tuvikene (Rannu, Estonia), A. Soirinsuo (Helsinki, Finland) and A. Hochkirch (Trier, Germany)	116
Replying to Domingues et al., <i>Ecological Indicators</i> , 24, 245–255, http://dx.doi.org/10.1016/j.ecolind.2012.06.020 A.C. Brito, V. Brotas, M. Caetano, T.P. Coutinho, T. Moita (Lisboa, Portugal), A.A. Bordalo (Portugal), J. Icely (Bispo, Portugal and Coimbra, Portugal), J.M. Neto (Coimbra, Portugal) and J. Serôdio (Aveiro, Portugal)	123
Modelling the spatial distribution of linear landscape elements in Europe E.H. van der Zanden, P.H. Verburg (Amsterdam, The Netherlands) and C.A. Mûcher (Wageningen, The Netherlands).	125