

CONTENTS

Volume 174 March 2013

- 1 Life stage-specific effects of the fungicide pyrimethanil and temperature on the snail *Physella acuta* (Draparnaud, 1805) disclose the pitfalls for the aquatic risk assessment under global climate change**
A. Seeland, J. Albrand, J. Oehlmann, R. Müller
The aquatic risk of pesticides under climate change cannot be adequately assessed by recent strategies for the regular risk assessment of agrochemicals.
- 10 Global assessment of the effects of terrestrial acidification on plant species richness**
L.B. Azevedo, R. van Zelm, A.J. Hendriks, R. Bobbink, M.A.J. Huijbregts
Relationships of potential species richness loss along a soil pH gradient are proposed.
- 16 Unravelling remote sensing signatures of plants contaminated with gasoline and diesel: An approach using the red edge spectral feature**
I.D. Sanches, C.R. Souza Filho, L.A. Magalhães, G.C.M. Quitério, M.N. Alves, W.J. Oliveira
Leaf and canopy spectra of healthy plants were significantly different from spectra of plants contaminated with liquid hydrocarbons allowing the contaminations to be detected remotely.
- 28 Species-specific differences in the accumulation features of organohalogen contaminants and their metabolites in the blood of Japanese terrestrial mammals**
H. Mizukawa, K. Nomiyama, S. Nakatsu, S. Yachimori, T. Hayashi, Y. Tashiro, Y. Nagano, S. Tanabe
Cats showed specific residue pattern of OH-PCBs compare to other species and marine natural occurrence OH-/MeO-PBDEs were detected in terrestrial mammals.
- 38 Initial transport and retention behaviors of ZnO nanoparticles in quartz sand porous media coated with *Escherichia coli* biofilm**
X. Jiang, X. Wang, M. Tong, H. Kim
Biofilm enhanced ZnO nanoparticle deposition and altered spacial distribution in porous media.
- 50 Photosynthetic traits of Siebold's beech and oak saplings grown under free air ozone exposure in northern Japan**
M. Watanabe, Y. Hoshika, N. Inada, X. Wang, Q. Mao, T. Koike
Photosynthesis of beech is more sensitive to free air ozone exposure than that of oak.
- 57 Dechlorane Plus flame retardant in kingfishers (*Alcedo atthis*) from an electronic waste recycling site and a reference site, South China: Influence of residue levels on the isomeric composition**
L. Mo, J.-P. Wu, X.-J. Luo, Y.-X. Sun, X.-B. Zheng, Q. Zhang, F.-S. Zou, B.-X. Mai
DP residue levels influence the isomeric compositions in the investigated kingfishers.
- 63 Functioning of metal contaminated garden soil after remediation**
M. Jelusic, H. Grcman, D. Vodnik, M. Suhadolc, D. Lestan
EDTA soil washing effectively removed toxic metals and reduced their transfer from the soil to plant roots but did not prevent their accumulation in leaves.
- 71 Antibiotics in the offshore waters of the Bohai Sea and the Yellow Sea in China: Occurrence, distribution and ecological risks**
R. Zhang, J. Tang, J. Li, Q. Zheng, D. Liu, Y. Chen, Y. Zou, X. Chen, C. Luo, G. Zhang
Some antibiotics were ubiquitous in the offshore waters of the Bohai Sea and the Yellow Sea in China and posed medium or low ecological risks to some sensitive organisms.
- 78 Mannitol can mitigate negative effects of simulated acid mist and fluoranthene in juvenile Japanese red pine (*P. densiflora* Sieb. et Zucc.)**
I. Oguntimhin, S. Bandai, H. Sakugawa
Mannitol could be useful for protecting pine trees and other plants from air pollutants.

Continued on inside back cover

CONTENTS—Continued from outside back cover

- 85 Total resistance of native bacteria as an indicator of changes in the water environment**
M. Harnisz
The presence of doxycycline-resistant bacteria in rivers can be a robust indicator of anthropogenic stress.
- 93 Enhanced desorption of PCB and trace metal elements (Pb and Cu) from contaminated soils by saponin and EDDS mixed solution**
M. Cao, Y. Hu, Q. Sun, L. Wang, J. Chen, X. Lu
Significant synergistic effect on Pb, Cu and PCB desorption were achieved with the mixed solution of saponin and EDDS.
- 100 Calibration and field evaluation of polar organic chemical integrative sampler (POCIS) for monitoring pharmaceuticals in hospital wastewater**
E. Bailly, Y. Levi, S. Karolak
After calibration in tap water and hospital wastewater, POCIS were used to monitor pharmaceuticals in hospital sewage and were compared to TWA sampling.
- 106 Deposition and release kinetics of nano-TiO₂ in saturated porous media: Effects of solution ionic strength and surfactants**
I.G. Godinez, C.J.G. Darnault, A.P. Khodadoust, D. Bogdan
Previously deposited nano-TiO₂ serve as preferential sites for subsequent deposition and changes in solution chemistry cause nanodeposits to release a portion of nano-TiO₂.
- 114 Risk spreading, habitat selection and division of biomass in a submerged clonal plant: Responses to heterogeneous copper pollution**
X. Yan, H. Wang, Q. Wang, L.G. Rudstam
*Cu can spread among *V. natans* clones and the clones grow randomly and relative independent in heterogeneous Cu-contaminated sediment.*
- 121 Antioxidant defense system responses and DNA damage of earthworms exposed to Perfluorooctane sulfonate (PFOS)**
D. Xu, C. Li, Y. Wen, W. Liu
Perfluorooctane sulfonate (PFOS) induced oxidative stress and DNA damage in earthworms.
- 128 Impact of arsenic on uptake and bio-accumulation of antimony by arsenic hyperaccumulator *Pteris vittata***
K. Müller, B. Daus, J. Mattusch, D. Vetterlein, I. Merbach, R. Wennrich
*Roots of *Pteris vittata* cultivated under green-house conditioned in substrates containing antimonate accumulates Sb(V). This was promoted by As(V).*
- 134 Behaviour of Au-citrate nanoparticles in seawater and accumulation in bivalves at environmentally relevant concentrations**
C.A. García-Negrete, J. Blasco, M. Volland, T.C. Rojas, M. Hampel, A. Lapresta-Fernández, M.C. Jiménez de Haro, M. Soto, A. Fernández
*Concentration-dependent physico-chemical evolution of citrate-stabilized AuNPs in sea water media, and their uptake and sub-cellular location in *Ruditapes philippinarum*.*
- 142 Chronic effects of lead (Pb) on bone properties in red deer and wild boar: Relationship with vitamins A and D₃**
J. Rodríguez-Estival, P. Álvarez-Lloret, A.B. Rodríguez-Navarro, R. Mateo
The effect of Pb on vitamin A status could be involved in disturbances on bone biomineralization in red deer and wild boar living in Pb-polluted mining areas.
- 150 Bioavailability of nanoparticulate hematite to *Arabidopsis thaliana***
Y. Marusenko, J. Shipp, G.A. Hamilton, J.L.L. Morgan, M. Keebaugh, H. Hill, A. Dutta, X. Zhuo, N. Upadhyay, J. Hutchings, P. Herckes, A.D. Anbar, E. Shock, H.E. Hartnett
*Synthesized iron nanoparticles were not bioavailable to *Arabidopsis thaliana* in agar nutrient media.*
- 157 Ingestion of metal-nanoparticle contaminated food disrupts endogenous microbiota in zebrafish (*Danio rerio*)**
D.L. Merrifield, B.J. Shaw, G.M. Harper, I.P. Saoud, S.J. Davies, R.D. Handy, T.B. Henry
*Dietary exposure to manufactured Cu- and Ag-nanoparticles caused unique changes in endogenous gut microbiota in zebrafish *Danio rerio*.*
- 164 Bioaccumulation of polybrominated diphenyl ethers and several alternative halogenated flame retardants in a small herbivorous food chain**
Y.-Z. She, J.-P. Wu, Y. Zhang, Y. Peng, L. Mo, X.-J. Luo, B.-X. Mai
A modest biomagnification of PBDEs and DP occurred from rice plant to apple snails.
- 171 Breeding performance of blue tits (*Cyanistes caeruleus ultramarinus*) in relation to lead pollution and nest failure rates in rural, intermediate, and urban sites in Algeria**
Z. Brahmia, R. Scheifler, N. Crini, S. Maas, P. Giraudoux, S. Benyacoub
*Urbanization modifies the nestling morphology and breeding success of populations of the Algerian blue tit (*Cyanistes caeruleus ultramarinus*) through mechanisms other than lead pollution.*
- 179 Magnetic susceptibility measurements to detect coal fly ash from the Kingston Tennessee spill in Watts Bar Reservoir**
E.A. Cowan, K.C. Seramur, S.J. Hageman
An application of magnetic susceptibility for tracking the distribution of coal fly ash within a river system after the 2008 TVA spill at Kingston, Tennessee.

- 189 National inventory of alkylphenol ethoxylate compounds in U.S. sewage sludges and chemical fate in outdoor soil mesocosms**
A.K. Venkatesan, R.U. Halden
First study providing national inventories of alkylphenol surfactants in U.S. sewage sludges (SS), shows significant release of chemicals to U.S. soils through SS land application.
- 194 Photolytic debromination pathway of polybrominated diphenyl ethers in hexane by sunlight**
H. Wei, Y. Zou, A. Li, E.R. Christensen, K.J. Rockne
The PBDE debromination products and pathways identified in this work will assist in future studies on their environmental fate.
- 201 Metabolomic analysis of *Cryptosporidium parvum* oocysts in water: A proof of concept demonstration**
D.J. Beale, D. Marney, D.R. Marlow, P.D. Morrison, M.S. Dunn, C. Key, E.A. Palombo
- 204 Impacts of fish farm pollution on ecosystem structure and function of tropical headwater streams**
R.d.S. Rosa, A.C.F. Aguiar, I.G. Boëchat, B. Gücker
Moderate water pollution by small fish farms caused considerable eutrophication responses in tropical headwater streams.
- 214 Metabolism of the polycyclic musk galaxolide and its interference with endogenous and xenobiotic metabolizing enzymes in the European sea bass (*Dicentrarchus labrax*)**
D. Fernandes, G. Dimastrogiovanni, M. Blázquez, C. Porte
HCB is actively metabolised by sea bass and acts as a weak inhibitor of the synthesis of oxyandrogens in gonads of male fish.
- 222 Investigation of gold nanoparticles uptake and their tissue level distribution in rice plants by laser ablation-inductively coupled-mass spectrometry**
J. Koelmel, T. Leland, H. Wang, D. Amarasiriwardena, B. Xing
The tissue level uptake and spatial distribution of engineered gold nanoparticles (AuNP) by rice plants was demonstrated by LA-ICP-MS bioimaging.
- 229 Steady-state mass balance model for mercury in the St. Lawrence River near Cornwall, Ontario, Canada**
C.R. Lessard, A.J. Poulain, J.J. Ridal, J.M. Blais
A steady-state mass balance model is presented for mercury species in the St. Lawrence River near Cornwall, Ontario.
- 236 Distillation fraction-specific ecotoxicological evaluation of a paraffin-rich crude oil**
E. Erlacher, A.P. Loibner, R. Kendler, K.E. Scherr
A differentiated consideration of the prevailing crude oil distillation fractions and of model soil properties employed for ecotoxicity testing should be included into the risk assessment of crude oil contaminated sites.
- 244 Effects of ozone on crops in north-west Pakistan**
M.N. Ahmad, P. Bükér, S. Khalid, L. Van Den Berg, H.U. Shah, A. Wahid, L. Emberson, S.A. Power, M. Ashmore
Ozone concentrations in NW Pakistan have adverse effects on sensitive crop species.
- 250 Reducing the bioavailability of PCBs in soil to plant by biochars assessed with triolein-embedded cellulose acetate membrane technique**
Y. Wang, Y.-J. Wang, L. Wang, G.-D. Fang, L. Cang, H.M.S.K. Herath, D.-M. Zhou
The reduced PCBs concentrations in plant roots by biochars show good linear relationship with those in TECAM.
- 257 Incorporation and mineralization of TNT and other anthropogenic organics by natural microbial assemblages from a small, tropical estuary**
M.T. Montgomery, R.B. Coffin, T.J. Boyd, C.L. Osburn
Incorporation of TNT ring carbon into biomass of natural microbial assemblages occurs at a more rapid rate than mineralization of that ring carbon to CO₂.
- 265 Occurrence of ectoparasitocides in Australian beef cattle feedlot wastes**
H.M. Coleman, T. Trinh, N. Le-Minh, M. Klein, D.J. Roser, R.W. Tucker, R.M. Stuetz, G. Peters, S.J. Khan
The first reported study on concentrations of ectoparasitocides in waste from Australian beef cattle feedlot operations.
- 273 Effects of plant species on soil microbial processes and CH₄ emission from constructed wetlands**
Y. Wang, H. Yang, C. Ye, X. Chen, B. Xie, C. Huang, J. Zhang, M. Xu
The polyculture constructed wetland has the higher contribution to global warming.
- 279 Impact of elevated CO₂ and elevated O₃ on *Beta vulgaris* L.: Pigments, metabolites, antioxidants, growth and yield**
S. Kumari, M. Agrawal, S. Tiwari
Predicted levels of CO₂ have greater ameliorative potential against negative effects of elevated ozone compared to present day CO₂ against ambient ozone.
- 289 Characterization and influence of biochars on nitrous oxide emission from agricultural soil**
Z. Wang, H. Zheng, Y. Luo, X. Deng, S. Herbert, B. Xing
Biochars (200–600 °C) produced from giant reed reduced N₂O emissions from a soil due to enhanced N immobilization and decreased denitrification.
- 297 Risk assessment of metals in road-deposited sediment along an urban–rural gradient**
H. Zhao, X. Li
A new index integrating road-deposited sediment characteristics and potential ecological risk was developed to assess metal risk along the urban–rural gradient.

305 Occurrence of 95 pharmaceuticals and transformation products in urban groundwaters underlying the metropolis of Barcelona, Spain

R. López-Serna, A. Jurado, E. Vázquez-Suñé, J. Carrera, M. Petrović, D. Barceló

Ninety-five pharmaceuticals and transformation products are monitored in the groundwater from Barcelona metropolitan area.

Retraction

316 Retraction note: Dispersal of the radionuclide caesium-137 (^{137}Cs) from point sources in the Barents and Norwegian Seas and its potential contamination of the Arctic marine food chain: Coupling numerical ocean models with geographical fish distribution data [Environ. Pollut. 164 (2012) 1–10]

H.E. Heldal, F. Vikebø, G.O. Johansen