

CONTENTS

Volume 175 April 2013

- 1 Are plants useful as accumulation indicators of metal bioavailability?**
E. Remon, J.-L. Bouchardon, M. Le Guédard, J.-J. Bessoule, C. Conord, O. Faure
Plant communities as accumulation bioindicators of trace metal bioavailability in contaminated soils.
- 8 Effects of dietary selenium on the health and survival of captive wintering lesser scaup**
C. Brady, S. Petrie, M. Schummer, S. Badzinski, N. Belzile, Y.-W. Chen
Selenium alone was not the cause of the continental decline in scaup populations and other plausible hypothesis should be explored.
- 16 Occurrence of chlorinated paraffins in house dust samples from Bavaria, Germany**
B. Hilger, H. Fromme, W. Völkel, M. Coelhan
Medium-chain chlorinated paraffins (MCCPs) may be regarded as being present in house dust from Western European countries.
- 22 Antibiotics in the surface water of the Yangtze Estuary: Occurrence, distribution and risk assessment**
C. Yan, Y. Yang, J. Zhou, M. Liu, M. Nie, H. Shi, L. Gu
Widespread occurrence of antibiotic residues was investigated in the surface water of Yangtze Estuary over four seasons.
- 30 The effects of low-level prenatal lead exposure on birth outcomes**
X. Xie, G. Ding, C. Cui, L. Chen, Y. Gao, Y. Zhou, R. Shi, Y. Tian
Prenatal low-level exposure to lead was associated with decreasing birth weight and length.
- 35 1-D and 2-D NMR-based metabolomics of earthworms exposed to endosulfan and endosulfan sulfate in soil**
J. Yuk, M.J. Simpson, A.J. Simpson
*NMR-based metabolomics was used to investigate the toxicity and toxic mode of action of endosulfan, and its main degradation product, to the earthworm *Eisenia fetida* in soil.*
- 45 Detecting benthic community responses to pollution in estuaries: A field mesocosm approach**
A.L. O'Brien, M.J. Keough
Large-scale experimental study that demonstrates the ecological effects of pollution on estuarine communities.
- 56 Personal exposure to benzene of selected population groups and impact of commuting modes in Ho Chi Minh, Vietnam**
T.T.N. Lan, N.Q. Liem, N.T.T. Binh
Average non-occupational personal exposure to benzene in Ho Chi Minh is ~18 µg/m³, most of exposure is due to commuting. Exposure during commuting by motorcycle is 116 µg/m³.
- 64 Reduced bioaccumulation of PAHs by *Lactuca sativa* L. grown in contaminated soil amended with sewage sludge and sewage sludge derived biochar**
S. Khan, N. Wang, B.J. Reid, A. Freddo, C. Cai
Sewage sludge biochar significantly increased biomass yield and was more effective than sewage sludge in reducing PAH bioaccumulation from contaminated soil.
- 69 New approach to the ecotoxicological risk assessment of artificial outdoor sporting grounds**
O. Krüger, U. Kalbe, E. Richter, P. Egeler, J. Römbke, W. Berger
Column tests of complete artificial sporting ground assemblies may be a useful tool for their ecotoxicological risk assessment.
- 75 Reduction in the earthworm metabolomic response after phenanthrene exposure in soils with high soil organic carbon content**
J.R. McKelvie, M.W. Åslund, M.A. Celejewski, A.J. Simpson, M.J. Simpson
¹H NMR-based metabolomics is able to evaluate phenanthrene bioavailability to earthworms in soil.

- 82 **Stomata are less responsive to environmental stimuli in high background ozone in *Dactylis glomerata* and *Ranunculus acris***
S. Wagg, G. Mills, F. Hayes, S. Wilkinson, W.J. Davies
*Elevated background ozone reduces stomatal sensitivity to environmental stimuli in *Dactylis glomerata* and *Ranunculus acris*.*
- 92 **Dietary exposure of Antarctic krill to p,p'-DDE: Uptake kinetics and toxicological sensitivity in a key polar species**
A.H. Poulsen, P.F. Landrum, S. Kawaguchi, S.M. Bengtson Nash
This research demonstrates the importance of diet for p,p'-DDE accumulation in Antarctic krill and presents critical body residues comparable to values known for temperate species.
- 100 **Mycorrhizal colonization affects the elemental distribution in roots of Ni-hyperaccumulator *Berkheya coddii* Roessler**
E. Orłowska, W. Przybyłowicz, D. Orłowski, N.P. Mongwaketsi, K. Turnau, J. Mesjasz-Przybyłowicz
*Arbuscular mycorrhizal fungi significantly affect the distribution of elements in roots of Ni-hyperaccumulator *Berkheya coddii*.*
- 110 **Isosteric heats of sorption and desorption of phenanthrene in soils and carbonaceous materials**
G. Wang, P. Grathwohl
Temperature-dependent sorption isotherms of phenanthrene on carbonaceous materials are used to determine isosteric heat for analyzing the relative effect of adsorption and partitioning.
- 117 **Effects of linuron on a rooted aquatic macrophyte in sediment-dosed test systems**
H. Burešová, S.J.H. Crum, J.D.M. Belgers, P.I. Adriaanse, G.H.P. Arts
First sediment-spiked pesticide toxicity test with a rooted macrophyte.
- 125 **Assessing microbial carbon sources and potential PAH degradation using natural abundance ¹⁴C analysis**
N. Mahmoudi, R.R. Fulthorpe, L. Burns, S. Mancini, G.F. Slater
Natural abundance ¹⁴C analysis revealed that microbial carbon sources were derived primarily from natural organic matter rather than PAHs in contaminated soils.
- 131 **Global climate change: Impact of diurnal temperature range on mortality in Guangzhou, China**
J. Yang, H.-Z. Liu, C.-Q. Ou, G.-Z. Lin, Q. Zhou, G.-C. Shen, P.-Y. Chen, Y. Guo
Diurnal temperature range is an independent risk factor of mortality particularly for female, the elderly and people with low education level.
- 137 **Persistent organic pollutants and methoxylated polybrominated diphenyl ethers in different tissues of white-tailed eagles (*Haliaeetus albicilla*) from West Greenland**
V.L.B. Jaspers, C. Sonne, F. Soler-Rodriguez, D. Boertmann, R. Dietz, M. Eens, L.M. Rasmussen, A. Covaci
Significant differences were found between tissues for POPs, but not for MeO-PBDEs, with muscle displaying the highest and adipose tissue the lowest lipid-normalized POP levels.
- 147 **Evaluation of acute and developmental effects of difenoconazole via multiple stage zebrafish assays**
X. Mu, S. Pang, X. Sun, J. Gao, J. Chen, X. Chen, X. Li, C. Wang
The acute toxicity of difenoconazole to the three phases of zebrafish were ordered larvae > adult > embryo, and many drug-induced abnormalities during embryonic and larvae stages were observed.
- 158 **Concentrations and source contributions of particulate organic matter before and after implementation of a low emission zone in Munich, Germany**
R.M. Qadir, G. Abbaszade, J. Schnelle-Kreis, J.C. Chow, R. Zimmermann
Significant effect of implementation of LEZ was detected on the concentrations of EC and some of the POC after that; the EPA-PMF 3.0 model was used for studying the source apportionment.
- 168 **Effects of freeze-thaw cycling on metal-phosphate formation and stability in single and multi-metal systems**
E.G. Hafsteinsdóttir, D.A. White, D.B. Gore
Competing ions, associated with salts or other metals, do not prevent the formation of lead phosphates but slow or hinder the formation of zinc and copper phosphates in multi-metal systems.
- 178 **Acknowledgement to Referees**