

ORIGINAL ARTICLES**N. A. TINSLEY, R. E. ESTES, M. E. GRAY**

Validation of a nested error component model to estimate damage caused by corn rootworm larvae

161–169**C. R. R. HOOKS, J. HINDS, E. ZOBEL, T. PATTON**

Impact of crimson clover dying mulch on two eggplant insect herbivores

170–180**J. HOLUŠA, K. LUKÁŠOVÁ, R. WEGENSTEINER, W. GRODZKI, M. PERNEK, J. WEISER**Pathogens of the bark beetle *Ips cembrae*: microsporidia and gregarines also known from other *Ips* species**181–187****K. LUKÁŠOVÁ, J. HOLUŠA, M. TURČÁNI**Pathogens of *Ips amitinus*: new species and comparison with *Ips typographus***188–196****D. F. WARD, M. C. STANLEY**

Site occupancy and detection probability of Argentine ant populations

197–203**J. O. NYASANI, R. MEYHÖFER, S. SUBRAMANIAN, H.-M. POEHLING**Feeding and oviposition preference of *Frankliniella occidentalis* for crops and weeds in Kenyan French bean fields**204–213****A. KNIGHT, D. LIGHT, V. CHEBNY**

Monitoring codling moth (Lepidoptera: Tortricidae) in orchards treated with pear ester and sex pheromone combo dispensers

214–224**W. L. YEE**Preferences by *Rhagoletis indifferens* (Diptera, Tephritidae) for rectangles of various yellow colours and fluorescence**225–233****SHORT COMMUNICATION****D. PIESIK, D. ROCHAT, K. J. DELANEY, F. MARION-POLL**

Orientation of European corn borer first instar larvae to synthetic green leaf volatiles

234–240