

CONTENTS

The influence of farming technique on cropland: A new approach for the Ecological Footprint N. Passeri (Viterbo, Italy), M. Borucke (Oakland, CA, USA), E. Blasi, S. Franco (Viterbo, Italy) and E. Lazarus (Oakland, CA, USA).....	1
Development of pollution indices for the middle section of the Lower Seyhan Basin (Turkey) M. Golge, F. Yenilmez and A. Aksoy (Ankara, Turkey).....	6
Freshwater sediment pesticide biodegradation potential as an ecological indicator of microbial recovery following a decrease in chronic pesticide exposure: A case study with the herbicide diuron S. Pesce, C. Margoum (Villeurbanne, France), N. Rouard (Dijon France), A. Foulquier (Dijon, France) and F. Martin-Laurent (Dijon, France)	18
Methodological considerations on the coastal and transitional benthic indicators proposed for the Water Framework Directive I. Karakassis, P.D. Dimitriou, N. Papageorgiou, E.T. Apostolaki, N. Lampadariou (Crete, Greece) and K.D. Black (Scotland, UK).....	26
Quantifying uncertainties in biologically-based water quality assessment: A pan-European analysis of lake phytoplankton community metrics S.J. Thackeray (Lancaster, UK), P. Nöges (Rannu, Estonia), M.J. Dunbar (Wallingford, UK), B.J. Dudley (Midlothian, UK), B. Skjelbred (Oslo, Norway), G. Morabito (Verbania Pallanza, Italy), L. Carvalho (Midlothian, UK), G. Phillips (Reading, UK), U. Mischke (Berlin, Germany), J. Catalan (Blanes, Spain), C. de Hoyos (Madrid, Spain), C. Laplace (France), M. Austoni (Verbania Pallanza, Italy), B.M. Padedda (Sassari, Italy), K. Maileht (Rannu, Estonia), A. Pasztaleniec (Kolektorska, Poland), M. Järvinen (Jyväskylä, Finland), A.L. Solheim (Oslo, Norway) and R.T. Clarke (Dorset, UK).....	34
The usefulness of large body-size macroinvertebrates in the rapid ecological assessment of Mediterranean lagoons M. Pinna, G. Marini, I. Rosati (Lecce, Italy), J.M. Neto, J. Patrício, J.C. Marques (Coimbra, Portugal) and A. Basset (Lecce, Italy).....	48
The emergy ecological footprint for small fish farm in China S. Zhao, K. Song, F. Gui, H. Cai, W. Jin and C. Wu (Zhoushan, China).....	62
Macroinvertebrate communities as bioindicators of water quality in conventional and organic irrigated rice fields in Guanacaste, Costa Rica F. Rizo-Patrón V. (Guanacaste, Costa Rica), A. Kumar (San Jose, Costa Rica), M.B. McCoy Colton (Costa Rica), M. Springer (San José, Costa Rica) and F.A. Trama (Guanacaste, Costa Rica)	68
A tool to assess the ecological condition of tropical high Andean streams in Ecuador and Peru: The IMEERA index C. Villamarín, M. Rieradevall (Barcelona, Spain), M.J. Paul, M.T. Barbour (Owings Mills, MD, USA) and N. Prat (Barcelona, Spain).....	79
Bole epiphytic lichens as potential indicators of environmental change in subtropical forest ecosystems in southwest China S. Li (Kunming, China and Beijing, China), W.-Y. Liu (Kunming, China and Perth, Australia) and D.-W. Li (Jingdong, China).....	93
A scientometric assessment of 30 years of the Index of Biotic Integrity in aquatic ecosystems: Applications and main flaws R. Ruaro (Cascavel, Brazil) and É.A. Gubiani (Cascavel, Brazil and Toledo, Brazil)	105
Habitat and hydrology condition indices for the upper Mississippi, Missouri, and Ohio rivers D.L. Taylor, D.W. Bolgrien, T.R. Angradi, M.S. Pearson and B.H. Hill (Duluth, MN, USA).....	111

(Contents continued on inside back cover)

Can an Environmental Indicator valid both at the local and global scales be derived on a thermodynamic basis? E. Sciubba (Roma, Italy)	125
Toward river health assessment using species distributional modeling I. Grouns (Armidale, Australia), M. Rourke (Narrandera, Australia) and D. Gilligan (Batemans Bay, Australia)	138
More than the mean-A note on heterogeneity aspects in the assessment of water footprints R. Finger (Wageningen, The Netherlands)	145
Water quality assessment in shrimp culture using an analytical hierarchical process J.J. Carbajal-Hernández (México, Mexico and Puebla, Mexico), L.P. Sánchez-Fernández, L.A. Villa-Vargas (México, Mexico), J.A. Carrasco-Ochoa and J.Fco. Martínez-Trinidad (Puebla, Mexico)	148
Geometric morphology as an alternative for measuring the diversity of fish assemblages M. Farré, V.M. Tuset, F. Maynou, L. Recasens and A. Lombarte (Barcelona, Spain)	159
Development of a macroinvertebrate multimetric index for the assessment of low-land streams in the neotropics J.E. Helson and D.D. Williams (Toronto, Canada)	167
Urban ecosystem services assessment along a rural-urban gradient: A cross-analysis of European cities N. Larondelle and D. Haase (Germany)	179
Application of selected partial order tools to analyze fate and toxicity indicators of environmentally hazardous chemicals L. Carlsen (Roskilde, Denmark and Almaty Kazakhstan), R. Bruggemann (Berlin, Germany) and Y. Sailaukhanuly (Almaty Kazakhstan)	191
Deriving land surface phenology indicators from CO ₂ eddy covariance measurements A. Gonsamo, J.M. Chen (Toronto, Canada) and P. D'Odorico (Zurich, Switzerland)	203
The intercalibration of the Hellenic Evaluation System for Mediterranean rivers of type R-M1 and R-M2 at Northern and Central Greece Ch. Ntislidou, V. Artemiadou, L. Kanli (Thessaloniki, Greece), A. Buffagni (Italy) and M. Lazaridou (Thessaloniki, Greece)	208
Defining past ecological status and <i>in situ</i> reference conditions using benthic foraminifera: A case study from the Oslofjord, Norway J.K. Dolven, E. Alve, B. Rygg and J. Magnusson (Oslo, Norway)	219
Reference diatom assemblage response to restoration of an acid mine drainage stream J.B. Gray and M.L. Vis (Athens, OH, USA)	234
Food web indicators under the Marine Strategy Framework Directive: From complexity to simplicity? I. Rombouts, G. Beaugrand (Wimereux, France), X. Fizzala (Marseille, France), F. Gaill (Paris, France), S.P.R. Greenstreet (Aberdeen, UK), S. Lamare (Paris, France and La Rochelle, France), F. Le Loc'h (Sète, France), A. McQuatters-Gollop (Plymouth, UK), B. Mialet (La Rochelle, France), N. Niquil (Paris, France; La Rochelle, France and Caen, France), J. Perceley (La Défense, France), F. Renaud (Marseille, France and Sète, France), A.G. Rossberg (Suffolk, UK and Belfast, UK) and J.P. Féral (Marseille, France)	246
Selecting ecological indicators to compare maintenance costs related to the compensation of damaged ecosystem services A.-C. Vaissière, H. Levrel, C. Hily and D. Le Guyader (Plouzané, France)	255
Robustness and sensitivity of weighting and aggregation in constructing composite indices M.J. Dobbie (Brisbane, Australia) and D. Dail (Brisbane, Australia and Corvallis, OR, USA)	270
The maximum empower principle: An invisible hand controlling the self-organizing development of forest plantations in south China L. Li (Guangzhou, China and Shenzhen, China), H. Lu (Guangzhou, China), D.R. Tilley (College Park, MD, USA), H. Ren and W. Shen (Guangzhou, China)	278
Development of environmental performance indicators supported by an environmental information system: Application to the Norwegian defence sector O. Myhre, K. Fjellheim, H. Ringnes, T. Reistad, K.S. Longva (Kjeller, Norway) and T.B. Ramos (Caparica, Portugal)	293
Indicator plant species selection for monitoring the impact of climate change based on prediction uncertainty M. Higa, K. Nakao, I. Tsuyama, E. Nakazono, M. Yasuda (Ibaraki, Japan), T. Matsui (Hokkaido, Japan) and N. Tanaka (Ibaraki, Japan)	307
Composite indicator for the assessment of sustainability: The case of Cuban nature-based tourism destinations V. Pérez (Pinar del Río, Cuba and Seville, Spain), F. Guerrero (Seville, Spain), M. González, F. Pérez and R. Caballero (Málaga, Spain)	316
Provincial level-based energy evaluation of crop production system and development modes in China J. Tao, M. Fu, X. Zheng, J. Zhang and D. Zhang (Beijing, China)	325
Multivariate geostatistical methods for analysis of relationships between ecological indicators and environmental factors at multiple spatial scales M.C. Ribeiro, P. Pinho (Lisboa, Portugal), E. Llop (Barcelona, Spain), C. Branquinho, A.J. Sousa and M.J. Pereira (Lisboa, Portugal)	339
Relationships of hydrology and life-history strategies on the spatio-temporal habitat utilisation of fish in European temperate river floodplains A. Scharbert and J. Borcharding (Köln, Germany)	348

(Contents continued from inside back cover)

A non-phylogenetic alpha diversity approach on prokaryotic community structure in aquatic systems E. Smeti (Mytilene, Greece), K.Ar. Kormas (Nea Ionia, Greece) and S. Spatharis (Mytilene, Greece)	361
The possible combined effects of land-use changes and climate conditions on the spatial-temporal patterns of primary production in a natural protected area I. Petrosillo, T. Semeraro, N. Zaccarelli, R. Aretano and G. Zurlini (Lecce, Italy).....	367
Developing an indicator system to foster sustainability in strategic planning in China: A case study of Pudong New Area, Shanghai Y. Wang (Shanghai, China), K.-c. Lam (Shatin, China), M.K. Harder, W.-c. Ma and Q. Yu (Shanghai, China).....	376
New multi-metric Salt Marsh Sediment Microbial Index (SSMI) application to salt marsh sediments ecological status assessment B. Duarte, J. Freitas (Lisbon, Portugal), T. Couto (Coimbra, Portugal), J. Valentim, J.M. Dias, H. Silva (Aveiro, Portugal), J.C. Marques (Coimbra, Portugal) and I. Caçador (Lisbon, Portugal)	390
Development of a structurally dynamic model for ecosystem health prognosis of Baiyangdian Lake, China F. Xu, Z. Yang, B. Chen and Y. Zhao (Beijing, China).....	398
A causal examination of the effects of confounding factors on multimetric indices D.R. Schoolmaster Jr. (USA), J.B. Grace (Lafayette, LA, USA), E.W. Schweiger (Fort Collins, CO, USA), B.R. Mitchell (Woodstock, VT, USA) and G.R. Guntenspergen (Laurel, MD, USA)	411
Extending the viability theory framework of resilience to uncertain dynamics, and application to lake eutrophication C. Rougé, J.-D. Mathias and G. Deffuant (Aubière, France)	420
An empirical review of cultural ecosystem service indicators M. Hernández-Morcillo, T. Plieninger (Berlin, Germany) and C. Bieling (Freiburg, Germany)	434
Ranking field site management priorities according to their metal transfer to snails B. Pauget, F. Gimbert, M. Coeurdassier, N. Crini (Besançon, France), G. Pérès (Rennes, France), O. Faure (St-Etienne, France), F. Douay (Lille, France), A. Hitmi (Clermont-Ferrand, France), T. Beguiristain (France), A. Alaphilippe (France), M. Guernion (Rennes, France), S. Houot (France), M. Legras (Mont-Saint-Aignan, France), J.-F. Vian (Lyon, France), M. Hedde (Versailles, France), A. Bispo, C. Grand (Angers, France) and A. de Vaufleury (Besançon, France)	445
Regional nutrient thresholds in wadeable streams of New York State protective of aquatic life A.J. Smith (Troy, NY, USA), R.L. Thomas (Philadelphia, PA, USA), J.K. Nolan (Schenectady, NY, USA), D.J. Velinsky, S. Klein (Philadelphia, PA, USA) and B.T. Duffy (Troy, NY, USA).....	455
New synthetic indicators to assess community resilience and restoration success R. Jaunatre, E. Buisson, I. Muller, H. Morlon (France), F. Mesléard and T. Dutoit (France)	468
Greenhouse gas emissions and the technical efficiency of dairy farmers O.K. Shortall (Leicestershire, UK) and A.P. Barnes (Scotland, UK).....	478
Application of European biomonitoring techniques in China: Are they a useful tool? C. Pignata (Torino, Italy), S. Morin (Cestas, France), A. Scharl (Toulouges, France), D. Traversi, T. Schilirò, R. Degan (Torino, Italy), P. Bartley (Beijing, PR China), M. Tu, H. Liu (Wuhan, PR China), F. Peres (Toulouges, France), M. Coste (Cestas, France), W. Liu (Guangzhou, PR China) and G. Gilli (Torino, Italy)	489
Impacts of sewage outfalls on rocky shores: Incorporating scale, biotic assemblage structure and variability into monitoring tools N.E. O'Connor (Northern Ireland, UK and Dublin, Ireland)	501
Development of a multimetric index based on benthic macroinvertebrates for the assessment of natural wetlands in Southwest Ethiopia S.T. Mereta (Ghent, Belgium and Jimma, Ethiopia), P. Boets (Ghent, Belgium), L. De Meester (Leuven, Belgium) and P.L.M. Goethals (Ghent, Belgium)	510
Mathematical convergences of biodiversity indices B. Bandeira, J.-L. Jamet, D. Jamet and J.-M. Ginoux (France)	522
Hotspot and relationship identification in multiple landscape services: A case study on an area with intensive human activities J. Wu, Z. Feng (Shenzhen, China and Beijing, China), Y. Gao and J. Peng (Beijing, China).....	529
Assessment of <i>Lemna gibba</i> L. (duckweed) as a potential ecological indicator for contaminated aquatic ecosystem by boron mine effluent H. Böcük, A. Yakar (Eskişehir, Turkey) and O.C. Türker (Aksaray, Turkey)	538
Talitrid and Tylid crustaceans bioecology as a tool to monitor and assess sandy beaches' ecological quality condition S.C. Gonçalves (Coimbra, Portugal and Peniche, Portugal), P.M. Anastácio (Évora, Portugal) and J.C. Marques (Coimbra, Portugal)	549

- Colombini, I., Aloia, A., Bouslama, M.F., El Gtari, M., Fallaci, M., Ronconi, L., Scapini, F., Chelazzi, L., 2002. Small-scale spatial and seasonal differences in the distribution of beach arthropods on the Northwestern Tunisian coast. Are species evenly distributed along the shore? *Mar. Biol.* 140, 1001–1012.
- Colombini, I., Chelazzi, L., 2003. Influence of marine allochthonous input on sandy beach communities. *Oceanogr. Mar. Biol. Ann. Rev.* 41, 115–159.
- Colombini, I., Fallaci, M., Milanesi, F., Scapini, F., Chelazzi, L., 2003. Comparative diversity analysis in sandy littoral ecosystems of the western Mediterranean. *Estuar. Coast. Shelf S* 58, 93–104.
- Defeo, O., Cardoso, R.S., 2002. Macroecology of population dynamics and life history traits of the mole crab *Emerita brasiliensis* in Atlantic sandy beaches of South America. *Mar. Ecol. Prog. Ser.* 239, 169–179.
- Defeo, O., McLachlan, A., 2005. Patterns, processes and regulatory mechanisms in sandy beach macrofauna: a multi-scale analysis. *Mar. Ecol. Prog. Ser.* 295, 1–20.
- Dunn, M.A., Hogg, J.C., Kelly, A., 2005. Two cues for sex determination in *Gammarus duebeni*: adaptative variation in environmental sex determination? *Limnol. Oceanogr.* 50 (1), 346–353.
- Gheskiere, T., Vincx, M., Weslawski, J.M., Scapini, F., Degraer, S., 2005. Meiofauna as a descriptor of tourism-induced changes at sandy beaches. *Mar. Environ. Res.* 60, 245–265.
- Gonçalves, S.C., Marques, J.C., Pardal, M.A., Bouslama, M.F., El Gtari, M., Charfi-Cheikhrouha, F., 2003. Comparison of *Talorchestia brito* (amphipoda, talitridae) biology, dynamics, and secondary production in Atlantic (Portugal) and Mediterranean (Tunisia) populations. *Estuar. Coast. Shelf S* 58, 901–916.
- Gonçalves, S.C., Pardal, M.A., Cardoso, P.G., Ferreira, S.M., Marques, J.C., 2005. Biology, population dynamics and secondary production of *Tylos europaeus* (Isopoda, Tylidae) on the western coast of Portugal. *Mar. Biol.* 147, 631–641.
- Gonçalves, S.C., Anastácio, P.M., Pardal, M.A., Cardoso, P.G., Ferreira, S.M., Marques, J.C., 2009. Sandy beach macrofaunal communities on the western coast of Portugal—Is there a steady structure under similar exposed conditions? *Estuar. Coast. Shelf S* 81, 555–568.
- Gonçalves, S.C., Marques, J.C., 2011. The effects of season and wrack subsidy on the community functioning of exposed sandy beaches. *Estuar. Coast. Shelf S* 95, 165–177.
- Gormsen, E., 1997. The impact of tourism on coastal areas. *Geoj.* 42 (1), 39–54.
- Kato, Y., Kobayashi, K., Watanabe, H., Iguchi, T., 2011. Environmental sex determination in the Branchiopod Crustacean *Daphnia magna*: deep conservation of a *doublesex* gene in the sex-determining pathway. *PLoS Genet.* 7 (3), 1–12.
- Kitahara, M., Fujii, K., 1994. Biodiversity and community structure of temperate butterfly species within a gradient of human disturbance: an analysis based on the concept of generalist vs. specialist strategies. *Res. Popul. Ecol.* 36, 187–199.
- Kitahara, M., Sei, K., Fujii, K., 2000. Patterns in the structure of grassland butterfly communities along a gradient of human disturbance: further analysis based on the generalist/specialist concept. *Popul. Ecol.* 42, 135–144.
- Korpelainen, H., 1990. Sex ratios and conditions required for environmental sex determination in animals. *Biol. Rev.* 65, 147–184.
- Lagardère, J.P., 1966. Recherches sur la biologie et l'écologie de la macrofaune des substrats meubles de la côte des Landes et de la côte Basque. *Bull. Cent. Etud. Rech. Sci. Biarritz* 6, 143–209.
- Lercari, D., Defeo, O., 2003. Variation of a sandy beach macrobenthic community along a human-induced environmental gradient. *Estuar. Coast. Shelf S* 58, 17–24.
- Marques, J.C., Anastácio, P.M., 2002. Integration and interpretation of ecological data at population level. In: Scapini, F. (Ed.), *Baseline research for the integrated sustainable management of Mediterranean sensitive coastal ecosystems. A manual for coastal managers, scientists and all those studying coastal processes and management in the Mediterranean*. Istituto Agronomico per l'Oltremare, Florence, pp. 155–164.
- Marques, J.C., Gonçalves, S.C., Pardal, M.A., Chelazzi, L., Colombini, I., Fallaci, M., Bouslama, M.F., El Gtari, M., Charfi-Cheikhrouha, F., Scapini, F., 2003. Comparison of *Talitrus saltator* (Amphipoda, Talitridae) biology, dynamics, and secondary production in Atlantic (Portugal) and Mediterranean (Italy and Tunisia) populations. *Estuar. Coast. Shelf S* 58, 127–148.
- Marvier, M., Kareiva, P., Neubert, M.G., 2004. Habitat destruction, fragmentation, and disturbance promote invasion by habitat generalists in a multispecies metapopulation. *Risk Anal.* 24, 869–878.
- McLachlan, A., Brown, A.C., 2006. *The Ecology of Sandy Shores*, second ed. Elsevier, California.
- Montanari, A., 1995. The Mediterranean region: Europe's summer leisure space. In: Montanari, A., Williams, A.M. (Eds.), *European tourism: Regions, Spaces and Restructuring*. Wiley, Chichester, pp. 41–65.
- Persson, L.E., 1999. Growth and reproduction in two brackish water populations of *Orchestia gammarellus* (Amphipoda: Talitridae) in the Baltic. *J. Crustacean Biol.* 19, 53–59.
- Prato, E., Trono, A., Biandolino, F., 2009. Life history of *Talorchestia deshayesii* (Amphipoda, Talitridae) in the Ionian sandy beach (Southern Italy). *Braz. Arch. Biol. Technol.* 52, 911–922.
- Rodil, I.F., Lastra, M., Sánchez-Mata, A.G., 2006. Community structure and intertidal zonation of the macroinfauna in intermediate sandy beaches in temperate latitudes North coast of Spain. *Estuar. Coast. Shelf S* 67, 267–279.
- Scapini, F., Morgan, E., 2002. Bioassays for estimation of beach stability and ecosystem quality. In: Scapini, F. (Ed.), *Baseline Research for the Integrated Sustainable Management of Mediterranean Sensitive Coastal Ecosystems. A Manual for Coastal Managers, Scientists and All Those Studying Coastal Processes and Management in the Mediterranean*. Istituto Agronomico per l'Oltremare, Florence, pp. 120–122.
- Schlacher, T.A., Richardson, D., McLean, I., 2008. Impacts of off-road vehicles (ORVs) on macrobenthic assemblages on sandy beaches. *Environ. Manage.* 41, 878–892.
- Smith, L.R., 1996. *Ecology and field biology*, fifth ed. Addison-Wesley Educational Publishers, Inc., New York.
- Shalan, I.M., 2005. Sustainable tourism development in the Red Sea of Egypt threats and opportunities. *J. Clean. Prod.* 13, 83–87.
- Van Senus, P., 1988. Reproduction of the sandhopper, *Talorchestia capensis* (Dana) (Amphipoda, Talitridae). *Crustaceana* 55 (1), 93–103.
- Weslawski, J.M., Urban-Malinga, B., Kotwicki, L., Opaliński, K., Szymelfenig, M., Dutkowsk, M., 2000a. Sandy coastlines—are there conflicts between recreation and natural values? *Oceanol. Stud.* 29 (2), 5–18.
- Weslawski, J.M., Stanek, A., Siewert, A., Beer, N., 2000b. The sandhopper (*Talitrus saltator*, Montagu 1808) on the Polish Baltic coast. Is it a victim of increased tourism? *Oceanol. Stud.* 29 (1), 77–87.