

CONTENTS

<i>H. Gunnlaugsdottir and J. T. Tuomisto</i>	1–2	Foreword
<i>P. Goyens</i>	3	Preface
 QALIBRA papers		
<i>A. Hart, J. Hoekstra, H. Owen, M. Kennedy, M. J. Zeilmaker, N. de Jong and H. Gunnlaugsdottir</i>	4–17	Qalibra: A general model for food risk–benefit assessment that quantifies variability and uncertainty
<i>J. Hoekstra, A. Hart, H. Owen, M. Zeilmaker, B. Bokkers, B. Thorgilsson and H. Gunnlaugsdottir</i>	18–29	Fish, contaminants and human health: Quantifying and weighing benefits and risks
<i>M. J. Zeilmaker, J. Hoekstra, J. C. H. van Eijkeren, N. de Jong, A. Hart, M. Kennedy, H. Owen and H. Gunnlaugsdottir</i>	30–34	Fish consumption during child bearing age: A quantitative risk–benefit analysis on neurodevelopment
<i>J. Hoekstra, H. P. Fransen, J. C. H. van Eijkeren, J. Verkaik-Kloosterman, N. de Jong, H. Owen, M. Kennedy, H. Verhagen and A. Hart</i>	35–42	Benefit–risk assessment of plant sterols in margarine: A QALIBRA case study
<i>N. de Jong, J. Verkaik-Kloosterman, H. Verhagen, H. C. Boshuizen, B. Bokkers and J. Hoekstra</i>	43–49	An appeal for the presentation of detailed human derived data for dose–response calculations in nutritional science
 BENERIS papers		
<i>O. Leino, A. K. Karjalainen and J. T. Tuomisto</i>	50–58	Effects of docosahexaenoic acid and methylmercury on child’s brain development due to consumption of fish by Finnish mother during pregnancy: A probabilistic modeling approach
<i>O. Leino, H. Kiviranta, A. K. Karjalainen, C. Kronberg-Kippilä, H. Sinkko, E. H. Larsen, S. Virtanen and J. T. Tuomisto</i>	59–69	Pollutant concentrations in placenta
<i>A. K. Karjalainen, A. Hallikainen, T. Hirvonen, H. Kiviranta, M. Knip, C. Kronberg-Kippilä, O. Leino, O. Simell, H. Sinkko, J. T. Tuomisto, R. Veijola, E.-R. Venäläinen and S. M. Virtanen</i>	70–77	Estimated intake levels for Finnish children of methylmercury from fish