

WEED RESEARCH

Insights

- 159 *A R Collins, D Thalmann & H Müller-Schärer* Cytotypes of *Centaurea stoebe* found to differ in root growth using growth pouches
- 164 *T Shilo, B Rubin, J E Ephrath & H Eizenberg* Continuous non-destructive monitoring of *Cyperus rotundus* development using a minirhizotron

Research Papers

- 169 *B W Arentoft, A Ali, J C Streibig & C Andreasen* A new method to evaluate the weed-suppressing effect of mulches: a comparison between spruce bark and cocoa husk mulches
- 176 *K Tsuji, M Hosokawa, S Morita, R Miura & T Tominaga* Resistance to paraquat in *Mazus pumilus*
- 183 *L Andersson, U Boström, J Forkman, I Hakman, J Liew & E Magnuski* Sprouting capacity from intact root systems of *Cirsium arvense* and *Sonchus arvensis* decrease in autumn
- 192 *N Orloff, Z J Miller, F D Menalled, M E Burrows & J M Mangold* Joint effects of biotic and abiotic stressors on winter wheat suppression of *Bromus tectorum*
- 201 *L N Kolb & E R Gallandt* Modelling population dynamics of *Sinapis arvensis* in organically grown spring wheat production systems
- 213 *C Zambrano-Navea, F Bastida & J L Gonzalez-Andujar* A hydrothermal seedling emergence model for *Conyza bonariensis*
- 221 *H Costa, V Medeiros, E B Azevedo & L Silva* Evaluating ecological-niche factor analysis as a modelling tool for environmental weed management in island systems