

CONTENTS

Volume 178 July 2013

- 1 Plant Species Sensitivity Distributions for ozone exposure**
T.M.W.J. van Goethem, L.B. Azevedo, R. van Zelm, F. Hayes, M.R. Ashmore, M.A.J. Huijbregts
Species Sensitivity Distributions offer opportunities in ozone risk assessment to both derive critical levels and estimate the affected fraction of a plant community.
- 7 Assessing urban habitat quality using spectral characteristics of *Tilia* leaves**
A.R. Khavanin Zadeh, F. Veroustraete, J.A.N. Buytaert, J. Dirckx, R. Samson
Leaf reflectance and dorsiventral properties of Linden tree species can be used to assess urban habitat quality.
- 15 Removal effectiveness and mechanisms of naphthalene and heavy metals from artificially contaminated soil by iron chelate-activated persulfate**
D.Y.S. Yan, I.M.C. Lo
This study focuses on the potencies and mechanisms of naphthalene and metal removal from contaminated soil by FeEDTA/FeEDDS-activated persulfate.
- 23 Fuzzy model for risk assessment of persistent organic pollutants in aquatic ecosystems**
X. Seguí, E. Pujolasus, S. Betrò, A. Àgueda, J. Casal, W. Ocampo-Duque, I. Rudolph, R. Barra, M. Páez, E. Barón, E. Eljarrat, D. Barceló, R.M. Darbra
The risk of POPs for aquatic organisms was assessed at several sites around the world, using a fuzzy-based model to provide useful results for decision-makers.
- 33 Anomalous concentrations of rare earth elements in the moss–soil system from south-central Poland**
S. Dołęgowska, Z.M. Migaszewski
The REE anomalies in different environmental sample matrices and methods of their interpretation were presented.
- 41 Integrated use of antioxidant enzymes and oxidative damage in two fish species to assess pollution in man-made hydroelectric reservoirs**
M.M. Sakuragui, M.G. Paulino, C.D.S. Pereira, C.S. Carvalho, H. Sadauskas-Henrique, M.N. Fernandes
In realistic environmental conditions, a mix of organochlorines and metals inhibited antioxidant defenses, causing oxidative stress in wild fish species.
- 52 Comparative effects of several cyclodextrins on the extraction of PAHs from an aged contaminated soil**
M.A. Sánchez-Trujillo, E. Morillo, J. Villaverde, S. Lacorte
Cyclodextrin solutions are useful and interesting tools for the decontamination of soils polluted by PAHs.
- 59 Sorption, dissolution and pH determine the long-term equilibration and toxicity of coated and uncoated ZnO nanoparticles in soil**
P.L. Waalewijn-Kool, M. Diez Ortiz, N.M. van Straalen, C.A.M. van Gestel
ZnO-NP addition causes Zn porewater concentrations to increase for more than one year, with higher toxicity for coated than for uncoated ZnO-NP.
- 65 Differential effects of metal contamination on the transcript expression of immune- and stress-response genes in the Sydney Rock oyster, *Saccostrea glomerata***
D.A. Taylor, E.L. Thompson, S.V. Nair, D.A. Raftos
qPCR identified significant down-regulation in stress- and immune-response genes in oysters exposed to metals, which could lead to increased disease susceptibility.
- 72 Developmental effects of aerosols and coal burning particles in zebrafish embryos**
A. Olivares, B.L. van Drooge, M. Casado, E. Prats, M. Serra, L.T. van der Ven, J.H. Kamstra, T. Hamers, S. Hermsen, J.O. Grimalt, B. Piña
Aerosols and coal burning particles showed a strong developmental toxicity in zebrafish, in a degree that cannot be directly predicted from chemical analyses or single-cell bioassays.

Continued on inside back cover

CONTENTS—Continued from outside back cover

- 80 Influence of land use on human bioaccessibility of metals in smelter-impacted soils**
A. Pelfrène, C. Waterlot, F. Douay
An empirical model, predicting the human bioaccessibility of metals in smelter-contaminated agricultural topsoils, was tested on other soil uses (urban and woody habitat topsoils).
- 89 The impact of aluminium smelter shut-down on the concentration of fluoride in vegetation and soils**
K.M. Brougham, S.R. Roberts, A.W. Davison, G.R. Port
Fluoride is hazardous to grazing herbivores, but when deposition stops, F-levels in plants and soil fall rapidly, some returning to background concentrations within a year.
- 97 Reclamation of copper-contaminated soil using EDTA or citric acid coupled with dissolved organic matter solution extracted from distillery sludge**
C.-C. Liu, Y.-C. Lin
Washing the soil with the two-phase chelate-DOM method could remove quantities of Cu and reserve most of fertility in soil.
- 102 Correlation analysis of the urban heat island effect and the spatial and temporal distribution of atmospheric particulates using TM images in Beijing**
L.Y. Xu, X.D. Xie, S. Li
Generally, if UHI is below 0.1 in the weak heat island or green island range, then DVI increases with the increase in UHI, and vice versa.
- 115 Associations of prenatal exposure to phenols with birth outcomes**
R. Tang, M.-j. Chen, G.-d. Ding, X.-j. Chen, X.-m. Han, K. Zhou, L.-m. Chen, Y.-k. Xia, Y. Tian, X.-r. Wang
Prenatal phenol exposure was sex-specifically related to birth outcomes.
- 121 Expression of zinc and cadmium responsive genes in leaves of willow (*Salix caprea* L.) genotypes with different accumulation characteristics**
C. Konlechner, M. Türkteş, I. Langer, M. Vaculík, W.W. Wenzel, M. Puschenreiter, M.-T. Hauser
*Expression analyzes reveal different responses to Cd and Zn exposure of *S. caprea* genotypes with different heavy metal accumulation abilities.*
- 128 Does plant uptake or low soil mineral-N production limit mineral-N losses to surface waters and groundwater from soils under grass in summer?**
A. Bhatti, C.J. McClean, M.S. Cresser
Low mineral-N production in soils under grass limits summer N losses to surface- and ground-waters.
- 135 Addition of contaminant bioavailability and species susceptibility to a sediment toxicity assessment: Application in an urban stream in China**
H. Li, B. Sun, X. Chen, M.J. Lydy, J. You
Field sediment toxicity caused by current-use pesticides could be more accurately evaluated by incorporating bioavailability measurements into the toxic unit analysis.
- 142 Maternal effects of inducible tolerance against the toxic cyanobacterium *Microcystis aeruginosa* in the grazer *Daphnia carinata***
X. Jiang, W. Yang, S. Zhao, H. Liang, Y. Zhao, L. Chen, R. Li
*Maternal exposure to the toxic *Microcystis aeruginosa* increased offspring fitness in one of three *Daphnia carinata* clones and carried a cost.*
- 147 Effects of insecticidal crystal proteins (Cry proteins) produced by genetically modified maize (Bt maize) on the nematode *Caenorhabditis elegans***
S. Höss, R. Menzel, F. Gessler, H.T. Nguyen, J.A. Jehle, W. Traunspurger
Insecticidal Cry proteins expressed by genetically modified maize act on nematodes via a similar mode of action as nematicidal Cry proteins, however, at concentrations far above expected soil levels.
- 152 Degradation of polycyclic aromatic hydrocarbons by microbial consortia enriched from three soils using two different culture media**
M. Wu, L. Chen, Y. Tian, Y. Ding, W.A. Dick
Bioremediation of oil-contaminated soils was most effective when using inoculum of microbial consortia from the same or similar soil as the soil being bioremediated.
- 159 The use of biodiesel blends on a non-road generator and its impacts on ozone formation potentials based on carbonyl emissions**
M. Chai, M. Lu, F. Liang, A. Tzillah, N. Dendramis, L. Watson
The study found that B50 resulted in lower total carbonyl emission rates and ozone formation potential resultant from these compounds, whereas both increased with B100.
- 166 Influence of different types of nanomaterials on their bioaccumulation in a paddy microcosm: A comparison of TiO₂ nanoparticles and nanotubes**
M.-K. Yeo, D.-H. Nam
TiO₂ nanomaterials can accumulate in the organisms of an artificial ecosystem.

- 419 Adaptive microbial population shifts in response to a continuous ethanol blend release increases biodegradation potential**
J. Ma, C.W. Nossa, Z. Xiu, W.G. Rixey, P.J.J. Alvarez
Pyrosequencing analysis discerned adaptive microbial population shifts that increase natural attenuation potential of an ethanol-blended fuel release.
- 426 Assessing the combined risks of PAHs and metals in urban soils by urbanization indicators**
C. Peng, Z. Ouyang, M. Wang, W. Chen, X. Li, J.C. Crittenden
The health risks of PAHs and HMs in residential soils were connected to building age, population density and road density of the community as well as its distance from urban center.
- 433 Spatial and temporal variations in pentachlorophenol dissipation at the aerobic–anaerobic interfaces of flooded paddy soils**
J. Lin, Y. Xu, P.C. Brookes, Y. He, J. Xu
The dissipation of PCP decreased with increasing soil depth in the surface layer, and varied between soils in the deep layer.
- 441 Validating the use of embryonic fish otoliths as recorders of sublethal exposure to copper in estuarine sediments**
N.C. Barbee, A. Greig, S.E. Swearer
We use sediments spiked with Cu to examine the uptake of heavy metals into embryonic fish otoliths as a novel tool for tracking pollution exposure history.
- 447 Environmental lead exposure risks associated with children’s outdoor playgrounds**
M.P. Taylor, D. Camenzuli, L.J. Kristensen, M. Forbes, S. Zahran
Playgrounds in Port Pirie are seriously contaminated by smelter emissions, with levels of surface dust and hand dust that pose a significant risk of harm to human health.
- 455 Dioxin and phthalate uptake and assimilation by the green mussel *Perna viridis***
W.-X. Wang, Q. Zhang
Aqueous uptake was the predominant pathway for dioxin and phthalate accumulation in marine green mussels.
- 463 Urban metabolism: A review of research methodologies**
Y. Zhang
The author reviews research progress in the field of urban metabolism, and based on her literature review, proposes directions for future research.
- 474 Update on the cometabolism of organic pollutants by bacteria**
A. Nzila
Cometabolism, the use of microorganisms to biodegrade pollutants without using them as growth-substrates, provides an effective approach to remove recalcitrant pollutants.
- 483 The physical impacts of microplastics on marine organisms: A review**
S.L. Wright, R.C. Thompson, T.S. Galloway
Here we review microplastic uptake in marine organisms and assess individual, population and community level effects, highlighting the most susceptible species.
- 493 Maternally transferred mercury in wild largemouth bass, *Micropterus salmoides***
D.K. Sackett, D.D. Aday, J.A. Rice, W.G. Cope
Here we report higher maternal transfer and higher ratios of egg to muscle tissue Hg than previously observed for teleost fish.
- 498 Silver nanoparticle toxicity in sea urchin *Paracentrotus lividus***
L. Šiller, M.-L. Lemloh, S. Piticharoenphun, B.G. Mendis, B.R. Horrocks, F. Brümmer, D. Medaković
Silver nanoparticles cause dose dependent developmental defects in sea urchin and they are more toxic than their equivalent Ag⁺ ion dose.