

## BEHAVIOUR, WELFARE &amp; ENVIRONMENT

Traditional and commercial turkeys show similar susceptibility to foot pad dermatitis and behavioural evidence of pain. *P.M. Hocking and K. Wu.* 281

Volatile gas concentrations in turkey houses estimated by Fourier Transform Infrared Spectroscopy (FTIR). *D. Witkowska.* 289

Identifying risk factors for eggshell contamination by *Bacillus cereus* group bacteria in French laying farms. *A.Z. Kone, S. Jan, C. Le Marechal, N. Grosset, M. Gautier, J. Puterflam and F. Baron.* 298

## GENETICS

The relationship of plumage colours with MC1R (Melanocortin 1 Receptor) and ASIP (Agouti Signaling Protein) in Japanese quail (*Coturnix coturnix japonica*). *X.H. Zhang, Y.Z. Pang, S.J. Zhao, H.W. Xu, Y.L. Li, Y. Xu, Z. Guo and D.D. Wang.* 306

A single-nucleotide polymorphism in the 3' untranslated region of the *LPIN1* gene and association analysis with performance traits in chicken. *S.P. Zhang, S.Y. Li, W. Chen, W.W. Lu and Y.Q. Huang.* 312

## IMMUNOLOGY, HEALTH &amp; DISEASE

Potential of a commercially available water acidification product for reducing *Campylobacter* in broilers prior to slaughter. *P.N. Haughton, J. Lyng, S. Fanning and P. Whyte.* 319

Identification of antibodies against hydropericardium syndrome in wild birds. *S. Manzoor, Z. Hussain, S.U. Rahman and I. Hussain.* 325

## MEAT &amp; EGG SCIENCE

Carcass characteristics, composition, physico-chemical, microbial and sensory quality of emu meat. *B.M. Naveena, A.R. Sen, M. Muthukumar, P.S. Girish, Y. Praveen Kumar and M. Kiran.* 329

## NUTRITION &amp; METABOLISM

Influence of pellet diameter and length on the quality of pellets and performance, nutrient utilisation and digestive tract development of broilers fed on wheat-based diets. *M.R. Abdollahi, V. Ravindran, T.J. Wester, G. Ravindran and D.V. Thomas.* 337

Effects of exogenous xylanase on performance, nutrient digestibility and caecal thermal profiles of broilers given wheat-based diets. *A.J. Cowieson and H.V. Masey O'Neill.* 346

Apilarnil reduces fear and advances sexual development in male broilers but has no effect on growth. *Ö. Altan, B. Yücel, Z. Açıkgöz, Ç. Seremet, M. Kösoğlu, N. Turgan and A.M. Özgönül.* 355

Effect of two microbial phytases on mineral availability and retention and bone mineral density in low-phosphorus diets for broilers. *T.K. Chung, S.M. Rutherford, D.V. Thomas and P.J. Moughan.* 362

The influence of diet composition on egg and chick traits in captive Greater Rhea females. *M.C. Lábaque, M.B. Martella, D.M. Maestri and J.L. Navarro.* 374

Effect of intravenously injected zinc on tissue zinc and metallothionein gene expression of broilers. *S.F. Shen, R.L. Wang, L. Lu, S.F. Li, S.B. Liu, J.J. Xie, L.Y. Zhang, M.L. Wang and X.G. Luo.* 381

## PHARMACOLOGY

Pharmacokinetics of enrofloxacin after multiple subcutaneous and intramuscular administrations in adult ostriches. *J.J. De Lucas, J. Solano, F. González, C. Ballesteros, M.I. San Andrés, C. Martín Von Kauffmann and C. Rodríguez.* 391

## PHYSIOLOGY, ENDOCRINOLOGY &amp; REPRODUCTION

Expression of TRPV6 and CaBP-D28k in the egg shell gland (uterus) during the oviposition cycle of the laying hen. *J.H. Yang, Z.H. Zhao, J.F. Hou, Z.L. Zhou, Y.F. Deng and J.J. Dai.* 398

## SHORT COMMUNICATION

Riboflavin requirements of white Pekin ducks from hatch to 21 d of age. *J. Tang, M. Xie, J. Yang, Z.G. Wen, Y.W. Zhu, W. Huang and S.S. Hou.* 407