

Removal of intensive agriculture from the landscape improves aquatic ecosystem health	
B. Gagliardi and V. Pettigrove (Australia)	1
Bird diversity relates to agri-environment schemes at local and landscape level in intensive farmland	
M. Hiron, Å. Berg, S. Eggers, J. Josefsson and T. Pärt (Sweden)	9
Forecasting the potential impacts of CAP-associated land use changes on farmland birds at the national level	
F. Chiron, K. Princé, M.L. Paracchini, C. Bulgheroni and F. Jiguet (France, USA, Italy)	17
Spatial-temporal dynamics of water soluble phosphorus in the topsoil of a low mountain range catchment	
I. Kistner, G. Ollesch, R. Meissner and M. Rode (Germany)	24
Amount, distribution and driving factors of soil organic carbon and nitrogen in cropland and grassland soils of southeast Germany (Bavaria)	
M. Wiesmeier, R. Hübner, F. Barthold, P. Spörlein, Uwe Geuß, E. Hangen, A. Reischl, B. Schilling, M. von Lützwow and I. Kögel-Knabner (Germany)	39
Effect of fish species on methane and nitrous oxide emission in relation to soil C, N pools and enzymatic activities in rainfed shallow lowland rice-fish farming system	
P. Bhattacharyya, D.P. Sinhababu, K.S. Roy, P.K. Dash, P.K. Sahu, R. Dandapat, S. Neogi and S. Mohanty (India)	53
Response of ground beetle (Coleoptera, Carabidae) communities to changes in agricultural policies in Sweden over two decades	
A. Rusch, R. Bommarco, P. Chiverton, Sandra Öberg, H. Wallin, S. Wikteliu and B. Ekbom (Sweden, France, Norway)	63
A comparative study of the seed banks of abandoned paddy fields along a chronosequence in Japan	
S. Yamada, Y. Kitagawa and S. Okubo (Japan)	70