

Regular papers

Factors biasing the correlation structure of patch level landscape metrics S. Szabó, Z. Túri and S. Márton (Debrecen, Hungary).....	1
Ecosystem service state and trends at the regional to national level: A rapid assessment J. Helfenstein (Zurich, Switzerland) and F. Kienast (Birmensdorf, Switzerland).....	11
Towards a unique landscape description for multi-species studies: A model comparison with common birds in a human-dominated French region A. Mimet, N. Maurel, V. Pellissier, L. Simon and R. Julliard (Paris, France)	19
The effect of size, weight, body compartment, sex and reproductive status on the bioaccumulation of 19 trace elements in rope-grown <i>Mytilus galloprovincialis</i> J. Richir and S. Gobert (Liège, Belgium)	33
Modeling Canada yew (<i>Taxus canadensis</i> Marsh.) distribution and abundance in the boreal forest of northeastern Ontario, Canada J.P. Dech, S. Mayhew-Hammond, A.L. James and B. Pokharel (North Bay, Canada)	48
The environmental efficiency of Wanjiang demonstration area: A Bayesian estimation approach M. Song and Y. Guan (Bengbu, China)	59
An application of Social Values for Ecosystem Services (SOLVES) to three national forests in Colorado and Wyoming B.C. Sherrouse (Baltimore, MD, USA), D.J. Semmens (Denver, CO, USA) and J.M. Clement (Laramie, WY, USA).....	68
Landscape water potential as a new indicator for monitoring macrostructural landscape changes J. Skaloš, K. Berchová, J. Pokorný, T. Sedmidubský, E. Pecharová and I. Trpáková (Prague 6, Czech Republic).....	80
An improved approach for modeling spatial distribution of water use profit—A case study in Tuhai Majia Basin, China Y. Huang, D. Jiang, D. Zhuang, Y. Zhu and J. Fu (Beijing, China).....	94
The Theorized Urban Gradient (TUG) method—A conceptual framework for socio-ecological sampling in complex urban agglomerations S. Qureshi (Berlin, Germany and Birmingham, UK), D. Haase (Berlin, Germany) and R. Coles (Birmingham, UK)	100
A comparison of methods for estimating common vole (<i>Microtus arvalis</i>) abundance in agricultural habitats D. Jareño, J. Viñuela (Ciudad Real, Spain), J.J. Luque-Larena (Palencia, Spain), L. Arroyo, B. Arroyo (Ciudad Real, Spain) and F. Mougeot (Ciudad Real, Spain and Almería, Spain).....	111
Evaluating the ecological effects of social agent scenarios for a housing development in the Atlantic Forest E. Hardt, R.F. dos Santos and E.F.L. Pereira-Silva (São Paulo, Brazil).....	120
Quantifying and mapping ecosystem services: Demand and supply of pollination in the European Union C.J.E. Schulp (Amsterdam, The Netherlands), S. Lautenbach (Bonn, Germany) and P.H. Verburg (Amsterdam, The Netherlands).....	131
Evaluating performance of macroinvertebrate-based adjusted and unadjusted multi-metric indices (MMI) using multi-season and multi-year samples K. Chen (Nanjing, People's Republic of China), R.M. Hughes (Corvallis, OR, USA), S. Xu, J. Zhang (Nanjing, People's Republic of China), D. Cai (Yichang, People's Republic of China) and B. Wang (Nanjing, People's Republic of China)	142

Standing herbage mass: An integrated indicator of management practices for examining how fertility and defoliation regime shape the functional structure of species-rich grasslands M. Duru, P. Cruz, P. Ansquer (Castanet Tolosan, France) and M.L. Navas (Montpellier, France)	152
Forest fragmentation and landscape connectivity change associated with road network extension and city expansion: A case study in the Lancang River Valley S. Liu, Y. Dong, L. Deng, Q. Liu, H. Zhao and S. Dong (Beijing, China)	160
The application of remote sensing for marine protected area management D. Kachelriess, M. Wegmann (Germany), M. Gollock and N. Pettorelli (London, UK)	169
A comparison of survey methods to evaluate macrophyte index of biotic integrity performance in Minnesota lakes B. Vondracek, J.D. Koch and M.W. Beck (St. Paul, MN, USA)	178
Taxonomic surrogacy, numerical resolution and responses of stream macroinvertebrate communities to ecological gradients: Are the inferences transferable among regions? J. Heino (Finland)	186
Leaf decomposition and ecosystem metabolism as functional indicators of land use impacts on tropical streams E.F. Silva-Junior (São João del-Rei, Brazil), T.P. Moulton (Rio de Janeiro, Brazil), I.G. Boëchat and B. Gücker (São João del-Rei, Brazil)	195
The use of environmental performance indicators and size effect: A study of industrial companies F. Hourneaux Jr. (São Paulo, Brazil), H.A. Hrdlicka (Rio Tinto, Brazil), C.M. Gomes (Santa Maria, Brazil) and I. Kruglianskas (São Paulo, Brazil)	205
Measuring progress towards sustainable intensification: How far can secondary data go? A.P. Barnes and S.G. Thomson (Edinburgh, UK)	213
Land availability vs conversion by use type: A new approach for land take monitoring L. Salvati (Rome, Italy)	221
Using stable nitrogen isotopes in <i>Patella</i> sp. to trace sewage-derived material in coastal ecosystems P.Ž. Rožič, T. Dolenc, S. Lojen (Ljubljana, Slovenia), G. Kniewald (Zagreb, Croatia) and M. Dolenc (Ljubljana, Slovenia)	224
Stand-total tree-ring measurements and forest inventory documented climate-induced forest dynamics in the semi-arid Altai Mountains X. Wu, H. Liu, L. He, Z. Qi (Beijing, China), O.A. Anenkhonov (Ulan Ude, Russia), A.Yu. Korolyuk (Novosibirsk, Russia), Y. Yu and D. Guo (Beijing, China)	231
Dimension reduction and data sharpening of high-dimensional vegetation data: An application to Swiss mire monitoring S. Ghosh, U. Graf, K. Ecker, O. Wildi, H. Kuchler, E. Feldmeyer-Christe and M. Kuchler (Birmensdorf, Switzerland)	242
Agricultural non-point source pollution in the Yongding River Basin W. Guo (Zhengzhou, P.R. China and Beijing, P.R. China), Y. Fu, B. Ruan, H. Ge and N. Zhao (Beijing, P.R. China)	254
Using <i>Pinus uncinata</i> to monitor tropospheric ozone in the Pyrenees S.C. Kefauver (Davis, CA, USA and Barcelona, Spain), J. Peñuelas, A. Ribas, M. Díaz-de-Quijano (Barcelona, Spain) and S. Ustin (Davis, CA, USA)	262
Morphological abnormalities in periphytic diatoms as a tool for biomonitoring of heavy metal pollution in a river L.K. Pandey (Varanasi, India), D. Kumar (Srinagar, India), A. Yadav, J. Rai and J.P. Gaur (Varanasi, India)	272
Development of a 'sediment-stress' functional-level indicator for the seagrass <i>Halophila ovalis</i> K. Kilminster, V. Forbes (Perth, Australia) and M. Holmer (Odense M, Denmark)	280
Harmonization of the Land Cover Classification System (LCCS) with the General Habitat Categories (GHC) classification system V. Kosmidou, Z. Petrou (Thessaloniki, Greece), R.G.H. Bunce (Tartu, Estonia), C.A. Múcher, R.H.G. Jongman, M.M.B. Bogers (Wageningen, The Netherlands), R.M. Lucas (Wales, UK), V. Tomaselli, P. Blonda (Bari, Italy), E. Padoa-Schioppa (Milano, Italy), I. Manakos and M. Petrou (Thessaloniki, Greece)	290
Estimation of soil organic matter by geostatistical methods: Use of auxiliary information in agricultural and environmental assessment C. Piccini, A. Marchetti and R. Francaviglia (Rome, Italy)	301
Urban sprawl scatterplots for Urban Morphological Zones data L. Altieri, D. Cocchi, G. Pezzi (Bologna, Italy), E.M. Scott (Glasgow, UK) and M. Ventrucci (Bologna, Italy)	315
Long term variation of an amphipod species' population secondary production as indicator of incomplete resilience in a temperate estuary N. Leite, A. Guerra, A. Almeida, J.C. Marques and I. Martins (Coimbra, Portugal)	324
Physico-chemical characterization of street dust and re-suspended dust on plant canopies: An approach for finger printing the urban environment S.S. Ram (Kolkata, India and Cambridge, UK), R.V. Kumar (Cambridge, UK), P. Chaudhuri, S. Chanda (Kolkata, India), S.C. Santra (Nadia, India), M. Sudarshan and A. Chakraborty (Kolkata, India)	334
Isotopic metrics as a tool for assessing the effects of mine pollution on stream food webs K.L. Hogsden and J.S. Harding (Christchurch, New Zealand)	339
A "lost biodiversity" approach to revealing major anthropogenic threats to regional freshwater ecosystems A. Yoshioka (Bunkyo-ku, Japan), Y. Miyazaki (Odawara-shi, Japan), Y. Sekizaki (Chiyoda-ku, Japan), S.-i. Suda (Bunkyo-ku, Japan), T. Kadoya (Tsukuba, Japan) and I. Washitani (Bunkyo-ku, Japan)	348

Macrofaunal community abundance and diversity and talitrid orientation as potential indicators of ecological long-term effects of a sand-dune recovery intervention D.H. Nourisson (Firenze, Italy), F. Bessa (Coimbra, Portugal), F. Scapini (Firenze, Italy) and J.C. Marques (Coimbra, Portugal)	356
Ranking matrices as operational tools for the environmental risk assessment of genetically modified crops on non-target organisms A. Hilbeck, G. Weiss (Rängsdorf, Germany), B. Oehen (Rängsdorf, Germany and Frick, Switzerland), J. Römbke, S. Jänsch (Flörsheim, Germany), H. Teichmann (Bonn, Germany), A. Lang (Basel, Switzerland), M. Otto and B. Tappeser (Bonn, Germany)	367
Indicators of responsible investing B. Scholtens (Groningen, Netherlands and Scotland, UK)	382
Detecting anthropogenic effects on a vulnerable species, the freshwater blenny (<i>Salaria fluviatilis</i>): The importance of considering key ecological variables M. Laporte (Montpellier, France and Trois-Rivières, Canada), A. Bertolo (Trois-Rivières, Canada), P. Berrebi (Montpellier, France) and P. Magnan (Trois-Rivières, Canada)	386
Evaluating relationships between mercury concentrations in air and in Spanish moss (<i>Tillandsia usneoides</i> L.) K.T. Sutton, R.A. Cohen and S.P. Vives (Statesboro, GA, USA)	392
Wetland and landscape indices for assessing the condition of semiarid Mediterranean saline wetlands under agricultural hydrological pressures J. Martínez-López, M.F. Carreño, J. Martínez-Fernández and M.A. Esteve (Murcia, Spain)	400
Measuring basal soil respiration across Europe: Do incubation temperature and incubation period matter? R.E. Creamer, R.P.O. Schulte, D. Stone (Johnstown Castle, Ireland), A. Gal (Gödöllő, Hungary), P.H. Krogh (Denmark), G. Lo Papa (Italy), P.J. Murray (Devon, UK), G. Pérès (France), B. Foerster (Germany), M. Rutgers (The Netherlands), J.P. Sousa (Portugal) and A. Winding (Denmark)	409
Energy evaluation vs. life cycle-based embodied energy (solar, tidal and geothermal) of wood biomass resources E. Neri (Siena, Italy), B. Rugani, E. Benetto (Esch-sur-Alzette, Luxembourg) and S. Bastianoni (Siena, Italy)	419
Windborne: Can liverworts be used as indicators of altitudinal gradient in the Brazilian Atlantic Forest? N.D. dos Santos (Campinas, Brazil), D.P. da Costa (Rio de Janeiro, Brazil), L.S. Kinoshita and G.J. Shepherd (Campinas, Brazil)	431
Using statistical tests on relative ecological indicator values to compare vegetation units – Different approaches and weighting methods C. Tölgyesi, Z. Bátori and L. Erdős (Szeged, Hungary)	441
Using benthic indicators to assess status at HUC levels: A robust hierarchical modeling approach C. Velasco-Cruz and E.P. Smith (Blacksburg, VA, USA)	447
Aggregated indices for trends in eutrophication of different types of fresh water in the Netherlands P.J.T.M. van Puijenbroek (Bilthoven, Netherlands), P. Cleij (Utrecht, Netherlands) and H. Visser (Bilthoven, Netherlands)	456
Factors explaining the patterns of benthic chlorophyll- <i>a</i> distribution in a large agricultural Iberian watershed (Gudiana river) G. Urrea-Clos, E. García-Berthou and S. Sabater (Girona, Spain)	463
Enhancing the performance and interpretation of freshwater biological indices: An application in arid zone streams J.J. Vander Laan and C.P. Hawkins (Logan, UT, USA)	470
Embryo development of the benthic amphipod <i>Monoporeia affinis</i> as a tool for monitoring and assessment of biological effects of contaminants in the field: A meta-analysis M. Reutgard, A.-K. Eriksson Wiklund, M. Breitholtz and B. Sundelin (Stockholm, Sweden)	483
Lead adsorption capacity of some moss species used for heavy metal analysis B. Koz (Giresun, Turkey) and U. Cevik (Trabzon, Turkey)	491
Analysis of spatial scales for ecosystem services: Application of the lacunarity concept at landscape level in Galicia (NW Spain) J.V. Rocas-Díaz, E.R. Díaz-Varela and P. Álvarez-Álvarez (Spain)	495
Theoretical exploration for the combination of the ecological, energy, carbon, and water footprints: Overview of a footprint family K. Fang, R. Heijungs and G.R. de Snoo (Leiden, The Netherlands)	508
Simplicity is the ultimate sophistication: Building capacity to meet the challenges of the Water Framework Directive M. Kelly (Bowburn, UK)	519

Convergence between ANPP estimation methods in grasslands — A practical solution to the comparability dilemma	
J.C. Ruppert (Cologne, Germany and Bonn, Germany) and A. Linstädter (Cologne, Germany)	524
Combined toxicological effects of pesticides: A fish multi-biomarker approach	
C. Bacchetta, A. Rossi, A. Ale, M. Campana, M.J. Parma and J. Cazenave (Santa Fe, Argentina)	532
Accounting for capacity and flow of ecosystem services: A conceptual model and a case study for Telemark, Norway	
M. Schröter (Wageningen, The Netherlands), D.N. Barton (Oslo, Norway), R.P. Remme and L. Hein (Wageningen, The Netherlands)	539
Spatial assessment and mapping of biodiversity and conservation priorities in a heavily modified and fragmented production landscape in north-central Victoria, Australia	
H. Baral (Richmond, Australia), R.J. Keenan (Parkville, Australia), S.K. Sharma (Adelaide, Australia), N.E. Stork (Nathan, Australia) and S. Kasel (Richmond, Australia)	552
Mapping correlations between nitrogen concentrations in atmospheric deposition and mosses for natural landscapes in Europe	
W. Schröder, R. Pesch, S. Schönrock (Vechta, Germany), H. Harmens, G. Mills (Bangor, UK) and H. Fagerli (Oslo, Norway)	563
Biotic congruence in humid tropical forests: A multi-taxa examination of spatial distribution and responses to forest disturbance	
R.B. de Andrade (USA), J. Barlow (UK), J. Louzada, L. Mestre, J. Silveira, F.Z. Vaz-de-Mello (Brazil) and M.A. Cochrane (USA)	572
Social indicators are predictors of airborne outdoor exposures in Berlin	
U. Franck (Leipzig, Germany), H.-J. Klimeczek (Berlin, Germany) and A. Kindler (Leipzig, Germany)	582
Spatial distribution of wild boar population abundance: Basic information for spatial epidemiology and wildlife management	
P. Acevedo (Vairão, Portugal, Bellaterra, Spain and Ciudad Real, Spain), F. Quirós-Fernández (Ciudad Real, Spain), J. Casal (Bellaterra, Spain) and J. Vicente (Ciudad Real, Spain)	594
Role of thermal niche in the cellular response to thermal stress: Lipid peroxidation and HSP70 expression in coastal crabs	
D. Madeira (Lisboa, Portugal and Caparica, Portugal), L. Narciso (Cascais, Portugal), H.N. Cabral (Lisboa, Portugal), M.S. Diniz (Caparica, Portugal) and C. Vinagre (Lisboa, Portugal)	601
The temperature recorded by simulated mobile receptors is an indicator for the thermal exposure of the urban inhabitants	
U. Schlink, A. Kindler, K. Großmann, N. Schwarz and U. Franck (Leipzig, Germany)	607
Effectiveness of entropy-based functions in the analysis of ecosystem state and development	
A. Ludovisi (Perugia, Italy)	617
A multi-scale method to assess pesticide contamination risks in agricultural watersheds	
F. Macary, S. Morin (Cestas, France), J.-L. Probst (Castanet-Tolosan, France) and F. Saudubray (Cestas Cedex, France)	624
On the feasibility of a consumer-based allocation method in national GHG inventories	
D. Caro, S. Bastianoni, S. Borghesi and F.M. Pulselli (Siena, Italy)	640
Linking biodiversity indicators, ecosystem functioning, provision of services and human well-being in estuarine systems: Application of a conceptual framework	
R. Pinto (Coimbra, Portugal), V.N. de Jonge (Hull, UK) and J.C. Marques (Coimbra, Portugal)	644
Key features for saproxylic beetle diversity derived from rapid habitat assessment in temperate forests	
C. Bouget (Nogent-sur-Vernisson, France), L. Larrieu (Castanet-Tolosan, France and Auzerville Tolosane, France) and A. Brin (Toulouse, France)	656
Using eco-efficiency as an indicator for sustainable urban development: A case study of Chinese provincial capital cities	
K. Yin, R. Wang (Beijing, China), Q. An (Hefei, China), L. Yao (Beijing, China) and J. Liang (Changsha, China)	665
Parameters selection for water quality index in the assessment of the environmental impacts of land-based trout farms	
M.A.T. Koçer and H. Sevgili (Antalya, Turkey)	672
Identification of key assessment indicators of the zero waste management systems	
A.U. Zaman (Adelaide, Australia)	682
Niche segregation amongst sympatric species at exposed sandy shores with contrasting wrack availabilities illustrated by stable isotopic analysis	
F. Bessa, A. Baeta and J.C. Marques (Coimbra, Portugal)	694
Dung beetle communities as biological indicators of riparian forest widths in southern Brazil	
G. Viegas, C. Stenert, U.H. Schulz and L. Maltchik (São Leopoldo, Brazil)	703

The assessment of local sustainability using fuzzy logic: An expert opinion system to evaluate environmental sanitation in the Algarve region, Portugal D. Canavese (Matinhos, Brazil), N.R.S. Ortega (São Paulo, Brazil) and M. Queirós (Lisbon, Portugal).....	711
Live benthic foraminiferal faunas from the French Mediterranean Coast: Towards a new biotic index of environmental quality C. Barras, F.J. Jorissen (Angers Cedex, France), C. Labrune (Banyuls/Mer, France), B. Andral (La Seyne-sur-Mer Cedex, France) and P. Boissery (Lyon, France).....	719
Corrigendum to “A new stochastic dynamic tool to improve the accuracy of mortality estimates for bats killed at wind farms” [Ecol. Indicators 34 (2013) 428–440] R. Bastos, M. Santos and J.A. Cabral (Vila Real, Portugal)	744
Corrigendum to “Geographic position of sample grid but not the removal of uncommon species affect multivariate analyses of diverse assemblages. The case of oribatid mites (Acari: Oribatida)” [Ecol. Indicat. 34 (2013) 172–180] E. Franklin (Manaus, Brazil), J. de Moraes (Brazil), V.L. Landeiro (Cuiabá, Brazil), J.L.P. de Souza, P.A.C.L. Pequeno (Brazil), W.E. Magnusson and J.W. de Moraes (Manaus, Brazil)	745
Special Section: Assessment of rural livelihoods in South-West China based on environmental, economic, and social indicators	
Assessment of rural livelihoods in South-West China based on environmental, economic, and social indicators S. Herrmann (Hannover, Germany) and J.M. Fox (Honolulu, HI, USA).....	746
Landscape transformation through the use of ecological and socioeconomic indicators in Xishuangbanna, Southwest China, Mekong Region J. Xu, R.E. Grumbine (Kunming, China) and P. Beckschäfer (Göttingen, Germany)	749
CLUENaban-A land use change model combining social factors with physical landscape factors for a mountainous area in Southwest China S. Wehner (Passau, Germany), S. Herrmann and K. Berkhoff (Hannover, Germany).....	757
Farm types as an interface between an agroeconomical model and CLUE-Naban land change model: Application for scenario modelling T.M. Gibreel (Gießen, Germany), S. Herrmann, K. Berkhoff (Hannover, Germany), E.-A. Nuppenau and A. Rinn (Gießen, Germany)	766
Designing a sustainable land use scenario based on a combination of ecological assessments and economic optimization M. Cotter (Stuttgart, Germany), K. Berkhoff (Hannover, Germany), T. Gibreel (Gießen, Germany), A. Ghorbani, R. Golbon (Stuttgart, Germany), E.-A. Nuppenau (Gießen, Germany) and J. Sauerborn (Stuttgart, Germany)	779
Developing indicators of economic value and biodiversity loss for rubber plantations in Xishuangbanna, southwest China: A case study from Menglun township Z.-F. Yi (Yunnan, China, Beijing, China and Kunming, China), C.H. Cannon (Yunnan, China and Lubbock, TX, USA), J. Chen, C.-X. Ye (Yunnan, China) and R.D. Swetnam (Cambridge, UK)	788