

# CONTENTS

Seasonality of above-ground net primary productivity along an Andean altitudinal transect in Peru <b>C. A. J. Girardin, Y. Malhi, K. J. Feeley, J. M. Rapp, M. R. Silman, P. Meir, W. Huaraca Huasco, N. Salinas, M. Mamani, J. E. Silva-Espejo, K. García Cabrera, W. Farfan Rios, D. B. Metcalfe, C. E. Doughty and L. E. O. C. Aragão</b>	503
Frugivorous weevils are too rare to cause Janzen–Connell effects in New Guinea lowland rain forest <b>Richard Cvrtecka, Katerina Sam, Erik Brus, George D. Weiblen and Vojtech Novotny</b>	521
Spatial and temporal variation in amphibian metacommunity structure in Chiapas, Mexico <b>Leticia Margarita Ochoa-Ochoa and Robert James Whittaker</b>	537
The association between invasive <i>Lantana camara</i> and seedlings/saplings of a plant community in Mudumalai Tiger Reserve, India <b>Gaius Wilson, Ajay A. Desai, Dalice A. Sim, Monica A. M. Gruber and Philip J. Lester</b>	551
Influences of forest structure and landscape features on spatial variation in species composition in a palm community in central Amazonia <b>Lilian Figueiredo Rodrigues, Renato Cintra, Carolina Volkmer Castilho, Ocírio de Sousa Pereira and Tânia P. Pimentel</b>	565
Effects of patch size on liana diversity and distributions in the tropical montane evergreen forests of the Nilgiri Mountains, southern India <b>Dharmalingam Mohandass, Alice C. Hughes, Mason Campbell and Priya Davidar</b>	579
Termitaria as preferred browsing patches for black rhinoceros ( <i>Diceros bicornis</i> ) in Chipinge Safari Area, Zimbabwe <b>Justice Muvengwi, Hilton G. T. Ndagurwa, Tatenda Nyenda and Innocent Mlambo</b>	591
Arbuscular mycorrhizas increase survival, precocity and flowering of herbaceous and shrubby species of early stages of tropical succession in pot cultivation <b>Artur Berbel Lirio Rondina, Luis Eduardo Azevedo Marques Lescano, Ricardo de Almeida Alves, Enio Massao Matsuura, Marco Antonio Nogueira and Waldemar Zangaro</b>	599
<i>Short Communications</i>	
Rapid collapse of a population of <i>Dieffenbachia</i> spp., plants used for tadpole-rearing by a poison-dart frog ( <i>Oophaga pumilio</i> ) in a Costa Rican rain forest <b>Mark J. McKone, Jonathan W. Moore, Christopher W. Harbison, Ian C. Holmen, Hillary C. Lyons, Kristine M. Nachbor, Julia L. Michalak, Maurine Neiman, Julia L. Nicol and George R. Wheeler</b>	615
Consumption of lipid-rich seed arils improves larval development in a Neotropical primarily carnivorous ant, <i>Odontomachus chelifer</i> (Ponerinae) <b>Claudia Bottcher and Paulo S. Oliveira</b>	621
Epiphyte assemblages respond to host life-form independently of variation in microclimate in lower montane cloud forest in Panama <b>Jennifer C. Sanger and James B. Kirkpatrick</b>	625